

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number  
WO 01/83749 A2

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
C07K 14/705, G01N 33/68, C12Q 1/68, C07K 16/28,  
A61K 38/17, A01K 67/027 // A61P 3/04, 3/10, 25/32

(21) International Application Number: PCT/US01/13387

(22) International Filing Date: 25 April 2001 (25.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/200,794 28 April 2000 (28.04.2000) US  
60/221,419 28 July 2000 (28.07.2000) US  
60/247,443 10 November 2000 (10.11.2000) US

(71) Applicants (for all designated States except US):  
WARNER-LAMBERT COMPANY [US/US]; 201 Tabor  
Road, Morris Plains, NJ 07950 (US). THE MONELL  
CHEMICAL SENSES CENTER [US/US]; 3500 Market  
Street, Philadelphia, PA 19104 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BACHMANOV,  
Alexander, A. [RU/US]; The Monell Chemical Senses  
Center, 3500 Market Street, Philadelphia, PA 19104 (US).  
BEAUCHAMP, Gary, K. [US/US]; The Monell Chemical  
Senses Center, 3500 Market Street, Philadelphia, PA 19104  
(US). CHATTERJEE, Aurobindo [IN/US]; 2106 Clinton  
Avenue, Apt. A, Alameda, CA 94501 (US). DE JONG,  
Pieter, J. [US/US]; 631 Catamaran Street #4, Foster  
City, CA 94404 (US). LI, Shanru [CN/US]; The Monell  
Chemical Senses Center, 3500 Market Street, Philadelphia,  
PA 19104 (US). Li, Xia [CN/US]; The Monell Chemical

Senses Center, 3500 Market Street, Philadelphia, PA 19104  
(US). OHMEN, Jeffrey, D. [US/US]; 1817 Versailles  
Avenue, Alameda, CA 94501 (US). REED, Danielle,  
R. [US/US]; The Monell Chemical Senses Center, 3500  
Market Street, Philadelphia, PA 19104 (US). ROSS, David  
[US/US]; 2651 Barbers Point Road, Apt. D., Alameda,  
CA 94501 (US). TORDOFF, Michael, Guy [GB/US];  
The Monell Chemical Senses Center, 3500 Market Street,  
Philadelphia, PA 19104 (US).

(74) Agents: FEDERMAN, Evan, J.; Warner-Lambert Com-  
pany, 201 Tabor Road, Morris Plains, NJ 07950 et al. (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 01/83749 A2

(54) Title: GENE AND SEQUENCE VARIATION ASSOCIATED WITH SENSING CARBOHYDRATE COMPOUNDS AND  
OTHER SWEETENERS

(57) Abstract: The present invention relates to the discovery of a gene and its sequence variation associated with preference for carbohydrates, other sweet compounds, or ethanol. The present invention also relates to the study of metabolic pathways to identify other genes, receptors, and relationships that contribute to differences in sensing of carbohydrates or ethanol. The present invention also relates to germline or somatic sequence variations and its use in the diagnosis and prognosis of predisposition to diabetes, other obesity related disorders, or ethanol consumption. The present invention also provided probes or primers specific for the detection and analysis of such sequence variation. The present invention also relates to method for screening drugs for inhibition or restoration of gene function as antidiabetic, antiobesity, or antialcohol consumption therapies. The present invention relates to other antidiabetic, antiobesity disorder, or antialcohol consumption therapies, such as gene therapy, protein replacement therapy, etc. Finally, the present invention relates to a method for identifying sweeteners or alcohols utilizing the gene and its variations.

-1-

## GENE AND SEQUENCE VARIATION ASSOCIATED WITH SENSING CARBOHYDRATE COMPOUNDS AND OTHER SWEETENERS

### FIELD OF THE INVENTION

The present invention relates generally to the field of mouse and human genetics and sensing of extracellular carbohydrates. Specifically, the present invention relates to the discovery of a gene and its sequence variation associated  
5 with a differential preference for sweet compounds in laboratory strains of mice.

### BACKGROUND OF THE INVENTION

The ability to sense extra-cellular carbohydrates, transduce this sensory information, and relay it to the brain, is carried out by membrane bound receptors in taste papillae. Many approaches to identify the sweet receptor or receptors have  
10 been tried, but the problem has proved, until recently, to be difficult.

Mammals vary in their *ad libitum* consumption of sweeteners. To investigate the genetic contribution to this complex behavior, behavioral, electrophysiological, and genetic studies were conducted using two strains of mice that differ markedly in their preference for sucrose and saccharin (Bachmanov  
15 et al., *Behavior Genetics*, 1996;26:563-573).

Recently published data indicates that the ability to sense carbohydrates is linked to obesity. These studies demonstrated that sensation of simple carbohydrates is suppressible by the adipose hormone, leptin.

These studies demonstrated that a locus on the telomere of mouse  
20 chromosome 4 accounts for ~40% of the genetic variability in sucrose and saccharin intake, and that the effect of this locus is to enhance or retard the gustatory neural response to sucrose.

## SUMMARY OF THE INVENTION

The present invention provides a gene and its sequence variation associated with a preference for carbohydrate compounds, other sweeteners, or alcohol.

5 The present invention provides a gene and its sequence variation associated a differential response by the pancreas and/or muscle in response to dietary carbohydrates.

10 The present invention also relates to sequence variation and its use in the diagnosis and prognosis of predisposition to diabetes, other obesity-related disorders, or alcohol consumption.

15 The present invention also relates to the study of taste to identify molecules responsible for signal transduction, other receptors and genes and relationships that contribute to taste preference.

20 The present invention also relates to the study of diabetes to identify molecules responsible for sensing extra-cellular carbohydrate, other receptors and genes and relationships that contribute to a diabetic state.

The present invention also relates to a sequence variation and its use in the identification of specific alleles altered in their specificity for carbohydrate compounds.

25 The present invention also relates to a recombinant construct comprising SAC1 (also referred to as *Sac*) polynucleotide suitable for expression in a transformed host cell.

The present invention also provides primers and probes specific for the detection and analysis of the SAC1 locus.

30 The present invention also relates to kits for detecting a polynucleotide comprising a portion of the SAC1 locus.

The present invention also relates to transgenic animals, which carry an altered SAC1 allele, such as a knockout mouse.

The present invention also relates to methods for screening drugs for inhibition or restoration of SAC1 function as a taste receptor.

The present invention also relates to identification of sweeteners or alcohols using the SAC1 gene and its sequence variations.

The present invention also relates to methods for screening drugs for inhibition or restoration of SAC1 function in homeostatic regulation of glucose 5 levels.

The present invention also relates to methods for screening drugs for modification of SAC1 function in the consumption of alcohol.

Finally, the present invention provides therapies directed to diabetic or obesity disorders. Therapies of diabetes and obesity include gene therapy, protein 10 replacement, protein mimetics, and inhibitors.

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1A shows genetic mapping of the SAC1 locus, using 632 F2 mice from a cross between the B6 (high preference) and 129 (low preference) strains. Mapping results were obtained with MAPMAKER/QTL Version 1.1, using an 15 unconstrained model. A black triangle at the bottom indicates peak LOD score at M134G01 marker. Horizontal line at the bottom shows a 1-LOD confidence interval.

Fig. 1B shows SAC1-containing chromosomal region defined by a donor fragment of the 129.B6-*Sac<sup>b</sup>* partially congenic mice. The partially congenic 20 strains were constructed by identifying several founder F2 mice with small fragments of the telomeric region of mouse chromosome 4 from the B6 strain and successive backcrossing to the 129 strain. Presence and size of donor fragment were determined by genotyping polymorphic markers in mice from the N4, N6, N7, N4F4, and N3F5 generations.

Fig. 1C shows average daily saccharin consumption by N6, N7, N4F4, and N3F5 segregating partially congenic 129.B6-*Sac* mice in 4-days two-bottle tests with water (means  $\pm$  SE). The open bar indicates intakes of mice that did not inherit the donor fragment. The black bar indicates intakes of mice with one or two copies of the donor fragment, which is flanked by 280G12-T7 proximally and

-4-

*D4Mon1* distally. The complete donor fragment is represented by overlapping sequences of the BAC RPCI-23-118E21 and a genomic clone (Accession AF185591), as indicated at the bottom. The size of the SAC1-containing donor fragment is 194,478 kb.

5 Fig. 1D shows BAC contig of distal chromosome 4 in the SAC1 region. Using  $^{32}\text{P}$  radioactively labeled probes from the nonrecombinant interval, a mouse BAC library (RPCI-23) was screened; positive clones were confirmed by PCR analysis and only clones positive by hybridization and by PCR are included in the contig. BAC ends were sequenced and PCR primers designed. The STS 10 content of each BAC, using all BAC ends was determined. BAC size was determined by digesting the BAC with *NotI*, and the insert size determined using pulse field gel electrophoresis.

15 Fig. 1E shows genes contained within the SAC1 nonrecombinant interval. Arrows indicate predicted direction of transcription. See Table 1 for a description 15 of gene prediction, and details concerning function.

Fig. 2A shows the mouse SAC1 gene (mSac; Accession AF311386), its 20 human ortholog (hSac), and the previously described gene T1R1, now Gpr70, are aligned above. Residues shaded in black are identical between at least two identical residues; residues in gray indicate conservative changes. The human 25 ortholog was identified by sequence homology search within the htgs database (Accession AC026283). The amino acid sequence of the human ortholog was predicted using GENSCAN. The amino acid sequence of mouse *Gpr70* was obtained by constructing primers based upon the nucleotide sequence, and taste cDNA was amplified and sequenced. This amino acid and nucleotide sequence for 25 *Gpr70* differed slightly from the initial report; the sequence reported in this paper has been deposited in GenBank (AF301161, AF301162). The location of the missense mutation is indicated by an \*.

Fig. 2B shows structure of the SAC1 gene. The six exons are shown as black boxes.

30 Fig. 2C shows conformation of a protein predicted from the *Sac* gene. To determine the transmembrane regions, the hydrophobicity was determined using

-5-

the computer program HMMTOP, and drawn with TOPO. The missense mutation is denoted with an asterisk.

Fig. 3 shows saccharin and sucrose preferences by mice from inbred strains with two different haplotypes of the *Sac* gene. The haplotype found in the 5 B6 mice and the other high sweetener-prefering inbred strains consisted of four variants, two variants were 5' of the predicted translation start codon, one variant was a missense mutation (Ile61Thr), and the last variant was located in the intron between exon 2 and 3. The strains with the B6-like haplotype of *Sac* strongly preferred saccharin ( $82 \pm 4\%$ ) and sucrose ( $86 \pm 6\%$ ), whereas strains with the 10 129-like haplotype were indifferent to these solutions ( $57 \pm 2\%$  and  $54 \pm 1\%$  respectively,  $p = 0.0015$ ).

Fig. 4A shows tissue expression of the SAC1 gene. Note that cDNA was obtained from a commercial source for the multiple tissue panel, with the exception of tongue cDNA, which was as isolated by the investigator, as described 15 within the text. Relative band intensities may differ due to differences in cDNA isolation methods or concentration.

Fig. 4B shows RNA from human fungiform papillae was obtained from biopsy material, reversed transcribed, and the resulting bands from genomic and cDNA were amplified using primers, described in the text. The bands were 20 excised from the agarose gel, purified and reamplified. The PCR product was sequenced to confirm that the bands amplified the human otholog to *Sac*.

Fig. 5 shows amino acid sequence alignment of the mouse cDNA sequence for the SAC1 gene and the cDNA for a calcium sensing metabotropic receptor. Dark areas indicated regions of shared similarity.

25 Fig. 6 plots the hydrophobicity of the SAC1 amino acid sequence as predicted by the computer program Top Pred. Note the seven transmembrane domains characteristic of G-protein coupled receptors.

## DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

The present invention employs the following definitions:

As used herein, the terms "polynucleotide" and "nucleic acid" refer to 5 naturally occurring polynucleotides, e.g., DNA or RNA. These terms do not refer to a specific length. Thus, these terms include oligonucleotide, primer, probe, etc. These terms also refer to analogs of naturally occurring polynucleotides. The polynucleotide may be double stranded or single stranded. The polynucleotides may be labeled with radiolabels, fluorescent labels, enzymatic labels, proteins, 10 haptens, antibodies, sequence tags.

For example, these terms include RNA, cDNA, genomic DNA, synthetic forms, and mixed polymers, both sense and antisense strands, and may be chemically or biochemically modified or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those skilled in the art. Such 15 modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), pendent moieties (e.g., 20 polypeptides), intercalators (e.g., acridine, psoralen, etc.), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids, etc.). Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Such 25 molecules are known in the art and include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule.

As used herein, the term "polynucleotide amplification" refers to a broad range of techniques for increasing the number of copies of specific polynucleotide sequences. Typically, amplification of either or both strand of the target nucleic acid comprises the use of one or more nucleic acid-modifying enzymes, such as a 30 DNA polymerase, a ligase, an RNA polymerase, or an RNA-dependent reverse transcriptase. Examples of polynucleotide amplification reaction include, but not

limited to, polymerase chain reaction (PCR), nucleic acid sequence based amplification (NASB), self-sustained sequence replication (3SR), strand displacement activation (SDA), ligase chain reaction (LCR), Q $\beta$  replicase system, and the like.

5 As used herein, the term "primer" refers to a nucleic acid, e.g., synthetic polynucleotide, which is capable of annealing to a complementary template nucleic acid (e.g., the SAC1 locus) and serving as a point of initiation for template-directed nucleic acid synthesis. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to hybridize  
10 with a template. Typically, a primer will include a free hydroxyl group at the 3' end. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 12 to 30 nucleotides. The term primer pair means a set of primers including a 5' upstream primer that hybridizes with the 5' end of the target sequence to be amplified and a 3' downstream primer that hybridizes  
15 with the complement of the 3' end of the target sequence to be amplified.

The present invention includes all novel primers having at least eight nucleotides derived from the SAC1 locus for amplifying the SAC1 gene, its complement or functionally equivalent nucleic acid sequences. The present invention does not include primers which exist in the prior art. That is, the present  
20 invention includes all primers having at least 8 nucleotides with the proviso that it does not include primers existing in the prior art.

"Target polynucleotide" refers to a single- or double-stranded polynucleotide which is suspected of containing a target sequence, and which may be present in a variety of types of samples, including biological samples.

25 "Antibody" refers to polyclonal and/or monoclonal antibody and fragments thereof, and immunologic binding equivalents thereof, which are capable of specifically binding to the SAC1 polypeptides and fragments thereof or to polynucleotide sequences from the SAC1 region, particularly from the SAC1 locus or a portion thereof. Antibody may be a homogeneous molecular entity, or a mixture such as a serum product made up of a plurality of different molecular entities.

Antibodies may be produced by in vitro or in vivo techniques well-known in the art. For example, for production of polyclonal antibodies, an appropriate target immune system, typically mouse or rabbit, is selected. Substantially purified antigen is presented to the immune system. Typical sites for injection are 5 in footpads, intramuscularly, intraperitoneally, or intradermally. Polyclonal antibodies may then be purified and tested for immunological response, e.g., using an immunoassay.

For production of monoclonal antibodies, protein, polypeptide, fusion protein, or fragments thereof may be injected into mice. After the appropriate 10 period of time, the spleens may be excised and individual spleen cells fused, typically, to immortalized myeloma cells under appropriate selection conditions. Thereafter, the cells are clonally separated and the supernatants of each clone tested for their production of an appropriate antibody specific for the desired 15 region of the antigen. Affinities of monoclonal antibodies are typically  $10^{-8} \text{ M}^{-1}$  or preferably  $10^{-9}$  to  $10^{-10} \text{ M}^{-1}$  or stronger.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides, or alternatively, to selection of libraries of antibodies in phage or similar vectors.

Frequently, antibodies are labeled by joining, either covalently or non- 20 covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, magnetic particles, and the like. Also, recombinant immunoglobulins may be produced.

25 "Binding partner" refers to a molecule capable of binding another molecule with specificity, as for example, an antigen and an antigen-specific antibody or an enzyme and its inhibitor. Binding partners are known in the art and include, for example, biotin and avidin or streptavidin, IgG and protein A, receptor-ligand couples, and complementary polynucleotide strands. In the case of 30 complementary polynucleotide binding partners, the partners are normally at least about 15, 20, 25, 30, 40 bases in length.

-9-

A "biological sample" refers to a sample of tissue or fluid suspected of containing an analyte (e.g., polynucleotide, polypeptide) including, but not limited to, e.g., plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, blood cells, organs, 5 tissue and samples of in vitro cell culture constituents. A biological sample is typically from human or other animal.

"Encode." A polynucleotide is said to "encode" a polypeptide if, in its native state or when manipulated by methods well-known to those skilled in the art, it can be transcribed and/or translated to produce the mRNA and/or the 10 polypeptide or a fragment thereof. The antisense strand is the complement of such a nucleic acid, and the encoding sequence can be deduced therefrom.

"Isolated" or "substantially pure" polynucleotide or polypeptide (e.g., an RNA, DNA, protein) is one which is substantially separated from other cellular components which naturally accompany a native human nucleic acid or protein, 15 e.g., ribosomes, polymerases, many other human genome sequences and proteins. The term embraces a nucleic acid or peptide sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems.

20 "SAC1 Allele" refers to normal alleles of the SAC1 locus as well as alleles carrying variations that predispose individuals to develop obesity, diabetes, or for alcohol consumption or alcoholism.

"SAC1 Locus" refers to polynucleotides, which are in the SAC1 region, that are likely to be expressed in normal individual, certain alleles of which 25 predispose an individual to develop obesity, diabetes, or alcohol consumption or alcoholism. The SAC1 locus includes coding sequences, intervening sequences and regulatory elements controlling transcription and/or translation. The SAC1 locus includes all allelic variations of the DNA sequence.

The DNA sequences used in this invention will usually comprise at least 30 about 5 codons (15 nucleotides), 7, 10, 15, 20, or 30 codons, and most preferably, at least about 35 codons. One or more introns may also be present. This number of

-10-

nucleotides is usually about the minimal length required for a successful probe that would hybridize specifically with a SAC1 locus.

"SAC1 Region" refers to a portion of mouse chromosome 4 bounded by the markers 280G12-T7 and D4Mon1 GenBank Accession number is YG7772

5 (SEQ ID NO: 652) and is

GCAGTGAGCTGCAGAGTTGCAGAATGAGGGCACTCTAAACTCATCAA  
GTGAGGAGGCCCTCCCTCACACTCCAGATGGCTGATAGGTGGCATTA  
CATGGTC(CA)<sub>n</sub>CGCGCGCACCGCCTCAGATGCAATCTCCACATTCTATA  
ACCAGATGTCTGGTAGGCCT. The CA sequence in the middle is

10 variable in length. In the B6 mouse, n = 19, while in the 129 mouse, n = 16. This region contains the SAC1 locus, including the SAC1 gene. GenBank accession number for the SAC1 gene is AF311386.

15 As used herein, a "portion" or "fragment" of the SAC1 gene, locus, region, or allele is defined as having a minimal size of at least about 15 nucleotides, or preferably at least about 20, or more preferably at least about 25 nucleotides, and may have a minimal size of at least about 40 nucleotides.

20 As used herein, the term "polypeptide" refers to a polymer of amino acids without referring to a specific length. This term includes to naturally occurring protein. The term also refers to modifications, analogues and functional mimetics thereof. For example, modifications of the polypeptide may include glycosylations, acetylations, phosphorylations, and the like. Analogues of polypeptide include unnatural amino acid, substituted linkage, etc. Also included are polypeptides encoded by DNA which hybridize under high or low stringency conditions, to the nucleic acids of interest.

25 Modification of polypeptides includes those substantially homologous to primary structural sequence, e.g., in vivo or in vitro chemical and biochemical modifications or incorporation unusual amino acids. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labeling, e.g., with radionuclides, and various enzymatic  
30 modifications, as will be readily appreciated by those well-skilled in the art. A variety of methods for labeling polypeptides and of substituents or labels useful for such purposes are well-known in the art, and include radioactive isotopes such

as  $^{32}\text{P}$ , ligands which bind to labeled antiligands (e.g., antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands which can serve as specific binding pair members for a labeled ligand. The choice of label depends on the sensitivity required, ease of conjugation with the primer, stability requirements, 5 and available instrumentation. Methods of labeling polypeptides are well-known in the art (see Sambrook et al., 1989 or Ausubel et al., 1992).

Besides substantially full-length polypeptides, the present invention provides for biologically active fragments of the polypeptides. Significant biological activities include ligand-binding, immunological activity, and other 10 biological activities characteristic of SAC1 polypeptides. Immunological activities include both immunogenic function in a target immune system, as well as sharing of immunological epitopes for binding, serving as either a competitor or substitute antigen for an epitope of the SAC1 protein. As used herein, "epitope" refers to an antigenic determinant of a polypeptide. An epitope could comprise three amino 15 acids in a spatial conformation that is unique to the epitope. Generally, an epitope consists of at least five such amino acids, and more usually consists of at least 8 to 10 such amino acids. Methods of determining the spatial conformation of such amino acids are known in the art.

For immunological purposes, tandem-repeat polypeptide segments may be 20 used as immunogens, thereby producing highly antigenic proteins. Alternatively, such polypeptides will serve as highly efficient competitors for specific binding.

Fusion proteins comprise SAC1 polypeptides and fragments. Homologous polypeptides may be fusions between two or more SAC1 polypeptide sequences or between the sequences of SAC1 and a related protein. Likewise, heterologous 25 fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. For example, ligand-binding or other domains may be "swapped" between different new fusion polypeptides or fragments. Such homologous or heterologous fusion polypeptides may display, for example, altered strength or specificity of binding. Fusion partners include 30 immunoglobulins, bacterial  $\beta$ -galactosidase, trpE, protein A,  $\beta$ -lactamase,  $\alpha$ -amylase, alcohol dehydrogenase, and yeast  $\alpha$  mating factor.

-12-

Fusion proteins will typically be made by either recombinant nucleic acid methods or may be chemically synthesized. Techniques for the synthesis of polypeptides are known in the art.

Functional mimetics of a native polypeptide may be obtained using known methods in the art. For example, polypeptides may be least about 50% homologous to the native amino acid sequence, preferably in excess of about 70%, and more preferably at least about 90% homologous. Substitutions typically contain the exchange of one amino acid for another at one or more sites within the polypeptide, and may be designed to modulate one or more properties of the polypeptide, such as stability against proteolytic cleavage, without the loss of other functions or properties. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. Preferred substitutions are ones which are conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well-known in the art and typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and tyrosine, phenylalanine.

Certain amino acids may be substituted for other amino acids in a polypeptide structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules or binding sites on proteins interacting with a polypeptide. Since it is the interactive capacity and nature of a polypeptide which defines that polypeptide's biological functional activity, certain amino acid substitutions can be made in a protein sequence, and its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydrophobic amino acid index in conferring interactive biological function on a protein is generally understood in the art. Alternatively, the substitution of like amino acids can be made effectively on the basis of hydrophilicity.

-13-

A peptide mimetic may be a peptide-containing molecule that mimics elements of protein secondary structure. The underlying rationale behind the use of peptide mimetics is that the peptide backbone of proteins exists chiefly to orient amino acid side chains in such a way as to facilitate molecular interactions, such 5 as those of antibody and antigen, enzyme and substrate or scaffolding proteins. A peptide mimetic is designed to permit molecular interactions similar to the natural molecule. A mimetic may not be a peptide at all, but it will retain the essential biological activity of a natural polypeptide.

Polypeptides may be produced by expression in a prokaryotic cell or 10 produced synthetically. These polypeptides typically lack native post-translational processing, such as glycosylation. Polypeptides may be labeled with radiolabels, fluorescent labels, enzymatic labels, proteins, haptens, antibodies, sequence tags.

“SAC1 polypeptide” refers to a protein or polypeptide encoded by the SAC1 locus, variants, fragments or functional mimics thereof. A SAC polypeptide 15 may be that derived from any of the exons described herein which may be in isolated and/or purified form. The length of SAC1 polypeptide sequences is generally at least about 5 amino acids, usually at least about 10, 15, 20, 30 residues.

“Alcohol consumption” relates to the intake and/or preference of an animal 20 for ethanol.

“Diabetes” refers to any disorder that exhibits phenotypic features of an increased or decreased level of a biological substance associated with glucose or fatty acid metabolism. The term “carbohydrate” refers to simple mono and disaccharides.

25 The terms “sequence variation” or “variant form” encompass all forms of polymorphism and mutations. A sequence variation may range from a single nucleotide variation to the insertion, modification, or deletion of more than one nucleotide. A sequence variation may be located at the exon, intron, or regulatory region of a gene.

30 Polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A biallelic polymorphism has two forms. A triallelic polymorphism has three forms. A

-14-

polymorphic site is the locus at which sequence divergence occurs. Diploid organisms may be homozygous or heterozygous for allelic forms. Polymorphic sites have at least two alleles, each occurring at frequency of greater than 1% of a selected population. Polymorphic sites also include restriction fragment length 5 polymorphisms, variable number of tandem repeats (VNTRs), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide repeats, simple sequence repeats, and insertion elements. The first identified allelic form may be arbitrarily designated as the reference sequence and other allelic forms may be designated as alternative or variant alleles. The allelic form 10 occurring most frequently in a selected population is sometimes referred to as the wild type form or the consensus sequence.

Mutations include deletions, insertions and point mutations in the coding and noncoding regions. Deletions may be of the entire gene or of only a portion of the gene. Point mutations may result in stop codons, frameshift mutations, or 15 amino acid substitutions. Somatic mutations are those which occur only in certain tissues, such as liver, heart, etc. and are not inherited in the germline. Germline mutations can be found in any of a body's tissues and are inherited.

"Operably linked" refers to a juxtaposition wherein the components are in a relationship permitting them to function in their intended manner. For instance, a 20 promoter is operably linked to a coding sequence if the promoter affects its transcription or expression.

The term "probes" refers to polynucleotide of any suitable length which allows specific hybridization to the target region. Probes may be attached to a label or reporter molecule using known methods in the art. Probes may be selected 25 by using homologous polynucleotides. Alternatively, polynucleotides encoding these or similar polypeptides may be synthesized or selected by use of the redundancy in the genetic code. Various codon substitutions may be introduced, e.g., by silent changes (thereby producing various restriction sites) or to optimize expression for a particular system. Mutations may be introduced to modify the 30 properties of the polypeptide, perhaps to change ligand-binding affinities, interchain affinities, or the polypeptide degradation or turnover rate.

-15-

Probes comprising synthetic oligonucleotides or other polynucleotides of the present invention may be derived from naturally occurring or recombinant single- or double-stranded polynucleotides, or be chemically synthesized. Probes may also be labeled by nick translation, Klenow fill-in reaction, or other methods  
5 known in the art.

Portions of the polynucleotide sequence having at least about 8 nucleotides, usually at least about 15 nucleotides, and fewer than about 6 kb, usually fewer than about 1.0 kb, from a polynucleotide sequence encoding SAC1 are preferred as probes.

10 The terms "isolated," "substantially pure," and "substantially homogeneous" are used interchangeably to describe a protein or polypeptide which has been separated from components which accompany it in its natural state. A monomeric protein is substantially pure when at least about 60% to 75% of a sample exhibits a single polypeptide sequence. A substantially pure protein  
15 will typically comprise about 60% to 90% W/W of a protein sample, more usually about 95%, and preferably will be over about 99% pure. Protein purity or homogeneity may be indicated by a number of means well-known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes, higher  
20 resolution may be provided by using HPLC or other means well-known in the art which are utilized for purification.

A SAC1 protein is substantially free of naturally associated components when it is separated from the native contaminants which accompany it in its natural state. Thus, a polypeptide which is chemically synthesized or synthesized  
25 in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well-known in the art.

"Recombinant nucleic acid" is a nucleic acid which is not naturally occurring, or which is made by the artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by either chemical synthesis means, or by the artificial manipulation of isolated

-16-

segments of nucleic acids, e.g., by genetic engineering techniques. Such is usually done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid 5 segments of desired functions to generate a desired combination of functions.

“Regulatory sequences” refers to those sequences normally within 100 kb of the coding region of a locus, but they may also be more distant from the coding region, which affect the expression of the gene (including transcription of the gene, and translation, splicing, stability or the like of the messenger RNA).

10 “Substantial homology or similarity.” A nucleic acid or fragment thereof is of substantially homologous (“or substantially similar”) to another if, when optimally aligned (with appropriate nucleotide insertions or deletions) with the other nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 60% of the nucleotide bases, usually at least about 70%, 15 more usually at least about 80%, preferably at least about 90%, and more preferably at least about 95-98% of the nucleotide bases.

Identity means the degree of sequence relatedness between two polypeptide or two polynucleotides sequences as determined by the identity of the match between two strings of such sequences. Identity can be readily calculated 20 (Lesk A.M., ed., *Computational Molecular Biology*, New York: Oxford University Press, 1988; Smith D.W., ed., *Biocomputing: Informatics and Genome Projects*, New York: Academic Press, New York, 1993; Griffin A.M., and Griffin H.G., eds., *Computer Analysis of Sequence Data*, Part 1, New Jersey: Humana Press, 1994; von Heinje G., *Sequence Analysis in Molecular Biology*, 25 Academic Press, 1987; and Gribskov M. and Devereux J., eds., *Sequence Analysis Primer*, New York: M Stockton Press, 1991).

Alternatively, substantial homology or similarity exists when a nucleic acid or fragment thereof will hybridize to another nucleic acid (or a complementary strand thereof) under selective hybridization conditions, to a 30 strand, or to its complement. Selectivity of hybridization exists when hybridization which is substantially more selective than total lack of specificity occurs. Typically, selective hybridization will occur when there is at least about

-17-

55% homology over a stretch of at least about 14 nucleotides, preferably at least about 65%, more preferably at least about 75%, and most preferably at least about 90%. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will often be over a stretch of at least about 5 9 nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides, and preferably at least about 36 or more nucleotides.

Nucleic acid hybridization will be affected by such conditions as salt concentration, temperature, or organic solvents, in addition to the base 10 composition, length of the complementary strands, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. Stringent temperature conditions will generally include temperatures in excess of 30°C, typically in excess of 37°C, and preferably in excess of 45°C. Stringent salt conditions will ordinarily be less than 15 1000 mM, typically less than 500 mM, and preferably less than 200 mM. However, the combination of parameters is much more important than the measure of any single parameter.

The terms "substantial homology" or "substantial identity," when referring to polypeptides, indicate that the polypeptide or protein in question exhibits at 20 least about 30% identity with an entire naturally-occurring protein or a portion thereof, usually at least about 70% identity, and preferably at least about 95% identity.

Homology, for polypeptides, is typically measured using sequence analysis software (see, e.g., the Sequence Analysis Software Package of the Genetics 25 Computer Group, University of Wisconsin Biotechnology Center). Protein analysis software matches similar sequences using measures of homology assigned to various substitutions, deletions and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, 30 glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

"Substantially similar function" refers to the function of a modified nucleic acid or a modified protein, with reference to the wild-type SAC1 nucleic acid or

-18-

wild-type SAC1 polypeptide. The modified polypeptide will be substantially homologous to the wild-type SAC1 polypeptide and will have substantially the same function. The modified polypeptide may have an altered amino acid sequence and/or may contain modified amino acids. In addition to the similarity of

5. function, the modified polypeptide may have other useful properties, such as a longer half-life. The similarity of function (activity) of the modified polypeptide may be substantially the same as the activity of the wild-type SAC1 polypeptide. Alternatively, the similarity of function (activity) of the modified polypeptide may be higher than the activity of the wild-type SAC1 polypeptide. The modified

10 polypeptide is synthesized using conventional techniques, or is encoded by a modified nucleic acid and produced using conventional techniques. The modified nucleic acid is prepared by conventional techniques. A nucleic acid with a function substantially similar to the wild-type SAC1 gene function produces the modified protein described above.

15 A polypeptide "fragment," "portion," or "segment" is a stretch of amino acid residues of at least about 5 to 7 contiguous amino acids, often at least about 7 to 9 contiguous amino acids, typically at least about 9 to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids.

20 The polypeptides of the present invention, if soluble, may be coupled to a solid-phase support, e.g., nitrocellulose, nylon, column packing materials (e.g., Sepharose beads), magnetic beads, glass wool, plastic, metal, polymer gels, cells, or other substrates. Such supports may take the form, for example, of beads, wells, dipsticks, or membranes.

25 "Target region" refers to a region of the nucleic acid which is amplified and/or detected. The term "target sequence" refers to a sequence with which a probe or primer will form a stable hybrid under desired conditions.

II. Positional Cloning of Mouse SAC1 Gene and the Discovery of a Gene and Its Sequence Variation Associated With Altered Sensation for Carbohydrates

30

Inbred strains of mice differ in their intake of sweeteners (Bachmanov A.A., Reed D.R., Tordoff M.G., Price R.A., and Beauchamp G.K. Intake of ethanol, sodium chloride, sucrose, citric acid, and quinine hydrochloride solutions by mice: a genetic analysis. *Behavior Genetics*, 1996;26:563-573; Lush I.E., The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet Res*, 1989;53:95-99; Lush I. The genetics of bitterness, sweetness, and saltiness in strains of mice. In *Genetics of Perception and Communication*, Vol. 3, eds. Wysocki C. and Kare M., New York: Marcel Dekker, 1991:227-235; Capretta P.J. Saccharin and saccharin-glucose ingestion in two inbred strains of *Mus musculus*. *Psychon. Sci.*, 1970;21:133-135; Nachman M. The inheritance of saccharin preference. *Journal of Comp Physiol Psychol*, 1959;52:451-457). Breeding and linkage experiments suggest that a single gene, the *Sac* locus (for saccharin intake), accounts for a large proportion of the genetic variance (Fuller J.L. Single-locus control of saccharin preference in mice. *Journal of Heredity*, 1974;65:33-36; Capeless C.G. and Whitney G. The genetic basis of preference for sweet substances among inbred strains of mice: preference ratio phenotypes and the alleles of the *Sac* and *dpa* loci. *Chem Senses*, 1995;20:291-298; Bachmanov A.A. et al. Sucrose consumption in mice: major influence of two genetic loci affecting peripheral sensory responses. *Mammalian Genome*, 1997;8:545-548; Belknap J.K. et al. Single-locus control of saccharin intake in BXD/Ty recombinant inbred (RI) mice: some methodological implications for RI strain analysis. *Behav Genet*, 1992;22:81-100; Blizzard D.A., Kotlus B., and Frank M.E. Quantitative trait loci associated with short-term intake of sucrose, saccharin and quinine solutions in laboratory mice. *Chem Senses*, 1999;24:373-85). Using genetic and physical mapping methods, an interval of 194 kb was identified at the telomeric end of mouse chromosome 4 that contains the *Sac* locus. BAC sequencing within this interval led to the identification of a gene that has a 30% amino acid homology with other putative taste receptors (Hoon M.A. et al. Putative mammalian taste receptors: a class of taste-specific GPCRs with distinct topographic selectivity. *Cell*, 1999;96:541-551). This gene is expressed in mouse tongue. Mutation detection on this gene revealed a missense mutation (Ile61Thr) with four other sequence variants define a haplotype found in mice

-20-

with low sweetener preference (129, Balb/c, AKR, and DBA2). An alternative five variant haplotype is found in mice with a high preference for sweet fluids (B6, SWR, IS, ST, and SEA). A human ortholog of this gene exists, and is expressed in human taste papillae. We therefore suggest that this gene is a sweet 5 taste receptor, and variation within this gene is responsible for the phenotype of the *Sac* locus.

To identify this locus, mice from the high sweetener preference (C57BL/6ByJ; B6) and the low sweetener preference (129P3/J; formerly 129/J, abbreviated here as 129) were used as parental strains to produce an F2 10 generation. The F2 mice were phenotyped for sweetener preference using 96-hour two-bottle taste tests and genotyped with markers polymorphic between the B6 and 129 strains (Fig. 1A). The results of this analysis indicated peak linkage near marker D18346 with the B6 allele having a dominant mode of inheritance. Using recombinant mice from the F2 generation, 129.B6-*Sac* partially congenic 15 mice were created, using genotypic (B6 allele at D18346; Fig. 1B) and phenotypic (high saccharin intake; Fig. 1C) characteristics as selection criteria for each generation. Genotyping of partially congenic mice with polymorphic markers defined the *Sac* nonrecombinant interval. Radiation hybrid mapping was conducted with additional markers (R74924, D18402, D18346, Agrin, V2r2 and 20 D4Ert296e). These markers were amplified using DNA and mouse and hamster control DNA in the T31 mouse radiation hybrid panel, scored for the presence or absence of an appropriately sized band, and the data analyzed by the Jackson Laboratory. All markers were within the SAC1 confidence interval suggested by the initial linkage analysis, and were used in subsequent analyses.

25 A BAC library was screened with markers within the nonrecombinant interval, and a contig was developed (Fig. 1D). A BAC clone was selected for sequencing (RPCI-23-118E21, 246 kb). Within this BAC, a gene with a 30% homology to T1R1 (a putative taste receptor) was discovered (Fig. 2A), along with other ESTs and known genes (Table 1). The human ortholog to this gene was 30 identified from a BAC available in the public *hg* database, and the predicted protein sequence was aligned with SAC1 and T1R1. SAC1 is 858 amino acids in length and contains six exons; the intron and exon boundaries were determined by

-21-

sequencing of the mouse tongue cDNA (Fig. 2B). The secondary structure of this protein with regards to transmembrane domains was predicted (Fig. 2C).

To determine whether this gene might contain functional polymorphisms that could account for the behavioral differences between the two strains, 11.8 kb 5 of sequence, including the SAC1 gene and several kb up and downstream were amplified with PCR primers and then sequenced using DNA from the high and low preferring strains (Lush I.E., The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet Res* 1989;53:95-99; Lush I. The genetics of bitterness, sweetness, and saltiness in strains of mice. In *Genetics of Perception* 10 and *Communication*, Vol. 3, eds. Wysocki C. and Kare M., New York: Marcel Dekker, 1991:227-235). Many variants existed between these strains, and of these, five variants were found in the low preferring strains but not in the high preferring strain. One of these variants results in a missense mutation (Ile61Thr; Fig. 2). The other four variants were in non-coding regions (T>A -2383 nt; A>G -183 nt; 15 A>G +134 nt; T>C +651 nt, between exon 2 and 3). These five variants will be referred to as the 129-like or B6-like haplotypes. Additional inbred strains of mice with known saccharin and sucrose preferences (Lush I.E., The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet Res*, 1989;53:95-99; Lush I. The genetics of bitterness, sweetness, and saltiness in 20 strains of mice. In *Genetics of Perception and Communication*, Vol. 3, eds. Wysocki C. and Kare M., New York: Marcel Dekker, 1991:227-235; Lush I.E. and Holland G. The genetics of tasting in mice. V. Glycine and cyclohexamide. *Genet Res*, 1988;52:207-212) were also sequenced. The 129-like haplotype was found in mice with lower sweetener preference and the B6-like haplotype was 25 found in mice with higher sweetener preference (Fig. 3).

B6 mice have higher maximal gustatory neural firing in response to sweeteners compared with 129 mice, as do the 129.B6-Sac partially congenic strains (Bachmanov A.A. et al. Sucrose consumption in mice: major influence of two genetic loci affecting peripheral sensory responses. *Mammalian Genome*, 30 1997;8:545-548). Thus, the SAC1 gene is likely to be expressed in tongue. To test this hypothesis, RNA from mouse and human tongue was extracted, reversed transcribed into cDNA and primers, chosen to span an intron, were used in a PCR

-22-

reaction. Genomic and cDNA yielded bands of different sizes, which were purified and sequenced (Figure 4AB). Sequencing results confirmed that the bands were derived from this gene with the appropriate intron/exon boundaries. Further analysis of expression in cDNA in mouse tissue, using commercially 5 available mouse cDNA, indicated this gene is also expressed is widely expressed. The broad range of tissue expression of this gene may indicate that other tissues use this receptor to sense extra cellular sugars (Fig. 4A).

Hoon et al. identified a gene, *Gpr70* (formerly TR1 or T1R1) as a putative sweet receptor based mainly on its expression in anterior tongue taste cells. Since 10 it also mapped to distal chromosome 4, it was a logical candidate for SAC1. However, we have shown that *Gpr70* is at least 4 cM proximal to SAC1 (Li X. et al. The saccharin preference locus (*Sac*) and the putative sweet taste receptor (*Gpr70*) gene have distinct locations on mouse chromosome 4. *Mammalian Genome*, 2001;12:13-16). Nevertheless, *Gpr70* could be an additional sweet 15 receptor and there could be others. It has been argued based upon human psychophysical studies and studies of sweet taste transduction mechanisms that there must be more than one sweet receptor. Other lines of evidence, however, are more consistent with the existence of one or a very few receptors (Bartoshuk L.M. Is sweetness unitary? An evaluation of the evidence for multiple sweeteners. In 20 *Sweetness*, ed. Dobbing, J., London: Springer-Verlag, 1987:33-46). At present no evidence has been found of a family of *Sac*-like receptors resembling the large family of bitter receptors recently reported (Matsunami H., Montmayeur J.P., and Buck L.B. A family of candidate taste receptors in human and mouse [see 25 comments]. *Nature*, 2000;404:601-604; Adler E. et al. A novel family of mammalian taste receptors [see comments]. *Cell*, 2000;100:693-702). The sweet substances that exist in nature, which presumably shaped the evolution of sweet receptor(s), are likely much more similar amongst themselves, mostly simple sugars, than are the vast array of structurally diverse bitter tasting compounds.

A receptor for the sugar trehalose has recently been identified in the fruit 30 fly, *Drosophila melanogaster*. Surprisingly, the trehalose and other fly taste receptors, have no homology with SAC1. The specialization of flies for the sugar trehalose may account for this divergence.

-23-

There may be multiple sweet receptors; evidence from across species comparisons, psychophysical cross adaptation, and sweetness competitors has been reviewed (Bartoshuk L.M. Is sweetness unitary? An evaluation of the evidence for multiple sweeteners. In *Sweetness*, ed. Dobbing, J., London: Springer-Verlag, 1987:33-46). The SAC1 gene accounts for ~40% of the genetic differences in sweet perception between these two particular strains of mice, but other receptors, and other alleles of these receptors may exist.

Because sucrose is perceived to be bad for human health, considerable resources are directed toward the discovery of high potency, low caloric sweeteners. Most of the most widely known high potency sweeteners were discovered serendipitously, i.e., the sweetener was synthesized for a different purpose and someone in the laboratory accidentally tasted it and discovered it was sweet (Walters E.D. The rational discovery of sweeteners. In *Sweeteners. Discovery, molecular design, and chemoreception*, eds. Walters D.E., Orthoefer F.T., and DuBois G.E., American Chemical Society, USA, 1991:1-11). More direct methods, however, have been employed to identify new sweet compounds, and the sweet receptor has been extensively modeled to predict which ligands will be sweet.

It is not known how or why different alleles of SAC1 arose in inbred strains of mice but their existence, in addition to providing us with a tool to identify a sweet receptor, raises the question of whether they might also characterize human populations. There appear to exist reliable individual differences in human sensitivity and preference for sweet sugars but whether these are genetically influenced remains to be determined. The identification of SAC1 should facilitate research in this area. Also, the observation that SAC1 is expressed in several tissues in addition to tongue raises the interesting possibility that it could be involved in other aspects of sugar recognition and that allelic variants in this gene could be related to diseases or conditions such as diabetes and obesity.

Alleles of the gene described in this application are likely to account for the SAC1 behavioral and neurological phenotype for four reasons. First, the SAC1 nonrecombinant region is small, less than 194 kb; this gene lies within this

-24-

nonrecombinant interval and the peak of LOD score corresponds closely with the location of the gene. Second, of the genes contained within this region, no others are viable candidates for SAC1. Third, this gene has sequence homology to other putative taste receptors, and is expressed in the tongue. Finally, a haplotype with a 5 missense mutation is found in mice with low sweetener preference but not in mice with high sweetener preference. These data strongly suggest that mutations of this gene account for differences in the acceptance and preference for sweeteners attributed to the SAC1 locus.

Among the multiple mechanisms involved in regulation of ethanol intake, 10 one of the least appreciated factors is the perception of its flavor (Nachman M., Larue C., Le Magnen J. The role of olfactory and orosensory factors in the alcohol preference of inbred strains of mice. *Physiology Behavior*, 1971;6:53-95). Although individual variability in the perception of ethanol flavor by adults and children was described over 60 years ago (Richter C.P. Alcohol as a food. *Quart. 15 J. Studies Alcohol*, 1941;1:650-62), the hypothesis that individual differences in alcohol chemosensory perception can affect alcohol intake did not receive due attention. As a result, the relationship between alcohol chemosensation and intake is not well-understood. Humans perceive ethanol flavor as a combination of components, including sweetness, bitterness, odor and irritation (burning 20 sensation), which depend on ethanol concentration (Green B.G. The sensitivity of the tongue to ethanol. *Ann. NY. Acad. Sci.*, 1987;510:315-7; Bartoshuk L.M., Conner E., Grubin D., Karrer T., Kochenbach K., Palsco M., et al. PROP supertasters and the perception of ethyl alcohol. *Chem. Senses*, 1993.). Rats detect sweet (sucrose-like) and bitter (quinine-like) sensory components in ethanol 25 (Kiefer S.W., Lawrence G.J. The sweet-bitter taste of alcohol: aversion generalization to various sweet-quinine mixtures in the rat. *Chem. Senses*, 1988;13:633-41; Kiefer S.W., Mahadevan R.S. The taste of alcohol for rats as revealed by aversion generalization tests. *Chem. Senses*, 1993;18:509-22) and probably perceive the other components detected by humans as well. 30 The relationship between ethanol and sweetener perception and consumption has been studied the most and is supported by several lines of evidence:

-25-

(a) Electrophysiological recordings from gustatory nerves indicate that lingual application of ethanol activates sweetener-responsive neural fibers (Hellekant G., Danilova V., Roberts T., Ninomiya Y. The taste of ethanol in a primate model: I. Chorda tympani nerve response in *Macaca mulatta*. *Alcohol*, 1997;14:473-84; Sako N., Yamamoto T. Electrophysiological and behavioral studies on taste effectiveness of alcohols in rats. *Am. J. Physiol.*, 1999;276:R388-96).

5 (b) Conditioned taste aversions generalize between ethanol and sucrose (Kiefer S.W., Lawrence G.J. The sweet-bitter taste of alcohol: aversion generalization to various sweet-quinine mixtures in the rat. *Chem. Senses*, 1988;13:633-41; Kiefer S.W., Mahadevan R.S. The taste of alcohol for rats as revealed by aversion generalization tests. *Chem. Senses*, 1993;18:509-22; Lawrence G.J., Kiefer S.W. Generalization of specific taste aversions to alcohol in the rat. *Chem. Senses*, 1987;12:591-9; Blizzard 10 D.A., McClearn G.E. Association between ethanol and sucrose intake in the laboratory mouse: exploration via congenic strains and conditioned taste aversion. *Alcohol. Clin. Exp. Res.*, 2000;24:253-8.), suggesting that ethanol and sucrose share the same taste property, most likely sweetness.

15 (c) Genetic associations between preferences for ethanol and sweeteners were found among some rat and mouse strains and within their segregating crosses (Overstreet D.H., Kampov-Polevoy A.B., Rezvani A.H., Murelle L., Halikas J.A., Janowsky D.S. Saccharin intake predicts ethanol intake in genetically heterogeneous rats as well as different rat strains. *Alcohol. Clin. Exp. Res.*, 1993;17:366-9; Sinclair J.D., Kampov-Polevoy A., 20 Stewart R., Li T-K. Taste preferences in rat lines selected for low and high alcohol consumption. *Alcohol*, 1992;9:155-60; Stewart R.B., Russell R.N., Lumeng L., Li T-K., Murphy J.M. Consumptions of sweet, salty, sour, and bitter solutions by selectively bred alcohol-preferring and alcohol- 25 nonpreferring lines of rats. *Alcohol. Clin. Exp. Res.*, 1994;18:375-81; Belknap J.K., Crabbe J.C., Young E.R. Voluntary consumption of alcohol in 30 15 inbred mouse strains. *Psychopharmacol.*, 1993;112:503-10;

-26-

Bachmanov A.A., Reed D.R., Tordoff M.G., Price R.A., Beauchamp G.K. Intake of ethanol, sodium chloride, sucrose, citric acid, and quinine hydrochloride solutions by mice: a genetic analysis. *Behav. Genet.*, 1996;26:563-73; Bachmanov A.A., Tordoff M.G., Beauchamp G.K.

5 Ethanol consumption and taste preferences in C57BL/6ByJ and 129/J mice. *Alcohol. Clin. Exp. Res.*, 1996;20:201-6), reviewed in (Kampov-Polevoy A.B., Garbutt J.C., Janowsky D.S. Association between preference for sweets and excessive alcohol intake: a review of animal and human studies. *Alcohol. Alcohol.*, 1999;34:386-95; Overstreet D.H.,

10 Rezvani A.H., Parsian A. Behavioural features of alcohol-preferring rats: focus on inbred strains. *Alcohol. Alcohol.*, 1999;34:378-85); with some exceptions (Phillips T.J., Crabbe J.C., Metten P., Belknap J.K. Localization of genes affecting alcohol drinking in mice. *Alcohol. Clin. Exp. Res.*, 1994;18:931-941; Parsian A., Overstreet D.H., Rezvani A.H.

15 Independent segregation of alcohol and saccharin intakes in F2 progeny from FH/ACI intercross (Abstract). *Alcohol. Clin. Exp. Res.*, 2000;24(Supplement):58A)).

(d) Human studies show that alcoholics have a stronger liking of concentrated sucrose compared with nonalcoholics (Kampov-Polevoy A.B., Garbutt J.C., Davis C.E., Janowsky D.S. Preference for higher sugar concentrations and Tridimensional Personality Questionnaire scores in alcoholic and nonalcoholic men. *Alcohol. Clin. Exp. Res.*, 1998;22:610-4; Kampov-Polevoy A.B., Garbutt J.C., Janowsky D. Evidence of preference for a higher concentration sucrose solution in alcoholic men. *American Journal of Psychiatry*, 1997;154:269-70).

20 There are several possible mechanisms that could underlie the association between sweetener and ethanol responses:

(a) *Common peripheral taste mechanisms*, which may involve the interaction of ethanol with a peripheral sweet taste transduction. At least one such common peripheral mechanism is mediated by the *Gpr98* gene (SAC1 locus) encoding a sweet taste receptor (as described below).

25

-27-

(b) *Common brain mechanisms.* The regulation of ingestive responses to ethanol and sweeteners may involve common opioidergic, serotonergic and dopaminergic brain neurotransmitter systems (Gosnell B.A., Majchrzak M.J. Centrally administered opioid peptides stimulate saccharin intake in nondeprived rats. *Pharm. Biochem. Behav.*, 1989;33:805-10; George S.R., Roldan L., Lui A., Naranjo C.A. Endogenous opioids are involved in the genetically determined high preference for ethanol consumption. *Alcohol. Clin. Exp. Res.*, 1991;15:668-72; Hubell C.L., Marglin S.H., Spitalnic S.J., Abelson M.L., Wild K.D., Reid L.D. 5 Opioidergic, serotonergic, and dopaminergic manipulations and rats' intake of a sweetened alcoholic beverage. *Alcohol*, 1991;8:355-67; Pucilowski O., Rezvani A.H., Janowsky D.S. Suppression of alcohol and saccharin preference in rats by a novel  $\text{Ca}^{2+}$  channel inhibitor, Goe 5438. *Psychopharmacol.*, 1992;107:447-52). These mechanisms could be 10 responsible for the emotional response to the pleasantness of ethanol or sweeteners, or the motivational mechanisms driving their intakes.

(c) *Common signals related to the caloric value of ethanol and sugars* (Gentry R.T., Dole V.P. Why does a sucrose choice reduce the consumption of alcohol in C57BL/6J mice? *Life Sci.*, 1987;40:2191-4). Ethanol is 15 metabolized in the body through some of the same pathways as carbohydrates and provides comparable energy. Thus, energy derived from carbohydrates and ethanol may have similar rewarding effects through the same hunger and satiety mechanisms.

(d) *Incidental genetic linkage.* Different genes affecting responses to ethanol 20 and sweeteners may reside nearby on the same chromosome. 25

Ethanol consumption is a complex trait, depending on multiple mechanisms of its regulation and determined by multiple genes. A body of evidence suggests that ethanol consumption may depend on perception of its flavor, and that there is an association between perception and consumption of 30 ethanol and sweet-tasting compounds. However, only a few genes have been identified as candidates affecting ethanol consumption.

-28-

The present invention provides that a gene, SAC1, is associated with the detection of a sensing of carbohydrates, other sweet compounds, and alcohols including ethanol. The sequence of the mouse SAC1 cDNA (SEQ ID NO: 1) is:

ATGCCAGCTTGGCTATCATGGGTCTAGCCTGGCTGCTTCTGGAGC  
5 TTGGGATGGGGCCTCTTGTCTGTACAGCAATTCAAGGCACAAG  
GGGACTACATACTGGCGGGCTATTCCCCTGGCTAACCGAGGAGG  
CCACTCTCAACCAGAGAACACAACCCAACAGCATCCGTGCAACAGGT  
TCTCACCCCTGGTTGTTCTGGCCATGGCTATGAAGATGGCTGTGGA  
GGAGATCAACAATGGATCTGCCTGCTCCCTGGCTGCGGCTGGCTA  
10 TGACCTATTGACACATGCTCCGAGCCAGTGGTCACCATGAAATCCAG  
TCTCATGTTCTGGCCAAGGTGGCAGTCAAAGCATTGCTGCCTACTG  
CAACTACACACAGTACCAACCCGTGTGCTGGCTGTACGGCCCCCA  
CTCATCAGAGCTTGCCTCATTACAGGCAAGTTCTCAGCTTCTCCTC  
ATGCCACAGGTCAAGCTATAGTGCCAGCATGGATCGGCTAAGTGACCGG  
15 GAAACGTTCCATCCTCTTCCGCACAGTGCCCAGTGACCGGGTGCAG  
CTGCAGGCAGTTGTGACTCTGTTGCAGAACTCAGCTGGAACTGGTG  
GCCGCCTTAGGGAGTGATGACTATGGCCGGAAGGTCTGAGCATT  
TTTCTAGTCTGCCAATGCACGAGGTATCTGCATCGCACATGAGGGC  
CTGGTGCACACATGACACTAGTGGCCAACAGTTGGCAAGGTGCTG  
20 GATGTACTACGCCAAGTGAACCAAAGTAAAGTACAAGTGGTGGTGTG  
TTGCCCTCTGCCGTGCTGTCTACTCCCTTTAGTTACAGCATCCATCA  
TGGCCTCTACCCAAGGTATGGTGGCCAGTGAGTCITGGCTGACATC  
TGACCTGGTCATGACACTTCCAATATTGCCGTGTGGCACTGTGCTT  
GGGTTTTGCAGCGGGTGCCTACTGCCGAATTTCCTATTATGTGG  
25 AGACTCACCTGCCCTGGCCGCTGACCCAGCATTCTGTGCCTACTGAA  
TGC GGAGTTGGATCTGGAGGAACATGTGATGGGCAACGCTGTCCACG  
GTGTGACGACATCATGCTGCAGAACCTATCATCTGGCTGTTGCAGAA  
CCTATCAGCTGGCAATTGCACCAACCAAAATTGCAACCTATGCAGC  
TGTGTACAGTGTGGCTCAAGCCCTCACAAACACCCCTACAGTGCAATGT  
30 CTCACATTGCCACGTATCAGAACATGTTCTACCCCTGGCAGCTCCTGGA  
GAACATGTACAATATGAGTTCCATGCTCGAGACTTGACACTACAGTT

-29-

TGATGCTGAAGGGAATGTAGACATGGAATATGACCTGAAGATGTGGGT  
GTGGCAGAGCCCTACACCTGTATTACATACTGTGGGCACCTCAACCGG  
CACCCCTCAGCTGCAGCAGTCTAAATGTACTGCCAGGCAACCAGGT  
GCCAGTCTCCCAGTGTCCCAGTGCAGAAGATGCCAGGTTGCCCG  
5 AGTAAAGGGCTTCATTCTGCTGCTATGACTGCGTGGACTGCAAGGC  
GGGCAGCTACCGGAAGCATCCAGATGACTTCACCTGTACTCCATGTAA  
CCAGGACCAGTGGTCCCCAGAGAAAAGCACAGCCTGCTTACCTCGCAG  
GCCCAAGTTCTGGCTGGGGGAGCCAGTTGTGCTGTCACTCCTCCTG  
CTGCTTGCGCTGGTGTGGCTAGCACTGGCTGCTCTGGGCTCTG  
10 TCCACCACTGGGACAGCCCTTGTCAGGCCTCAGGTGGCTCACAGT  
TCTGCTTGCGCTGATCTGCCCTAGGCCTCTGCCTCAGTGTCCCTCTG  
TTCCCAGGGCGGCCAAGCTCTGCCAGCTGCCCTGCACAACAACCAATG  
GCTCACCTCCCTCTCACAGGCTGCCCTGAGCACACTCTCCTGCAAGCAG  
CTGAGACCTTGTGGAGTCTGAGCTGCCACTGAGCTGGCAAACGGC  
15 TATGCAGCTACCTCGGGACTCTGGGCTGGTAGTGGTACTGTGG  
CCACTTTGTGGAGGCAGCACTATGTGCCCTGGTATTGATCGCTTCCC  
ACCAGAGGTGGTGACAGACTGGCAGTGCTGCCACAGAGGTACTGG  
AGCACTGCCACGTGCCCTGGTCAGCCTGGCTGGTGACATCA  
CCAATGCAATGTTAGCTTCCTGCTTCTGGCACCTCCTGGTACA  
20 GAGCCAGCCTGGCCGTACAACCGTGCCGTGGTCTCACCTCGCCAT  
GCTAGCTTATTTCATCACCTGGGCTCTTGTGCCCTCCTGGCCAAT  
GTGCAGGTGGCCTACCAGCCAGCTGTGCAGATGGGTGCTATCCTAGTC  
TGTGCCCTGGCATCCTGGTACCTCCACCTGCCCAAGTGCTATGTG  
TTCTTGGCTGCCAAAGCTCAACACCCAGGAGTTCTCCTGGGAAGGA  
25 ATGCCAAGAAAGCAGCAGATGAGAACAGTGGCGGTGGTGAGGCAGCT  
CAGGGACACAATGAATGA

The genomic DNA sequence of the mouse SAC1 gene (SEQ ID NO: 2)

is:

ATCTGAGCCTAGACACAGCACTGGTGCCAGGCAAACACTCCTGGGCC  
30 TACATGCTTGGG

-30-

GCCTCTCATATTCCAAAAGCTGTCTTGGTAAGATGAAGTTCCTCTG  
GCAGTGGCATG  
AGTGCAGGGCTCTTCCCTGCCCTCACCTGCTTCTGATAGTCTCT  
CTGCATACCA  
5 AACAGGCCCTGTCTCCTGGAAATGGAAACTATGAAATCAATAGCTG  
AGGCTTCTCTAG  
GAAAGCCTGCCCTGGTCAGTACAACCTGTTCACAGCTTCTATAGAAT  
AGTTACATCAGC  
CTTCTGAAGATGGCCTCTAGAGCACATGCACCCCCAAGATTCTAAGA  
10 TGTCAATACTAA  
CTGACCAAACCACCTCTAGCCAGCCCTGCTGCTCTGTTGCTGG  
TACCCAGGTGA  
CTGAGGACATGACTGGTGGAAAGGAAACTAGGCCCCTTGTCTGTCAGA  
TGGCCATACCCA  
15 GCATGGCTGATGCCAGTGTATAAGACCCCTACGCTTTCCACTGGTCTT  
AATGTTAAACC  
CTAGGACAGTGTCTCAGCATAGCTGGTGTGTGAATGCAAACATTG  
GGGCATATCTCT  
TCCATTAAGCACTGTGATATATGTAGTATTCCAACAAATAAATTATAC  
20 CTACATGATTG  
GGTATAGCATTCTGGATGGTCACAGGTGTGTCAAGGTGCTAATTAT  
GTGGGGGAAGAA  
CATAGAAATATATAGGTGGGGAGGGAGCTAACCCCTAGGAATAAGGCT  
AAAGCATGTGTCT  
25 CCAGTCCTGAAGACTCAAAGGGCAACGTGAATCATGAGACATGTCAG  
GAATGAGGAGT  
TGCCATGTATCTGCTTGTATCTTAATCATACATACACTATGAGA  
TCTGTGTTACC  
TCCATTTGCAGGTGAGAAAAGAAACACCTGAATGCCCTACCTAAAG  
30 GGCTAAGTGGGA  
AAATAGGTCTGAAGATAACCCAGGCAGTGTGACAAAGCAGGGAAAGA  
AAACTAGAGATGC

-31-

TTTCTTCATGGCAACAAACCTAGAGGGTACAACCTAGTGGTTCTTCTTG  
GTACTCCACTG  
TATACACCCCATCTGCTGGGCTGTACATTGTCTGACCATGCTTATAAC  
AAAAGTCACAT  
5 ACTACTAGCCAAGACTGAGAACTTAGAGCGACTGGCCAGAAAGTAAA  
GATACAACAGTTG  
ATATGTGTGCCACACACAGATCCATGTGTACATGTCTATTAAATTATGTG  
AACGTGCTTG  
TGGACATCCTCACAAAGCAGCAGGGAAATGCAAAGGTCAATTCCATAA  
10 CACCTGCTGGAC  
ACCATATGACATTGAGATTACCGGGGTGCCATTCCAACAAGAGTTAA  
TAGCTCCCCCTA  
TGTTTGGGTGCCAGAAACCTGATTGTTAGCAATAGCTCCCTCACATCC  
AGATTAAGAGG  
15 GGGATGGCTTAGCTAGGGTTACTATGATGAAACTATGACCAAAGCAAC  
TTGTGGTAAAAA  
GGGTGTATTGGCTTACACTCCATATCACTTCATCAAAGTGAGGACA  
GGAACCTCAAATA  
GAGTAGGAATTGGTGACAAGAGCTGATGTAGAGGCAATGCAGTGGT  
20 GCCACTTAGTGGC  
GCGCTCAGTCTGCTCCCTTCTTAATAGAATGCAAGACCAACAGCCAT  
GGGTGGCACCA  
CAATGGGACCGGGCCCTCCCATCGGTCACTAAGAAAATGCCCTACA  
GCCAGATCTTAT  
25 GGAGACATTTCTAACGGAGGCTCACTCCTTCAGATAACTCTATATC  
AAATTGACATA  
AACCAAGAACAGAGGAGGAGGCTAAGAAGGAAACTGCCAATTGCATAC  
ATGCACACACCTG  
GCCCTAGCAGCTGCAGGAAGCTATTGTTATGGCCTTTCTCATTTC  
30 ATGGACCAGCA  
TGAGCACTCTGCAGAGAGAGATGCCTGCATGCCTGCCAAGGCAGGAGT  
GCTTACACTGAA

-32-

GGTCAACAGGATGGCAGGGGGCTGCAGAGCTTCCAAGTGTCAAGAAC  
CCCAGCAGAAGAG  
CTGAGACCCCTGCCGAGGACTCAGGCAGGTTGGAAAGGCCAGGAAA  
TTCAGCCAGAGCT  
5. CTTCTTCAGATGGGTACCATCTGAAGGTTAGACCAGCTAGCCAGCTG  
TTGTTGAGGGAC  
CACCTCTGCAGCCCTACCTTGGAAAGATAGAAAGTGTCTGTGACA  
AGTATGGCCATT  
GTGCCCCCTTATTCCACAGTCAACAGAAACCTGGAATCCTGAACACT  
10 TCTGCAGCTTCT  
TTTTTACAGTCTGCCAGGTTGCTCTAGGAATGAAGGGTGCAGAGGC  
TTGGCGTAGGC  
AGGTGACAAGACCACAGTTAGTGGTCACAGCTGGCTTACTGGATCACT  
CTTGGACAGAGT  
15 TTGTTAGATATGGAGTGGAGTATAACACAAGGCATCAGGCGGGGATAT  
TGAATGTATCAC  
CGGAGCTCCTGGGGCTGGCAGCCAAGCACAGCAGTGGTTTGCTAA  
ACAAATCCACGG  
TTCCCTCCCTTGACGCAGTACATCTGTGGCTCCAACCCACACACCCA  
20 CCCATTGTTAG  
TGCTGGAGACTTCTACCTACCATGCCAGCTTGCTATCATGGGCTCA  
GCCTGGCTGCT  
TTCCTGGAGCTTGGATGGGGCCTCTTGTGTCTGTACAGCAATTCA  
AGGCACAAGGG  
25 GACTACATACTGGCGGGCTATTCCCTGGCTCAACCGAGGAGGCC  
ACTCTCAACCAG  
AGAACACAAACCCAACAGCATCCCGTGCAACAGGTATGGAGGCTAGTA  
GCTGGGGTGGGAG  
TGAACCGAAGCTTGGCAGCTTGGCTCCGTGGTACTACCAATCTGGGA  
30 AGAGGTGGTGAT  
CAGTTCCATGTGGCCTCAGGTCTCACCCCTGGTTTGTCTGGCCA  
TGGCTATGAAG

-33-

ATGGCTGTGGAGGAGATCAACAATGGATCTGCCTGCTCCCTGGGCTG  
CGGCTGGGCTAT  
GACCTATTGACACATGCTCCGAGCCAGTGGTACCATGAAATCCAGT  
CTCATGTTCTG  
5 GCCAAGGTGGCAGTCAAAGCATTGCTGCCTACTGCAACTACACACAG  
TACCAACCCCGT  
GTGCTGGCTGTCATCGGCCCCACTCATCAGAGCTGCCCTCATTACAG  
GCAAGTTCTC  
AGCTTCTTCATGCCACAGGTGAGCCCACCTCCTTGTGTTCTAAC  
10 CGATTGCACCC  
ATTGAGCTCTCATATCAGAAAGTGCCTCTTGATCACCAACAGGTAGCT  
ATAGTGCAGCA  
TGGATCGGCTAAGTGACCGGGAAACGTTCCATCCTCTCCGCACAG  
TGCCCAGTGACC  
15 GGGTGCAGCTGCAGGCAGTTGTGACTCTGTCAGAAACTCAGCTGGA  
ACTGGGTGGCCG  
CCTTAGGGAGTGATGACTATGGCCGGAAAGGTCTGAGCATCTTT  
CTAGTCTGGCCA  
ATGCACGAGGTATCTGCATGCACATGAGGGCCTGGTGCACAAACATG  
20 ACACTAGTGGCC  
AACAGTTGGCAAGGTGCTGGATGTACTACGCCAAGTGAACCAAAGT  
AAAGTACAAGTGG  
TGGTGCCTGCTGCTGCCGTGCTGTACTCCCTTTAGTTACAGC  
ATCCATCATG  
25 GCCTCTCACCAAGGTATGGGTGCCAGTGAGTCTGGCTGACATCTG  
ACCTGGTCATGA  
CACTTCCAATATTGCCGTGAGGCACGTGCTGGTTTGAGCG  
GGGTGCCCTAC  
TGCCTGAATTCTCCATTATGTGGAGACTCACCTGCCCTGGCCGCTGA  
30 CCCAGCATTCT  
GTGCCTCACTGAATGCGGAGTTGGATCTGGAGGAACATGTGATGGGC  
AACGCTGTCCAC

-34-

GGTGTGACGACATCATGCTGCAGAACCTATCATCTGGGCTGTTGCAGA  
ACCTATCAGCTG  
GGCAATTGCACCACCAAATATTGCAACCTATGCAGCTGTACAGTG  
TGGCTCAAGCCC  
5 TTCACAACACCCCTACAGTGCAATGTCTCACATTGCCACGTATCAGAAC  
ATGTTCTACCC  
GGCAGGTAAGGGTAGGGTTTTGCTGGGTTTGCCTGCTCCTGCAGG  
AACACTGAACCA  
GGCAGAGCCAAATCTGTTGTGACTGGAGAGGCCTTACCCCTGACTCCA  
10 CTCCACAGCTCC  
TGGAGAACATGTACAATATGAGTTCCATGCTCGAGACTTGACACTAC  
AGTTGATGCTG  
AAGGGAATGTAGACATGGAATATGACCTGAAGATGTGGGTGTGGCAG  
AGCCCTACACCTG  
15 TATTACATACTGTGGGCACCTTCAACGGCACCCCTCAGCTGCAGCAGT  
CTAAAATGTACT  
GGCCAGGCAACCAGGTAAGGACAAGACAGGCAAAAGGATGGTGGGT  
AGAACGCTTGTGCG  
TCTTGGGCCAGTGCTAGCCAAGGGAGGCCTAACCCAAGGCTCCATGT  
20 ACAGGTGCCAGT  
CTCCCAGTGTCCCCGCCAGTGCAAAGATGCCAGGTTGCCAGTAAA  
GGGCCTTCATTC  
CTGCTGCTATGACTGCGTGGACTGCAAGGCGGGCAGCTACCGGAAGCA  
TCCAGGTGAACC  
25 GTCTTCCCTAGACAGTCTGCACAGCCGGCTAGGGGGCAGAACGATTC  
AAGTCTGGCAAG  
CGCCCTCCCGGGGCTAATGTGGAGACAGTTACTGTGGGGCTGGCT  
GGGGAGGTGGT  
CTCCCAGCAGACCCCCACATTACTTTCTCCTTCCATCACTACAGA  
30 TGACTTCACCT  
GTACTCCATGTAACCAGGACCAGTGGTCCCCAGAGAAAAGCACAGCCT  
GCTTACCTCGCA

-35-

GGCCCAAGTTCTGGCTGGGGGGAGCCAGTTGTGCTGTCACTCCTCCT  
GCTGCTTGCC  
TGGTGCCTGGCTAGCACTGGCTGCTCTGGGCTCTGTCCACCACTG  
GGACAGCCCTC  
5 TTGTCCAGGCCTCAGGTGGCTCACAGTTCTGCTTGGCCTGATTCGCT  
AGGCCTCTTCT  
GCCTCAGTGTCCCTCTGTTCCCAGGGCGGCCAAGCTCTGCCAGCTGCCT  
TGCACAACAAC  
CAATGGCTCACCTCCCTCACAGGCTGCCTGAGCACACTCTTCCTGCA  
10 AGCAGCTGAGA  
CCTTTGTGGAGTCTGAGCTGCCACTGAGCTGGCAAACTGGCTATGCA  
GCTACCTTCGGG  
GACTCTGGGCCTGGCTAGTGGTACTGTTGCCACTTTGTGGAGGCAG  
CACTATGTGCCT  
15 GGTATTGATCGCTTCCCACCAAGAGGTGGTACAGACTGGTCAGTGC  
TGCCCACAGAGG  
TACTGGAGCACTGCCACGTGCGTCCCTGGTCAGCCTGGCTTGGTGC  
ACATCACCAATG  
CAATGTTAGCTTCCTCTGCTTCTGGCACTTCCTGGTACAGAGCCA  
20 GCCTGGCCGCT  
ACAACCGTGCCGTGGTCTCACCTCGCCATGCTAGCTTATTCATCAC  
CTGGGTCTCTT  
TTGTGCCCTCCTGGCCAATGTGCAGGTGGCTACCAGCCAGCTGTGC  
AGATGGGTGCTA  
25 TCCTAGTCTGTGCCCTGGCATTCTGGTCACCTCCACCTGCCAAGTG  
CTATGTGCTTC  
TTTGGCTGCCAAAGCTAACACCCAGGAGTTCTCCTGGAAAGGAATG  
CCAAGAAAGCAG  
CAGATGAGAACAGTGGCGGTGGTGGCAGCTCAGGGACACAATGAA  
30 TGACCACTGACCC  
GTGACCTTCCCTTAGGGAACCTAGCCCTACCAGAAATCTCCTAAGCC  
AACAAAGCCCCGA

-36-

ATAGTACCTCAGCCTGAGACGTGAGACACTTAACATAGACTTGGACT  
CCACTGACCTTA  
GCCTCACAGTGACCCCTCCCCAAACCCCCAAGGCCTGCAGTGCACAA  
GATGGACCCCTAT  
5 GAGCCCACCTATCCTTCAAAGCAAGATTATCCTGATCCTATTATGCC  
CACCTAAGGCC  
TGCCCAGGTGACCCACAAAAGGTTCTTGGGACTTCATAGCCATACTTT  
GAATTCAAGAAA  
TTCCCCAGGCAGACCATGGGAGACCAGAAGGTACTGCTGCCTGAACA  
10 TGCCCAGCCCTG  
AGCCCTCACTCAGCACCCCTGTCCAGGCGTCCCAGGAATAGAAGGCTGG  
GCATGTATGTGT  
GTGTGTGTGTGTGTGTGTGTGTGTGTGTATGTACGTATG  
TATGTATGTAT  
15 CAGGACAGAACAAAGAAAGACATCAGGCAGAGGACACTCAGGAGGTAG  
GCAACATCCAGCC  
TTCTCCATCCCTAGCTGAGCCCTAGCCTGTAGGAGAGAACCAAGGTCGC  
CGCCAGCACCTT  
GGACAGATCACACACAGGGTGGGGTCAGCACCAAGGCCAGGCCAG  
20 CCACGCGGGACCC  
CTGGAATCAGCTCTAGTACCAAGGACAGAAAAGTTGCCGCAAGGCC  
CTTACTGGCCAG  
CACCAAGGACAGAGCCACATGCCTAACGGCAAGGGACAAGAGCATC  
GTCCATCTGCAGG  
25 CAGGATCAGACCCGGGTCAAGTCTGGACTGGCCCCCACACCTGAATCC  
CGGAGCAGCTCA  
GCTGGAGAAAAGAGAAACAAGCCACACATCAGTCCCATAAAATTAAA  
CGCTTTTTAGT  
GTTAAAATAGCATTACACAGAAGCAGCATTACACAGAAGCAGCTC  
30 TATGTCAACTAC  
CCAGTCACTCAGACTTGCACACAGTGTCTAGTGTAGATGTGTGGGCC  
GCTGTGCCGGGA

-37-

TGGCAGTGGCACATGATGATGGGCAGCCACCAGAACAGAACAGAAC  
AGGGCCCAGCTCT  
GCAGCTCTGTGTTCACTGTCACCCACCACTGAGACTGAGACAGTGGC  
TAGGTGCCAGGT  
5 CTCTCTCCTGTCTCTCCTACTAGCTACCCCTCACATACCTCAGTACAA  
ACTGTGTTGTC  
ATGTGCCAAGTAGCAGGTGGGAAAGGGGCATGCAAACGTGCCCTTG  
GGTAACTAGCTG  
CCACCCCTAGAGCAGGCAGGCTAGCAATAAATAAATAAGTTAGACCCC  
10 ACCTGGGCAGCC  
AGAGAGGTTGAAGGCTCTGCTAACCCCTAAAAATCCCACCTTGGC  
CTGACAGGTGAG  
GCCCATGAACCTAGCGACAGTCAGCCTGTGTCCTGTGCACAGTTCTGT  
GAGGCTTGAG  
15 GCAAGGGGTACCAAGAGGCCAAGAGAGGCCTTCTTGTCTAAATGGAG  
GTCACCTCCAAA  
GAAGGGAACCAGGAGGTGGTCCCTGAGACTTGTGCTGAGGACTTAAA  
GTCAGAGATGTCT  
CCTTACAAGACTCTAGATACTTGAGCTGTACCACCATCAGCAGCCC  
20 CAAGAGCAGACA  
AAATGTCAAGCCAATATCCTGGTGGTATGGCTGCCCTCAGGCCCTCCT  
CTGTAGCCTGCT  
CCCTCTGCCCTGGCCCAGAGCCCACAGCTGATCTATCCTGGCTGCCA  
CCACCACGGCCA  
25 GCGCAGAGCTCCTGGCACAGCAGGAGCACAGACTCAGCCACAGGCAG  
CGCTGAAGACATT  
GGTGATCATCACATGATGTCCACAAAGAACTCACAGGGTTCCCAT  
GGCCTTTGGAA  
GGACTGGCGCTACCTGTAAGTTCTGGAGGGACAGCAGCCAGCTCCCG  
30 GACGGGTGGCCC  
TCCAGGTGGCCCACCCACTACTGCATAGGCCTTGTAAAGGGGTGCAG  
TGGGGGGAGCCC

-38-

TGGGGCAACAGCTGAAGCCTGACTTCGAGGGCTACTGCCACGGCTAAG  
CTGGCTGACAGG  
CCGCTCCCACCAGCCGGTGCTACCAGACCCACTGGTACTGTGTGGTCT  
GATTCACTGCC  
5 ACTACCCCCAGCTCCAGTTGCCCGGCGCTCCTCTCGGCCTGGGTCCG  
ATGGCTGCTCCG  
TGTGGACCCACTGCTCTGCTCCCTAGGGGGAGGGAAAGGGGACAACAG  
AGTCAGCACGAG  
GCCTGGCCACTCCAGGGCCACCAGCTGCTCCAGACAGTCAGGGCAG  
10 GACCTGGTAAGC  
CTGGAGATGGTAGGGGAATGGCAGCCATGCAGATAACCAGGAACAGCT  
GAGAGGCGAGAAG  
CTAGGGGCAGTGGCAGACAGCAGGGACAACAGGGCCAGCCTGGCAC  
CCCACACCTAAC  
15 CCAATGCTTGAACCAAGGGTTAATGTTACAGCTGAGAAACTAAAAACC  
AGCGAAGGCCCT  
GTGTGCCAGCATTCCCATTAGCCATCCTGGGTTCAACCACCCAAAGAC  
CCAACCAGGGTC  
CACCCAACCCCAGGACCCTGGTCATCTAATTGCTTAGCCCCTGTCCTG  
20 AAAGTAGTGGG  
AACCTGAAAACACGTGCTGGCTGGGACATGCTGAGAGGGACACAGG  
GGGACCTGGCTTA  
CCGGCCCGAGAGTCCACTCTGCTAGTCCTCAGTCTAAGGCTTGTCAG  
CACAAAGCAAG  
25 GGATAGCACAAGTCACACACCAGTCCAGTGCTCACCAATGGCTAATAG  
GACGATTTGGG  
CCAAGCTGAGCCTGGGTACATGCAAGGGCTGTCCATGGTCAGGATT  
ACTCGATAGCTT  
CCCCTGGGCTTGCACCCCTGGCCAAACCTCTCCTGAGTCTTCTCT  
30 GGACCTTGTA  
GCACAAGTGTGCCCCACTCTGCCTAAGACCTCCACATCAGTCCATCTCC  
TCCTGAGGGAC

-39-

ACCCACCCCTCAAGATCTTCAATATCCCTGGGATATGCTTAACACTGA  
TATGCTTAAC  
AGTGTGCTTGATACTCTTATCTGGCACTCTGTTGGATGCAGGCTCCA  
TAACTGATAAA  
5 GCCCATTCTCCCCTAGCTGGGCCTAGAGAGTGCCCCTACCTGCTAT  
CAGTGGTTACT  
TTCATTCTGCCATATCATCTCCTGGCCTCTGCCCTGCCACCTAGCAC  
ACCAGGCTGT  
CTTCCTATTCTCTAACGGCTTCTACCCACATCAGCCCCTCCCTGTCCCA  
10 CACACTGACTC  
TTGAGATGGAACCCACCGGGACTCAAACACACAGCAGGAGCACAGAG  
GGAAGCGTGGGG  
CCAGGCAGAGCGTGGGAGTGGGAGGGAGTGGGAGGGAGGGTGGCAC  
GCCTCTCACCTCA  
15 CTCTGCTGGCTCCCAGCACTGCCGTGCCAGCTGAAGCCAGGGTCC  
TGGTAAGCAGGC  
GGGAAGCAGGGCGGGGTCCTGGTACTGGTAGGGTAGCCTTGACC  
CAAGGGCCAGGGT  
ACTGATGGGTGGGCAGTGGGCCAGTGTGTCCCTGATCTGAGGCTCCA  
20 CTGGAGCCACTG  
TTGAGGTTCAAGGATGCGAGGTCTGGCAGGGAGGGAGGGAGGGAGGG  
GTAAGTGAAGGCA  
AATGAATGAGGCCACAGCAACCCCTACCCAACCGCACCCCTACTCACTA  
CTGCACAGGTG  
25 CCAAAGACATAGTAGCACTGCTCAGAAAAGGTGATCTTGTTCACGGTG  
TGCCTCAGGAAA  
CCGTGCTTCAGCATACTGCTGGCATACTTCTGCCTCCCTCGCTCCTT  
GAAGCCCTCC  
ACGTGTGTACAGCCAGTCCACCACATCCGCCCTGGCCACAGGTCC  
30 ATCAAAGTCAGG  
GTAGCTGAGCCCTGGGAAGCTACGCCAGAACATGAGGAACAGACGGGGC  
CCTTCCCCACACAG

-40-

CCAGGGACTCACCAATGACAGCATTGCAATGGTATCTTAAGCCACA  
TGCGGTCCCAGA  
TCTCCAGTCTGAGTCTGGCAACTGCATGACGCCGACAATGGCACTCA  
TGTCACTCTTCA  
5 CAGTCAGCGGTGCCTCCTCAAGCTCTGCAGAGCACACTCCCTGAGCC  
CAGGCTCACAGC  
GTGAACCTCCATGGGGTTGAGAGCAGGGGCCAGGGTCAAACCTCTTAT  
CTCCCATCCTTG  
GGAGATGCCCTCATCGAAACTTGAGCTAAGACCGGGAGATTCTTCCC  
10 CGTCCCACAGTG  
CAAGTCCACGTAGGCAAGGCAGCCCCCTCCCCCTCCCCGGAGAGAAC  
AGCTGTTAGCTA  
TGTTAGGTAGCAGAAAAGCAAAGCAGAGGCTGCCATGTCCTCCAAATT  
CCCCCCTCCGCA  
15 CAGGCCTGGCAGGACCCCTCAATTGAGATGACAGTATGGCCAGG  
CCTGGAGGGATA  
TGTACATGTATCTTGTGTACACATTGTGAAGGTGTTGGAAGCAAAC  
AAAACCTTCATA  
TGTAATGGGCCCTGTAATAGCTCTGATGAGCACCAAGCTCAAAGCT  
20 AGAACTGACCAT  
TGTCTTCAACCTCAGTTCTGGTGGGGGGGGGTCTGTGAGCTGC  
CACTTACGTGG  
GGCGCCAGGCAGTGGTAGTGAGGAAGAGCTGGTGCCTGTGAT  
GGCGCTGGAGCA  
25 GGGACTCGTACCATAGCGGGCAGGGCACCCGTCACTGCTGCTGTG  
GGACAGCCAGGC  
AGCCGGGTGATGGTCGCACTGGTCAGCTGCATAGTTCCACAGCA  
ACGGATTACAGG  
TGGTAAGTAGGGGGCAGCACAGAGGCAGACAAGAAAGACCCCCAGA  
30 CTGAACACAGAAA  
CCCCACCCCTACCCCCACCTTCCATGGGTAACTCACCCCTGGGATGGT  
GAAGTAGCTCC

-41-

GAGGGGTTGGGTCCCAGCACTGGCCACTGTGAGACTGATGGGCCTAC  
AGAGTTGAGCAG  
ACCATGTTGTAAGTGAGGCCCGCACAGCCCCTCCCATCCTGTGCCACT  
CCCACCCCCACT  
5 TGGCTCCCACCTCACCCCTGTCTGGGACACGATCTCCCGAAGCACCCGT  
ACAGCGTCGTCA  
TTGCTCATGTTCTCAAAGTTGACATCGTTCACCTACGGGGTTGTGGGG  
TCAGGGGTTGG  
TGGTGGGATGTGGGTGCCTCTGTCCCCACAGTCCCCACATGGCTCCA  
10 CCTGCAGCAAC  
ATGTCGCCCCGGCTCAATGCGGCCATCAGCAGCCACGGCCCCGCCCTTC  
ATGATGGATCCA  
ATGTAGATGCCGCCATCACCCCGGTCGGCTCTGGCCACGATGCTG  
ATGCCAGGAAG  
15 TGGTGCCTCTGCAGGAGGGGCCGTGAGCAGGCCCCAAAGCTCCCG  
AGGCTGTACCCA  
CCCCCAGCAGGCACCCACAGCCCACAAGGCCTCACCCATGTTGAGAGT  
GACGGTGATGAT  
GTTCAAGGGACATGGTGGAGTCTGTGATGCTGCTGAAGGAGGATGCCTG  
20 CGGAGGGACCCA  
GTGAGGGGCTGTGTGGCACCATTAGAGCAGACACCCCCACCCACCTG  
CTGCCTACCCGG  
TCTGTCTGCCTCAAGCGCTGCTCCGACGACGGCATTGTGCTCCGAA  
CTAGCCGAGAG  
25 GAGGTGCTCTGCTCTGGAGCTGCTCAGCCTGAGGCAGGAGTCAGAA  
AAGCACAAACAT  
GTATAACCAGCTGGACGCTCAACTACAAATCTCCAGCACGTACTGAC  
ATGTGCACACGT  
CACCCACCGGCTCGTATTGTCCTCCTCATCTGAGTCATAAAAGCTGCTA  
30 GATTCAAGCTC  
ACTGCTCAGTACAGTGGATGCACTGTCTGGAGGTAGTCCCAGGTCCCG  
CCGCCGATCCCC

-42-

TCTCGGGTGCCATTGGTCCGGGCAGCTGTGGGGACAGTAGGGTGGGT  
 ACGACTGTGGGA  
 CTTCAGTCCTAACAGAACATGCAGGTGGCCTGTGCATTCAAAGTTATGC  
 AGTAACCTCTGG  
 5   GGCACAGGGCTAGGAGTACCAAGGCTGGACCTCTACCCAAAGGATC  
 ACTGCTTGGAAAGA  
 ATATGTGGAATACTCCAGGCTGGAGTATACCAAAGGGATACCAAG  
 GG

The polypeptide sequence of mouse SAC1 (SEQ ID NO: 3) is:

10   MPALAIMGLSAAFLLELGMGASLCLSQQFKAQGDYILGGLFPLGSTEEAT  
 LNQRTQPNSIPCNRFSPGLFLAMAMKMAVEEINNGSALLPGLRLGYDLF  
 DTCSEPVVTMKSSLMFLAKVGSQSIAAYCNYTQYQPRVLAIGPHSSELA  
 LITGKFFSFFLMPQVSYSASMDRLSDRETFPSFFRTVPSDRVQLQAVVTLL  
 QNFSWNWVAALGSDDDYGREGLSIFSSLANARGICIAHEGLVPQHDTSGQ  
 15   QLGVLDVLRQVNQSKVQVVVLFASARAVYSLFSYSIHGLSPKVVVAS  
 ESWLTSVLVMTLPNIARVGTVLGFLQRGALLPEFSHYVETHLALAADPAF  
 CASLNAELDLEEHVMGQRCPRCDDIMLQNLSSGLLQNLSAQLHHQIFAT  
 YAAVYSVAQALHNTLQCNVSHCHVSEHVPWQLENMYNMSFHARDLT  
 LQFDAEGNVDMEYDLKMWVVWQSPTVLHTVGTNGTLQLQQSKMYWP  
 20   GNQVPVSQCSRQCKDGQVRRVKGFHSNCYDCVDCKAGSYRKHPDDFTC  
 TPCNQDQWSPEKSTA CLPRPKFLAWGEPVVLSSLLLCLVGLALAALG  
 LSVHHWDSPLVQASGGSQFCFGICLGLFCLSVLLFPGRPSSASCLAQQPM  
 AHLPLTGCLSTLFLQAAETFVESELPLSWANWLCSYLRGLWAWLVVLLA  
 TFVEAACAWYLIAPPEVTDWSVLPTEVLEHCHVRSWVSLGLVHITNA  
 25   MLAFLCFLGTFLVQSQPGRYNRARGLTFAMLAYFITWVSVPLLAVQV  
 AYQPAVQMGAILVCALGILVTFLPKCYVLLWLPKLNTQEFFLGRNAKK  
 AADENSGGGEAAQGHNE

The cDNA of human SAC1 (SEQ ID NO: 4) is:

ATGCTGGGCCCTGCTGTCCTGGGCCTCAGCCTCTGGCTCTCCTGCACC  
 30   CTGGGACGGGGGCCATTGTGCCTGTCACAGCAACTAGGATGAAGG

-43-

GGGACTACGTGCTGGGGGGCTGTTCCCCCTGGCGAGGCCGAGGAG  
GCTGGCCTCCGCAGCCGGACACGGCCCAGCAGCCCTGTGTGCACCAGG  
TTCTCCTCAAACGGCTGCTCTGGCACTGGCCATGAAAATGGCCGTG  
GAGGAGATCAACAACAAGTCGGATCTGCTGCCCGGGCTGCGCCTGGC  
5 TACGACCTCTTGATACGTGCTCGGAGCCTGTGGTGGCCATGAAGCCC  
AGCCTCATGTTCTGGCCAAGGCAGGCAGCCGCGACATGCCGCCTAC  
TGCAACTACACGCAGTACCAAGCCCCGTGTGCTGGCTGTATGGGCCC  
CACTCGTCAGAGCTGCCATGGTACCCGGCAAGTTCTCAGCTTCTCC  
TCATGCCCGAGGTCAAGCTACGGTCTAGCATGGAGCTGCTGAGCGCCC  
10 GGGAGACCTTCCCCCTCCTTCTTCCGCACCGTGCCAGCGACCGTGTGCA  
GCTGACGGCCGCCGGAGCTGCTGCAGGAGTTCGGCTGGAACGGT  
GGCCGCCCTGGCAGCGACGAGTACGGCCGGCAGGGCCTGAGCA  
TCTTCTCGGCCCTGGCCTCGGCACGCCGCATCTGCATCGCGCACGAGG  
GCCTGGTGCCGCTGCCCGTGCCATGACTCGCGCTGGGAAGGTGC  
15 AGGACGTCTGCACCAGGTGAACCAGAGCAGCGTGCAGGTGGTGTGCTG  
CTGTTGCCTCCGTGCACGCCGCCACGCCCTTCAACTACAGCATCA  
GCAGCAGGCTCTGCCCAAGGTGTGGTGGCCAGCGAGGCCTGGCTGA  
CCTCTGACCTGGTCATGGGCTGCCCGATGCCAGATGGCACGG  
TGCTTGGCTCCTCCAGAGGGGTGCCAGCTGCACGAGTTCCCCAGT  
20 ACGTGAAGACGCACCTGGCCCTGGCCACCGACCCGGCTCTGCTCTG  
CCCTGGCGAGAGGGAGCAGGGTCTGGAGGAGGACGTGGTGGCCAG  
CGCTGCCCGCAGTGTACTGCATCACGCTGCAGAACGTGAGCGCAGGG  
CTAAATCACCACCAGACGTTCTGTCTACGCAGCTGTATAGCGTG  
GCCCAAGGCCCTGCACAACACTCTTCAGTGCACGCCCTCAGGCTGCC  
25 GCGCAGGACCCCGTGAAGCCCTGGCAGCTCCTGGAGAACATGTACAAC  
CTGACCTTCCACGTGGCGGGCTGCCGCTGCCGTTGACAGCAGCGGA  
AACGTGGACATGGAGTACGACCTGAAGCTGTGGTGTGGCAGGGCTC  
AGTGCCCAAGGCTCCACGACGTGGCAGGTTAACGGCAGCCTCAGGAC  
AGAGCGCCTGAAGATCCGCTGGCACACGTCTGACAACCAGAACGCCGT  
30 GTCCCGGTGCTCGCGGAGTGCAGGAGGGCCAGGTGCGCCGGGTCA  
AGGGGTTCCACTCCTGCTGCTACGACTGTGTGGACTGCGAGGCAGGGCA  
GCTACCGGAAAACCCAGACGACATGCCCTGCACCTTGTGGCCAGG

-44-

ATGAGTGGTCCCCGGAGCGAACACGCTGCTCCGCCGCAGGTCTC  
 GGTTCTGGCATGGGCGAGCCGGCTGTGCTGCTGCTGCTCCTGCTGCT  
 GAGCCTGGCGCTGGCCTTGTGCTGGCTGCTTGGGCTGTTGTTAC  
 CATCGGGACAGCCCAGTGGTCAGGCCTCGGGGGGCCCTGGCCTGC  
 5 TTTGGCCTGGTGTGCCTGGCCTGGTCTGCCTCAGCGTCCCTGTTCC  
 CTGGCCAGCCCAGCCCTGCCGATGCCTGGCCAGCAGCCCTGTC  
 ACCTCCCGCTCACGGCTGCCTGAGCACACTCTCCTGCAGGCGGCCG  
 AGATCTTCGTGGAGTCAGAACTGCCTCTGAGCTGGCAGACCGGCTGA  
 GTGGCTGCCTGCAGGGGCCCTGGCCTGGCTGGTGGTGTGCTGGCCA  
 10 TGCTGGTGGAGGTGCACTGTGCACCTGGTACCTGGTGGCCTCCCGC  
 CGGAGGTGGTACGGACTGGCACATGCTGCCACGGAGGCGCTGGTG  
 CACTGCCGCACACGCTCCTGGTCAGCTCAGGCCCTAGCGCACGCCACC  
 AATGCCACGCTGGCCTTCTCTGCTTCCCTGGCACCTTCTGGTGC  
 GCCAGCCGGGCCGCTACAACCGTGCCGTGGCCTCACCTTGCCATGC  
 15 TGGCCTACTTCATCACCTGGCTCCTTGTGCCCTCCTGGCCAATGT  
 GCAGGTGGTCCTCAGGCCCGCCGTGCAGATGGCGCCCTGCTCTG  
 TGTCCCTGGCATTCTGGCTGCCCTCACCTGCCAGGTGTTACCTGCTC  
 ATGCGGCAGCCAGGGCTAACACCCCCGAGTTCTCCTGGGAGGGGGC  
 CCTGGGATGCCAAGGCCAGAATGACGGAACACAGGAAATCAGGG  
 20 GAAACATGAGTGA

The polypeptide sequence of human SAC1 substantially from the translated region of the human cDNA (SEQ ID NO: 5) is:

MLGPAVLGLSLWALLHPGTGAPLCLSQQLRMKGDYVLGGLFPLGEAEEA  
 GLRSRTRPSSPVCTRFSNGLLWALAMKMAVEEINNKSDLPLGLRLGYDL  
 25 FDTCEPVVAMKPSLMFLAKAGSRDIAAYCNYTQYQPRVLAIGPHSSEL  
 AMVTGKFFSFFLMPQVSYGASMELLSARETFPSFFRTVPSDRVQLTAAAE  
 LLQEFGWNWVAALGSDDEYGRQGLSIFSALASARGICIAHEGLVPLPRAD  
 DSRLGKVQDVLHQVNQSSVQVVLFAJVAAHALFNYSISSRLSPKVVW  
 ASEAWLTSSDLVMGLPGMAQMGTVLGFLQRGAQLHEFPQYVKTHLALAT  
 30 DPAFC SALGEREQGLEEDVVGQRCPQC DCITLQNV SAGLNHHQTF SVYAA  
 VYSVAQALHNTLQCNASGCPAQDPVKPWQLENMYNLTFHVGGLPLRF

-45-

DSSGNVDMYEYDLKLWWQGSVPRLHDVGRFNGSLRTERLKIRWHTSDN  
 QKPVSRCSRQCQEGQVRRVKGFHSCCYDCVDCEAGSYRQNPDDIACTFC  
 GQDEWSPERSTRCFRRRSRFLAWGEPAVLLLLLSSLALGLVLAALGLFV  
 HHRDSPLVQASGGPLACFGLVCLGLVCLSVLLFPQSPARCLAQQPLSHL  
 5 PLTGCLSTLFLQAAEIFVESELPLSWADRLSGCLRGPAWLVVLLAMLVE  
 VALCTWYLVAFPPEVVTDWHMLPTEALVHCRTRSWVSFGLAHATNATL  
 AFLCFLGTFLVRSQPGRYNRARGLTFAMLAYFITWVSFVPLLAVQVVLR  
 PAVQMGALLLCVLGILAAFHLPRCYLLMRQPGLNTPEFFLGGGGPGDAQG  
 QNDGNTGNQGKHE

10 III. SAC1 Is a G-Protein Coupled Receptor

The evidence that SAC is a G-protein coupled receptor (GPCR) comes from its sequence homology to other GPCR and the structure predicted for the amino acid sequence.

15 GPCRs (also known as 7-transmembrane receptors) bind extracellular ligands and transduce signals into the cell by coupling to intracellular G-proteins. GPCRs can be subdivided into more than 30 families on the basis of their ligands. Sac is most closely allied by sequence homology with the  $\text{Ca}^{++}$ -sensing, metabotropic receptors.

20 Proteins often contain several modules or domains, each with a distinct evolutionary origin and function. When the Sac cDNA sequence is queried against the Conserved Domain Database at NCBI, the following results are obtained:

Sequences producing significant alignments:		Score (bits)	E Value
Gnl   Pfam   pfam01094	ANF_receptor, Receptor family ligand binding region	145	73-36
Gnl   Pfam   pfam00003	7tm_3, 7-transmembrane receptor (metabotropic glutamate family)	87.0	3e-18

Note the ANF\_receptor family contains the metabotropic and calcium-sensing families of GCPs.

-46-

The closest sequence homology of the mouse SAC gene is to the  $\text{Ca}^{++}$  sensing receptors, all of which are GPCRs. An alignment between a calcium sensing GPCR (BAA09453) is shown in Fig. 5.

As described above, all GPCRs have a characteristic 7-transmembrane 5 domain. Figure 6 is a plot of the transmembrane domains of SAC1.

Table 1: Genes Predicted From the *Sac* Nonrecombinant Interval and Expression Data From NCBI

N	Gene or EST	Size (aa)	How Many ESTs From Tongue?	Suggested Protein Function
1	<i>Cyclin ania 6a</i>	~425	0/36	Potentially involved in differentiation and neural plasticity
2	<i>Slm1</i>	~189	0/29	Slm-1 is a Src substrate during mitosis
3	AA404005	446	0/61	Expressed in kidney
4	<i>Disheveled</i>	769	0/6	Segment polarity gene; knockouts have a behavioral phenotype
5	<i>Sac</i>	746 <sup>1</sup>	0/0 <sup>2</sup>	Sweet receptor
6	Mm.255556	216	0/5	Weakly similar to <i>Physcomitrella patens</i> glyceraldehyde 3-phosphate dehydrogenase in <i>C. elegans</i>
7	Mm.135238	524	0/5	Expressed in mammary gland and spleen
8	AA435261	328	0/1	Expressed in mouse two cell
9	Centaurin beta 2#	791	0/1	Regulators of membrane traffic and the actin cytoskeleton
10	Voltage gated Na <sup>+</sup> channel #	170	0/0	Gumarin reduces the perception of sweet, and may work by blocking sodium channels (Fletcher J.I., Chapman B.E., Mackay J.P., Howden M.E., and King G.F. The structure of versutoxin (delta-altracotoxin-Hv1) provides insights into the binding of site 3 neurotoxins to the voltage-gated sodium channel. <i>Structure</i> , 1997;5:1525-1535)
11	<i>Ubcc6p</i>	597	0/32	Essential for the degradation of misfolded and regulated proteins in the endoplasmic reticulum lumen and membrane
12	Mm.29140	402	0/2	Weakly similar to collagen alpha 1(XVIII) chain

The genomic sequence from AF183591 and RPCI-23-118E21, between the markers that flank the *Sac* nonrecombinant interval, was identified. The repetitive and low complexity sequences were removed, using Repeatmasker (Smit F. and Green P. Repeatmasker). The resulting sequence was analyzed by GENSCAN, which predicted 12 proteins. Of these 12 predicted proteins, one GENSCAN predicted protein was a chimera between two genes (cyclin ania 6a and Slm1). (These sequences were separated into their respective sequences.) The predicted proteins were submitted to a TBLASTN search through the *nr* and the *mouse* EST database at NCBI. Of the 12 predicted proteins, four were named genes, two genes were similar to other named genes (Centaurin beta 2 and the voltage gated Na<sup>+</sup> channel) and are denoted with an #. Three of the predicted proteins were represented as ESTs, and had Unigene cluster numbers. The remaining two predicted genes were identical to previously isolated mouse ESTs. When each predicted protein was blasted against the *mouse EST* database, the number of ESTs from tongue were compared with the number from other tissue sources. No ESTs from these genes appeared in the *mouse EST* database at NCBI.

- 1 Note that the GENSCAN prediction is not accurate; sequencing of the cDNA indicates *Sac* is 858 aa.
- 2 Note that TR1-like is expressed in tongue as detected by RT-PCR. Previously named genes are in *italics*, and ESTs or EST clusters in plain text.

#### IV. The *Sac* Locus and the *Gpr98* Sweet Taste Receptor Gene

A substantial effort has been devoted to positional cloning of a locus on distal Chr 4 with a major effect on sweetener intake. This locus has been previously described as the *Sac* (saccharin preference) locus, and it also explains ~8 % of the phenotypic variance in ethanol preferences within the B6 x 129 F<sub>2</sub> generation.

5 Details on positional cloning of the *Sac* locus are found above.

10 The effects of SAC1 (*Gpr98*) on ethanol intake Two lines of evidence point to the involvement of *Gpr98* in ethanol intake. First, 129.B6-*Sac* congenic mice homozygous for a 194-kb donor fragment from the B6 strain consumed more 10% ethanol solution than did congenic mice without the donor fragment (1.50 ± 0.15 and 1.19 ± 0.11 mL/day, respectively; *p* <0.05, one-tailed *t*-test).

15 Second, ethanol preference was related to sequence variations of *Gpr98*. Analysis of *Gpr98* sequences from genealogically remote or unrelated mouse strains indicated the presence of two haplotypes of single nucleotide polymorphisms within the *Gpr98* locus. One, 'B6-like' haplotype, was found in mouse strains with elevated sweetener preference and the other, '129-like' haplotype, was found in strains relatively indifferent to sweeteners as described above. Preferences for 10% ethanol for the same mouse strains were studied as described in Abstr. of the 20 23th RSA Meeting (June 2000, Denver, Colorado). We found that strains with the 'B6-like' haplotype had higher preferences for 10% ethanol (20 ± 4%, *n* = 14, strains C57BL/6J, C57L/J, CAST, FVB/NJ, KK/HIJ, NOD/LtJ, NZB/B1NJ, P/J, RBF/DnJ, RF/J, SEA/GnJ, SJL/J, SPRET/Ei and SWR/J) compared with strains having the '129-like' haplotype (12 ± 2%, *n* = 10, *p* <0.05, one-tailed *t*-test, strains 25 129P3/J, AKR/J, BALB/c, BUB/BnJ, C3H/HeJ, CBA/J, DBA/2J, LP/J, PL/J and RIIS/J).

#### V. Preparation of Recombinant or Chemically Synthesized Nucleic Acids, Vectors, Transformation, Host-Cells

30 Large amounts of the polynucleotides of the present invention may be produced by replication in a suitable host cell. Natural or synthetic polynucleotide

-49-

fragments coding for a desired fragment will be incorporated into recombinant polynucleotide constructs, usually DNA constructs, capable of introduction into and replication in a prokaryotic or eukaryotic cell. Usually the polynucleotide constructs will be suitable for replication in a unicellular host, such as yeast or bacteria, but may also be intended for introduction to (with and without integration within the genome) cultured mammalian or plant or other eukaryotic cell lines. The purification of nucleic acids produced by the methods of the present invention is described, e.g., in Ausubel et al., *Current Protocols in Molecular Biology*, Vol. 1-2, John Wiley & Sons, 1992 and Sambrook et al., *Molecular Cloning A Laboratory Manual*, 2nd Ed., Vols. 1-3, Cold Springs Harbor Press, 1989.

The polynucleotides of the present invention may also be produced by chemical synthesis, e.g., by the phosphoramidite method or the triester method, and may be performed on commercial, automated oligonucleotide synthesizers. A double-stranded fragment may be obtained from the single-stranded product of chemical synthesis either by synthesizing the complementary strand and annealing the strands together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polynucleotide constructs prepared for introduction into a prokaryotic or eukaryotic host may comprise a replication system recognized by the host, including the intended polynucleotide fragment encoding the desired polypeptide, and will preferably also include transcription and translational initiation regulatory sequences operably linked to the polypeptide encoding segment. Expression vectors may include, for example, an origin of replication or autonomously replicating sequence (ARS) and expression control sequences, a promoter, an enhancer and necessary processing information sites, such as ribosome-binding sites, RNA splice sites, polyadenylation sites, transcriptional terminator sequences, and mRNA stabilizing sequences. Secretion signals may also be included where appropriate, whether from a native SAC1 protein or from other receptors or from secreted polypeptides of the same or related species, which allow the protein to cross and/or lodge in cell membranes, and thus attain its functional topology, or be secreted from the cell. Such vectors may be prepared by

-50-

means of standard recombinant techniques well-known in the art and discussed, for example, in Sambrook et al., 1989 or Ausubel et al., 1992.

An appropriate promoter and other necessary vector sequences will be selected so as to be functional in the host, and may include, when appropriate, 5 those naturally associated with SAC1 genes. Examples of workable combinations of cell lines and expression vectors are described in Sambrook et al., 1989 or Ausubel et al., 1992. Many useful vectors are known in the art and may be obtained from commercial vendors. Promoters such as the trp, lac and phage promoters, tRNA promoters and glycolytic enzyme promoters may be used in 10 prokaryotic hosts. Useful yeast promoters include promoter regions for metallothionein, 3-phosphoglycerate kinase or other glycolytic enzymes such as enolase or glyceraldehyde-3-phosphate dehydrogenase, enzymes responsible for maltose and galactose utilization, and others. In addition, the construct may be joined to an amplifiable gene so that multiple copies of the gene may be made. For 15 appropriate enhancer and other expression control sequences, see also *Enhancers and Eukaryotic Gene Expression*, New York: Cold Spring Harbor Press, 1983. See also, e.g., US Patent Nos. 5,691,198; 5,735,500; 5,747,469 and 5,436,146.

Expression and cloning vectors will likely contain a selectable marker, a 20 gene encoding a protein necessary for survival or growth of a host cell transformed with the vector. The presence of this gene ensures growth of only those host cells which express the inserts. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxic substances, e.g., ampicillin, neomycin, methotrexate, etc.; (b) complement auxotrophic deficiencies; or 25 (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well-known in the art.

The vectors containing the nucleic acids of interest can be transcribed 30 in vitro, and the resulting RNA introduced into the host cell by well-known methods, e.g., by injection, or the vectors can be introduced directly into host cells by methods well-known in the art, which vary depending on the type of cellular host, including electroporation; transfection employing calcium chloride,

-51-

rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection; infection (where the vector is an infectious agent, such as a retroviral genome); and other methods. The introduction of the polynucleotides into the host cell by any method known in the art, including, inter alia, those described above, will be referred to herein as "transformation." The cells into which have been introduced nucleic acids described above are meant to also include the progeny of such cells.

Large quantities of the nucleic acids and polypeptides of the present invention may be prepared by expressing the SAC1 nucleic acids or portions thereof in vectors or other expression vehicles in compatible prokaryotic or eukaryotic host cells. The most commonly used prokaryotic hosts are strains of *Escherichia coli*, although other prokaryotes, such as *Bacillus subtilis* or *Pseudomonas* may also be used.

Mammalian or other eukaryotic host cells, such as those of yeast, filamentous fungi, plant, insect, or amphibian or avian species, may also be useful for production of the proteins of the present invention. Propagation of mammalian cells in culture is per se well-known. Examples of commonly used mammalian host cell lines are VERO and HeLa cells, Chinese hamster ovary (CHO) cells, and W138, BHK, and COS cell lines. An example of a commonly used insect cell line is SF9. However, it will be appreciated by the skilled practitioner that other cell lines may be appropriate, e.g., to provide higher expression, desirable glycosylation patterns, or other features.

Clones are selected by using markers depending on the mode of the vector construction. The marker may be on the same or a different DNA molecule, preferably the same DNA molecule. In prokaryotic hosts, the transformant may be selected, e.g., by resistance to ampicillin, tetracycline or other antibiotics. Production of a particular product based on temperature sensitivity may also serve as an appropriate marker.

#### VI. Diagnosis or Screening

Genetic analysis of obesity and diabetes and alcoholism or alcohol consumption is often complicated by the lack of a simple diagnostic mark. For

-52-

example, currently there is no single diagnostic marker for the diagnosis of obesity. Sequence variation of the SAC1 locus may indicate a predisposition to diabetes, obesity, and alcoholism and may provide a diagnostic mark.

5 In order to detect the presence of a SAC1 allele predisposing an individual to obesity, diabetes, or alcoholism, a biological sample may be prepared and analyzed for the presence or absence of susceptibility alleles of SAC1. Results of these tests and interpretive information may be returned to the health care professionals for communication to the tested individual. Such diagnoses may be performed by diagnostic laboratories. In addition, diagnostic kits may be  
10 manufactured and available to health care providers or to private individuals for self-diagnosis.

A basic format for sequence or expression analysis is finding sequences in DNA or RNA extracted from affected family members which create abnormal SAC1 gene products or abnormal levels of SAC1 gene product. The diagnostic or 15 screening method may involve amplification or molecular cloning of the relevant SAC1 sequences. For example, PCR based amplification may be used. Once amplified, the resulting nucleic acid can be sequenced or used as a substrate for DNA probes. Primers and probes specific for the SAC1 gene sequences may be used to identify SAC1 alleles.

20 The pairs of single-stranded DNA primers can be annealed to sequences within or surrounding the SAC1 gene in order to prime amplifying DNA synthesis of the SAC1 gene itself. The set of primers may allow synthesis of both intron and exon sequences. Allele-specific primers can also be used. Such primers anneal only to particular SAC1 mutant alleles, and thus will only amplify a product in the 25 presence of the mutant allele as a template.

In order to facilitate subsequent cloning of amplified sequences, primers 30 may have restriction enzyme site sequences appended to their 5' ends. Thus, all nucleotides of the primers are derived from SAC1 sequences or sequences adjacent to SAC1, except for the few nucleotides necessary to form a restriction enzyme site. Such enzymes and sites are well-known in the art. The primers themselves can be synthesized using techniques which are well-known in the art.

-53-

Generally, the primers can be made using oligonucleotide synthesizers which are commercially available.

The biological sample to be analyzed, such as blood, may be treated, if desired, to extract the nucleic acids. The sample nucleic acid may be prepared in various ways to facilitate detection of the target sequence; e.g., denaturation, restriction digestion, electrophoresis or dot blotting. The region of interest of the target nucleic acid is usually at least partially single-stranded to form hybrids with the probe. If the sequence is double-stranded, the sequence will probably need to be denatured. The target nucleic acid may be also be fragmented to reduce or eliminate the formation of secondary structures. The fragmentation may be performed using a number of methods, including enzymatic, chemical, thermal cleavage or degradation. For example, fragmentation may be accomplished by heat/Mg<sup>2+</sup> treatment, endonuclease (e.g., DNAase 1) treatment, restriction enzyme digestion, shearing (e.g., by ultrasound) or NaOH treatment.

Many genotyping and expression monitoring methods have been described previously. In general, target nucleic acid and probe are incubated under conditions which forms hybridization complex between the probe and the target sequence. The region of the probes which is used to bind to the target sequence can be made completely complementary to the targeted region of the SAC1 locus. Therefore, high stringency conditions may be desirable in order to prevent false positives. However, conditions of high stringency are typically used if the probes are complementary to regions of the chromosome which are unique in the genome. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, base composition, probe length, and concentration of formamide. Under certain circumstances, the formation of higher order hybrids, such as triplexes, quadruplexes, etc. may be desired to provide the means of detecting target sequences.

Detection, if any, of the resulting hybrid is usually accomplished by the use of labeled probes. Alternatively, the probe may be unlabeled, but may be detectable by specific binding with a ligand which is labeled, either directly or

-54-

indirectly. Suitable labels, and methods for labeling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation, random priming or kinase reaction), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies and the like. Variations of this basic scheme are known in the art, and include those variations that facilitate separation of the hybrids to be detected from extraneous materials and/or that amplify the signal from the labeled moiety.

Two-step label amplification methodologies are known in the art. These assays work on the principle that a small ligand (such as digoxigenin, biotin, or the like) is attached to a nucleic acid probe capable of specifically binding SAC1.

In one example, the small ligand attached to the nucleic acid probe is specifically recognized by an antibody-enzyme conjugate. In one embodiment of this example, digoxigenin is attached to the nucleic acid probe. Hybridization is detected by an antibody-alkaline phosphatase conjugate which turns over a chemiluminescent substrate. In a second example, the small ligand is recognized by a second ligand-enzyme conjugate that is capable of specifically complexing to the first ligand. A well-known embodiment of this example is the biotin-avidin type of interactions.

It is also contemplated within the scope of this invention that the nucleic acid probe assays of this invention will employ a cocktail of nucleic acid probes capable of detecting SAC1. Thus, in one example to detect the presence of SAC1 in a biological sample, more than one probe complementary to SAC1 is employed.

Predisposition to diabetes, obesity, or alcoholism can be ascertained by testing any fluid or tissue of a human for sequence variations of the SAC1 gene. For example, a person who has inherited a germline SAC1 mutation would be prone to develop obesity, diabetes, or alcoholism. This can be determined by testing DNA from any tissue of the person's body. Most simply, blood can be drawn and DNA extracted from the cells of the blood. In addition, prenatal diagnosis can be accomplished by testing fetal cells, placental cells or amniotic cells for mutations of the SAC1 gene.

-55-

5        The most definitive test for mutations in a candidate locus is to directly compare genomic SAC1 sequences from obese, diabetic, or alcoholic patients, with those from a control population. Alternatively, one could sequence messenger RNA after amplification, e.g., by PCR, thereby eliminating the necessity of determining the exon structure of the candidate gene.

10      Sequence variations from diabetic, obese, or alcoholic patients falling outside the coding region of SAC1 can be detected by examining the non-coding regions, such as introns and regulatory sequences near or within the SAC1 gene. An early indication that mutations in noncoding regions are important may come from Northern blot experiments that reveal messenger RNA molecules of abnormal size or abundance in obese or diabetic patients as compared to control individuals.

15      Alteration of SAC1 mRNA expression can be detected by any techniques known in the art (see above). These include Northern blot analysis, PCR amplification, RNase protection, and gene chip analysis. Diminished mRNA expression indicates an alteration of the wild-type SAC1 gene.

20      The diabetic, obese, or alcoholic condition can also be detected on the basis of the alteration of wild-type SAC1 polypeptide. For example, the presence of a SAC1 gene variant, which produces a protein having a loss of function, or altered function, may directly correlate to an increased risk of obesity or diabetes. Such variation can be determined by sequence analysis in accordance with conventional techniques. For example, antibodies (polyclonal or monoclonal) may be used to detect differences in, or the absence of, SAC1 polypeptides. Antibodies may immunoprecipitate SAC1 proteins from solution as well as react with SAC1 protein on Western or immunoblots of polyacrylamide gels. Antibodies may also detect SAC1 proteins in paraffin or frozen tissue sections, using immunocytochemical techniques. Immunoassay include, for example, enzyme linked immunosorbent assays (ELISA), radioimmunoassays (RIA), immunoradiometric assays (IRMA) and immunoenzymatic assays (IEMA), sandwich assays, etc.

-56-

Functional assays, such as protein binding determinations, can be used. Finding a mutant SAC1 gene product indicates alteration of a wild-type SAC1 gene.

VII. Drug, Sweetener, and Alcohol Preference Screening

5 This invention is also useful for screening compounds by using the SAC1 polypeptide or binding fragment thereof in any of a variety of drug, sweetener, and alcohol screening techniques.

10 The SAC1 polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, or borne on a cell surface. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant polynucleotides expressing the polypeptide or fragment, preferably in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, for the formation of complexes between a SAC1 polypeptide or fragment and the agent being tested, or examine the degree to which the formation 15 of a complex between a SAC1 polypeptide or fragment and a known ligand is interfered with by the agent being tested.

20 Thus, the present invention provides methods of screening for drugs and sweeteners comprising contacting such an agent with a SAC1 polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the SAC1 polypeptide or fragment, or (ii) for the presence of a complex between the SAC1 polypeptide or fragment and a ligand, by methods well-known in the art. In such competitive binding assays the SAC1 polypeptide or fragment is typically labeled. Free SAC1 polypeptide or fragment is separated from that 25 present in a protein:protein complex, and the amount of free (i.e., uncomplexed) label is a measure of the binding of the agent being tested to SAC1 or its interference with SAC1:ligand binding, respectively.

30 Other suitable techniques for drug, sweetener, and alcohol screening may provide high throughput screening for compounds having suitable binding affinity to the SAC1 polypeptides. For example, large numbers of different small peptide

-57-

test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with SAC1 polypeptide and washed. Bound SAC1 polypeptide is then detected by methods well-known in the art.

5       Purified SAC1 can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to the polypeptide can be used to capture antibodies to immobilize the SAC1 polypeptide on the solid phase.

10      This invention also contemplates the use of competitive drug, sweetener, and alcohol screening assays in which neutralizing antibodies capable of specifically binding the SAC1 polypeptide compete with a test compound for binding to the SAC1 polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants of the SAC1 polypeptide.

15      A further technique for drug, sweetener, and alcohol screening involves the use of host eukaryotic cell lines or cells which have a nonfunctional SAC1 gene. These host cell lines or cells are defective at the SAC1 polypeptide level. The host cell lines or cells are grown in the presence of the drug, sweetener, or alcohol compound. The rate of growth of the host cells is measured to determine if the compound is capable of regulating the growth of SAC1 defective cells.

20      Briefly, a method of screening for a substance which modulates activity of a polypeptide may include contacting one or more test substances with the polypeptide in a suitable reaction medium, testing the activity of the treated polypeptide and comparing that activity with the activity of the polypeptide in comparable reaction medium untreated with the test substance or substances. A difference in activity between the treated and untreated polypeptides is indicative of a modulating effect of the relevant test substance or substances.

25      Prior to or as well as being screened for modulation of activity, test substances may be screened for ability to interact with the polypeptide, e.g., in a yeast two-hybrid system. This system may be used as a coarse screen prior to testing a substance for actual ability to modulate activity of the polypeptide.

-58-

Alternatively, the screen could be used to screen test substances for binding to a SAC1 specific binding partner, or to find mimetics of a SAC1 polypeptide.

### VIII. Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact (e.g., agonists, antagonists, inhibitors) in order to fashion drugs which are, for example, more active or stable forms of the polypeptide, or which, e.g., enhance or interfere with the function of a polypeptide *in vivo*. In one approach, one first determines the three-dimensional structure of a protein of interest (e.g., SAC1 polypeptide) or, for example, of the SAC1-receptor or ligand complex, by x-ray crystallography, by computer modeling or most typically, by a combination of approaches. Less often, useful information regarding the structure of a polypeptide may be gained by modeling based on the structure of homologous proteins. An example of rational drug design is the development of HIV protease inhibitors. In addition, peptides (e.g., SAC1 polypeptide) are analyzed by an alanine scan. In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

It is also possible to isolate a target-specific antibody, selected by a functional assay, and then to solve its crystal structure. In principle, this approach yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

Thus, one may design drugs which have, e.g., improved SAC1 polypeptide activity or stability or which act as inhibitors, agonists, antagonists, etc. of SAC1 polypeptide activity. By virtue of the availability of cloned SAC1 sequences,

sufficient amounts of the SAC1 polypeptide may be made available to perform such analytical studies as x-ray crystallography. In addition, the knowledge of the SAC1 protein sequence provided herein will guide those employing computer modeling techniques in place of, or in addition to x-ray crystallography.

5 Following identification of a substance which modulates or affects polypeptide activity, the substance may be investigated further. Furthermore, it may be manufactured and/or used in preparation, i.e., manufacture or formulation; or a composition such as a medicament, pharmaceutical composition or drug. These may be administered to individuals.

10 Thus, the present invention extends in various aspects not only to a substance identified using a nucleic acid molecule as a modulator of polypeptide activity, in accordance with what is disclosed herein, but also a pharmaceutical composition, medicament, drug or other composition comprising such a substance, a method comprising administration of such a composition comprising such a substance, a method comprising administration of such a composition to a patient, e.g., for treatment of diabetes, obesity or alcohol consumption, use of such a substance in the manufacture of a composition for administration, e.g., for treatment of diabetes or alcohol consumption, and a method of making a pharmaceutical composition comprising admixing such a substance with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

15

20

25

A substance identified as a modulator of polypeptide function may be peptide or non-peptide in nature. Non-peptide "small molecules" are often preferred for many in vivo pharmaceutical uses. Accordingly, a mimetic or mimic of the substance (particularly if a peptide) may be designed for pharmaceutical use.

30

The designing of mimetics to a known pharmaceutically active compound is a known approach to the development of pharmaceuticals based on a "lead" compound. This might be desirable where the active compound is difficult or expensive to synthesize or where it is unsuitable for a particular method of administration, e.g., pure peptides are unsuitable active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary

-60-

canal. Mimetic design, synthesis and testing is generally used to avoid randomly screening large numbers of molecules for a target property.

There are several steps commonly taken in the design of a mimetic from a compound having a given target property. First, the particular parts of the compound that are critical and/or important in determining the target property are determined. In the case of a peptide, this can be done by systematically varying the amino acid residues in the peptide, e.g., by substituting each residue in turn. Alanine scans of peptide are commonly used to refine such peptide motifs. These parts or residues constituting the active region of the compound are known as its pharmacophore.

Once the pharmacophore has been found, its structure is modeled according to its physical properties, e.g., stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g., spectroscopic techniques, x-ray diffraction data and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of a pharmacophore, rather than the bonding between atoms) and other techniques can be used in this modeling process.

In a variant of this approach, the three-dimensional structure of the ligand and its binding partner are modeled. This can be especially used where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the design of the mimetic.

A template molecule is then selected onto which chemical groups which mimic the pharmacophore can be grafted. The template molecule and the chemical groups grafted onto it can conveniently be selected so that the mimetic is easy to synthesize, is likely to be pharmacologically acceptable, and does not degrade in vivo, while retaining the biological activity of the lead compound.

Alternatively, where the mimetic is peptide-based, further stability can be achieved by cyclizing the peptide, increasing its rigidity. The mimetic(s) found by this approach can then be screened to see whether they have the target property, or to what extent they exhibit it. Further optimization or modification can then be carried out to arrive at one or more final mimetics for in vivo or clinical testing.

IX. Gene Therapy

According to the present invention, a method is also provided of supplying wild-type SAC1 function to a cell which carries mutant SAC1 alleles. The wild-type SAC1 gene or a part of the gene may be introduced into the cell in a vector such that the gene remains extrachromosomal. In such a situation, the gene will be expressed by the cell from the extra chromosomal location. More preferred is the situation where the wild-type SAC1 gene or a part thereof is introduced into the mutant cell in such a way that it recombines with the endogenous mutant SAC1 gene present in the cell. Such recombination requires a double recombination event which results in the correction of the SAC1 gene mutation. Vectors for introduction of genes both for recombination and for extrachromosomal maintenance are known in the art, and any suitable vector may be used. Methods for introducing DNA into cells such as electroporation, calcium phosphate coprecipitation and viral transduction are known in the art, and the choice of method is within the competence of skilled practitioners.

As generally discussed above, the SAC1 gene or fragment, where applicable, may be employed in gene therapy methods in order to increase the amount of the expression products of such genes in diabetic or obese cells. Such gene therapy is particularly appropriate, in which the level of SAC1 polypeptide is absent or compared to normal cells. It may also be useful to increase the level of expression of a given SAC1 gene even in those situations in which the mutant gene is expressed at a "normal" level, but the gene product is not fully functional.

Gene therapy would be carried out according to generally accepted methods, for example, as described by *Therapy for Genetic Diseases*, T. Friedman, ed. Oxford University Press, 1991. Cells from a patient would be first analyzed by the diagnostic methods described above, to ascertain the production of SAC1 polypeptide in these cells. A virus or plasmid vector, containing a copy of the SAC1 gene linked to expression control elements and capable of replicating inside the sample cells, is prepared. Suitable vectors are known, such as disclosed in PCT publications WO 93/07282 and United States

-62-

Patent Nos. 5,252,479, 5,691,198, 5,747,469, 5,436,146 and 5,753,500. The vector is then injected into the patient.

Gene transfer systems known in the art may be useful in the practice of the gene therapy methods of the present invention. These include viral and nonviral transfer methods. A number of viruses have been used as gene transfer vectors, including papovaviruses, e.g., SV40, adenovirus, vaccinia virus, adeno-associated virus, herpes viruses including HSV and EBV; lentiviruses, Sindbis and Semliki Forest virus, and retroviruses of avian, murine, and human origin. Most human gene therapy protocols have been based on disabled murine retroviruses.

Nonviral gene transfer methods known in the art include chemical techniques such as calcium phosphate coprecipitation; mechanical techniques, for example microinjection; membrane fusion-mediated transfer via liposomes; and direct DNA uptake and receptor-mediated DNA transfer. Viral-mediated gene transfer can be combined with direct *in vivo* gene transfer using liposome delivery, allowing one to direct the viral vectors to the affected cells and not into the surrounding nondividing cells. Alternatively, the retroviral vector producer cell line can be injected into affected cells. Injection of producer cells would then provide a continuous source of vector particles.

In an approach which combines biological and physical gene transfer methods, plasmid DNA of any size is combined with a polylysine-conjugated antibody specific to the adenovirus hexon protein, and the resulting complex is bound to an adenovirus vector. The trimolecular complex is then used to infect cells. The adenovirus vector permits efficient binding, internalization, and degradation of the endosome before the coupled DNA is damaged. For other techniques for the delivery of adenovirus based vectors see United States Patent Nos. 5,691,198; 5,747,469; 5,436,146 and 5,753,500.

Liposome/DNA complexes have been shown to be capable of mediating direct *in vivo* gene transfer. While in standard liposome preparations the gene transfer process is nonspecific, localized *in vivo* uptake and expression may be accomplished following direct *in situ* administration.

Expression vectors in the context of gene therapy are meant to include those constructs containing sequences sufficient to express a polynucleotide that

-63-

has been cloned therein. In viral expression vectors, the construct contains viral sequences sufficient to support packaging of the construct. If the polynucleotide encodes SAC1, expression will produce SAC1. If the polynucleotide encodes an antisense polynucleotide or a ribozyme, expression will produce the antisense 5 polynucleotide or ribozyme. Thus in this context, expression does not require that a protein product be synthesized. In addition to the polynucleotide cloned into the expression vector, the vector also contains a promoter functional in eukaryotic cells. The cloned polynucleotide sequence is under control of this promoter. Suitable eukaryotic promoters include those described above. The expression 10 vector may also include sequences, such as selectable markers and other sequences described herein.

Receptor-mediated gene transfer, for example, may be accomplished by the conjugation of DNA (usually in the form of covalently closed supercoiled plasmid) to a protein ligand via polylysine. Ligands are chosen on the basis of the 15 presence of the corresponding ligand receptors on the cell surface of the target cell/tissue type. One appropriate receptor/ligand pair may include the estrogen receptor and its ligand, estrogen (and estrogen analogues). These ligand-DNA conjugates can be injected directly into the blood if desired and are directed to the target tissue where receptor binding and internalization of the DNA-protein 20 complex occurs. To overcome the problem of intracellular destruction of DNA, coinfection with adenovirus can be included to disrupt endosome function.

#### X. Peptide Therapy

Peptides which have SAC1 activity can be supplied to cells which carry mutant or missing SAC1 alleles. Protein can be produced by expression of the 25 cDNA sequence in bacteria, for example, using known expression vectors. Alternatively, SAC1 polypeptide can be extracted from SAC1-producing mammalian cells. In addition, the techniques of synthetic chemistry can be employed to synthesize SAC1 protein. Any of such techniques can provide the preparation of the present invention which comprises the SAC1 protein.

-64-

Preparation is substantially free of other human proteins. This is most readily accomplished by synthesis in a microorganism or in vitro.

5 Active SAC1 molecules can be introduced into cells by microinjection or by use of liposomes, for example. Alternatively, some active molecules may be taken up by cells, actively or by diffusion. Extra-cellular application of the SAC1 gene product may be sufficient. Molecules with SAC1 activity (for example, peptides, drugs or organic compounds) may also be used to effect such a reversal. Modified polypeptides having substantially similar function are also used for peptide therapy.

10 XI. Transformed Hosts

Similarly, cells and animals which carry a mutant SAC1 allele can be used as model systems to study and test for substances which have potential as therapeutic agents. These may be isolated from individuals with SAC1 mutations, either somatic or germline. Alternatively, the cell line can be engineered to carry 15 the mutation in the SAC1 allele.

Animals for testing therapeutic agents can be selected after mutagenesis of whole animals or after treatment of germline cells or zygotes. Such treatments include insertion of mutant SAC1 alleles, usually from a second animal species, as well as insertion of disrupted homologous genes. Alternatively, the endogenous 20 SAC1 gene of the animals may be disrupted by insertion or deletion mutation or other genetic alterations using conventional techniques to produce knockout or transplacement animals. A transplacement is similar to a knockout because the endogenous gene is replaced, but in the case of a transplacement the replacement is by another version of the same gene. After test substances have been 25 administered to the animals, the phenotype must be assessed. If the test substance prevents or suppresses the disease, then the test substance is a candidate therapeutic agent for the treatment of disease. These animal models provide an extremely important testing vehicle for potential therapeutic products.

In one embodiment of the invention, transgenic animals are produced 30 which contain a functional transgene encoding a functional SAC1 polypeptide or

-65-

variants thereof. Transgenic animals expressing SAC1 transgenes, recombinant cell lines derived from such animals and transgenic embryos may be useful in methods for screening for and identifying agents that induce or repress function of SAC1. Transgenic animals of the present invention also can be used as models for studying indications such as diabetes.

5 In one embodiment of the invention, a SAC1 transgene is introduced into a non-human host to produce a transgenic animal expressing a human or murine SAC1 gene. The transgenic animal is produced by the integration of the transgene into the genome in a manner that permits the expression of the transgene. Methods 10 for producing transgenic animals are generally described in US Patent No. 4,873,191 and in *Manipulating the Mouse Embryo; A Laboratory Manual*, 2nd edition (eds., Hogan, Beddington, Costantini and Long, New York: Cold Spring Harbor Laboratory Press, 1994).

15 It may be desirable to replace the endogenous SAC1 by homologous recombination between the transgene and the endogenous gene; or the endogenous gene may be eliminated by deletion as in the preparation of "knock-out" animals. Typically, a SAC1 gene flanked by genomic sequences is transferred by microinjection into a fertilized egg. The microinjected eggs are implanted into a host female, and the progeny are screened for the expression of the transgene. 20 Transgenic animals may be produced from the fertilized eggs from a number of animals including, but not limited to reptiles, amphibians, birds, mammals, and fish. Within a particularly preferred embodiment, transgenic mice are generated which express a mutant form of the polypeptide.

25 As noted above, transgenic animals and cell lines derived from such animals may find use in certain testing experiments. In this regard, transgenic animals and cell lines capable of expressing wild-type or mutant SAC1 may be exposed to test substances. These test substances can be screened for the ability to reduce overexpression of wild-type SAC1 or impair the expression or function of mutant SAC1.

-66-

**XII. Pharmaceutical Compositions and Routes of Administration**

The SAC1 polypeptides, antibodies, peptides and nucleic acids of the present invention can be formulated in pharmaceutical compositions, which are prepared according to conventional pharmaceutical compounding techniques. See, 5 for example, *Remington's Pharmaceutic. Sciences*, 18th Ed. (Easton, PA: Mack Publishing Co., 1990). The composition may contain the active agent or pharmaceutically acceptable salts of the active agent. These compositions may comprise, in addition to one of the active substances, a pharmaceutically acceptable excipient, carrier, buffer, stabilizer or other materials well-known in 10 the art. Such materials should be nontoxic and should not interfere with the efficacy of the active ingredient. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., intravenous, oral, intrathecal, epineural or parenteral.

For oral administration, the compounds can be formulated into solid or 15 liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, suspensions or emulsions. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, suspending agents, and the like in the case of oral liquid preparations (such as, for example, 20 suspensions, elixirs and solutions); or carriers such as starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations (such as, for example, powders, capsules and tablets). Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical 25 carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by standard techniques. The active agent can be encapsulated to make it stable to passage through the gastrointestinal tract while at the same time allowing for passage across the blood brain barrier. See for example, WO 96/11698.

30 For parenteral administration, the compound may be dissolved in a pharmaceutical carrier and administered as either a solution or a suspension.

-67-

Illustrative of suitable carriers are water, saline, dextrose solutions, fructose solutions, ethanol, or oils of animal, vegetative or synthetic origin. The carrier may also contain other ingredients, for example, preservatives, suspending agents, solubilizing agents, buffers and the like. When the compounds are being 5 administered intrathecally, they may also be dissolved in cerebrospinal fluid.

The active agent is preferably administered in a therapeutically effective amount. The actual amount administered, and the rate and time-course of administration, will depend on the nature and severity of the condition being treated. Prescription of treatment, e.g., decisions on dosage, timing, etc., is within 10 the responsibility of general practitioners or specialists, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of techniques and protocols can be found in *Remington's Pharmaceutical Sciences*.

15 Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cell, by the use of targeting systems such as antibodies or cell specific ligands. Targeting may be desirable for a variety of reasons, e.g., if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

20 Instead of administering these agents directly, they could be produced in the target cell, e.g., in a viral vector such as described above or in a cell based delivery system such as described in United States Patent No. 5,550,050 and PCT publications WO 92/19195, WO 94/25503, WO 95/01203, WO 95/05452, WO 96/02286, WO 96/02646, WO 96/40871, WO 96/40959 and WO 97/12635, 25 designed for implantation in a patient. The vector could be targeted to the specific cells to be treated, or it could contain regulatory elements which are more tissue specific to the target cells. The cell based delivery system is designed to be implanted in a patient's body at the desired target site and contains a coding sequence for the active agent. Alternatively, the agent could be administered in a precursor form for conversion to the active form by an activating agent produced 30 in, or targeted to, the cells to be treated. See for example, EP 425,731A and WO 90/07936.

-68-

## EXAMPLES

The following examples further illustrate the present invention. These examples are intended merely to be illustrative of the present invention and are not to be construed as being limiting.

5

### EXAMPLE 1

**Animal care and maintenance.** All animal protocols used in these studies were approved by the Monell Institutional Animal Care and Use Committee. Mice were housed in individual cages in a temperature- controlled room at 23°C on a 12-hour light:12-hour dark cycle. The animals had free access to deionized water 10 and Teklad Rodent Diet 8604 (Harlan Teklad, Madison, WI).

### EXAMPLE 2

**Breeding of F2 and partially congenic mice.** C57BL6/ByJ (B6) and 129P3/J (formerly named 129/J; abbreviated here as 129) mice were purchased from The Jackson Laboratory. The B6 and 129 mice were outcrossed to produce 15 the first filial generation of hybrids (F<sub>1</sub>), and these were intercrossed to produce the second hybrid generation (F<sub>2</sub>, n = 629).

To create the partially congenic lines, the F<sub>2</sub> mice were genotyped with several markers on the distal part of chromosome 4, and a few F<sub>2</sub> mice with recombinations in this region were used as founders of strains partially congenic 20 with the 129 strain. These F<sub>2</sub> founders were backcrossed to the 129 strain to produce the N<sub>2</sub> generation. Mice from this and subsequent backcross generations were phenotyped using 96-hour two-bottle tests with saccharin solutions, and 25 genotyped using markers on distal chromosome 4 and on other autosomes. Mice with high saccharin intake (with a fragment of distal chromosome 4 from the B6 strain and homozygous for 129 alleles of markers on other chromosomes) were selected for subsequent backcrossing. This marker-assisted selection resulted in a segregating 129.B6-Sac partially congenic strain. Three strains were created, with different overlapping fragments containing the SAC1 gene.

-69-

### EXAMPLE 3

#### Testing of sweet preference in the F2 and partially congenic mice.

Consumption of 120 mM sucrose and 17 mM saccharin (Sigma Chemical Company, St. Louis, MO) was measured in individually caged mice using 96-hour two-bottle tests, with water as the second choice. The positions of the tubes were switched every 24 hours. Fluid intakes were expressed per 30 g of body weight (the approximate weight of an adult mouse) per day, or as a preference score (ratio of average daily solution intake to total fluid intake, in percent).

### EXAMPLE 4

Genotyping of F2 mice and linkage analysis. Genomic DNA was purified from mouse tails by NaOH/Tris (Beier, personal communication; Truett G.E. et al., Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT) [In Process Citation]. *Biotechniques*, 2000;29:52, 54), or the phenol/chloroform method. All F2 mice were genotyped with all available polymorphic microsatellite markers (Research Genetics, Huntsville, AL) known to map near the SAC1 region with a protocol modified slightly from that described by Dietrich W. et al., A genetic map of the mouse suitable for typing intraspecific crosses. *Genetics*, 1991;131:423-447. The markers tested are as follows: D4Mit190, D4Mit42, D4Mit254, and D4Mit256. Analysis of this framework map using MAPMAKER/QTL 1.1 (Lander E. et al., MAPMAKER: An interactive complex package for constructing primary linkage maps of experimental and natural populations. *Genomics*, 1987;1:174-181), indicated that *Sac* mapped distal to D4Mit256, and therefore all available STS and EST were tested by SSCP (Orita M., Iwahana H., Kanazawa H., Hayashi K., and Sekiya T. Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proceedings of the National Academy of Sciences of the USA*, 1989:86) or direct sequencing, for polymorphisms between the B6 and 129 strains. Polymorphisms between strains were found for the following markers: D18346, AA410003 (K00231), V2r2, and D4Erdt296E, and the linkage analysis conducted again using these polymorphic makers.

-70-

#### EXAMPLE 5

**Genotyping of partially congenic mice.** Three partially congenic strains of mice were genotyped with all available markers, and those markers with two 129 alleles were excluded from the SAC1 nonrecombinant interval.

5

#### EXAMPLE 6

**Radiation hybrid mapping.** To generate additional markers to narrow the Sac nonrecombinant interval, several markers were tested using the T31 RH genome map. Primers from several sequences suggested through survey of the public databases were constructed and DNA from the T31 panel. Results were 10 scored using software at the Jackson Laboratory.

10

#### EXAMPLE 7

**Construction of BAC contig and marker development.** To construct a physical map of the SAC1 region, the RPCI-23 BAC library was screened with markers within and near the SAC1 nonrecombinant interval: each marker was 15 tested by whole cell PCR to confirm its presence. Only those markers positive by both hybridization and PCR are shown. Primers for the BAC ends were constructed from sequence obtained through TIGR ([www.tigr.org](http://www.tigr.org)) or by direct sequencing, when necessary. Each positive clone was tested for the presence of each BAC end (if the BAC end contained unique sequence), and the contig 20 oriented using SEGMAP, Version 3.48 (Green E.D. and Green P. Sequence-tagged site (STS) content mapping of human chromosomes: theoretical considerations and early experiences. *PCR Methods Appl.*, 1991;1:77-90). BAC end sequences was amplified in B6 and 129 strains, and analyzed by SSCP or direct sequencing. Those BAC ends polymorphic between 129 and B6 were tested 25 in the recombinant F2 and partially congenic mice, to further narrow the SAC1 nonrecombinant interval.

25

-71-

#### EXAMPLE 8

**Amplification of SAC1 and polymorphism detection.** After the SAC1 nonrecombinant interval was narrowed to less than 350 kb, a 246 kb BAC was chosen for sequencing which spanned most of the region (RPCI-23-118E21).

5 Within this BAC, there was a gene with homology to other taste receptors. Using 11.8 kb of sequence and the program GENSCAN (Burge C.B. and Karlin S. Finding the genes in genomic DNA. *Current Opinion Structural Biology*, 1998;8:346-354), a 858 amino acid protein, with 6 exons, was identified. Primers were constructed that amplified this gene, and an additional 2600 nt upstream and

10 5200 nt downstream were also amplified (primer sequence available upon request). These PCR products were sequenced using genomic DNA from B6 and 129 mouse strains, as well as other strains with either higher (SWR/J, C57L/J, IS, ST/bJ, SEA/GnJ) or lower (DBA/2J, AKR/J, BALB/cByJ) saccharin preference (Lush I.E., The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and

15 sucrose. *Genet Res.*, 1989;53:95-99; Lush I. The genetics of bitterness, sweetness, and saltiness in strains of mice. in *Genetics of perception and communication*, Vol. 3, eds. Wysocki C. and Kare M., New York: Marcel Dekker, 1991:227-235; Lush I.E. and Holland G. The genetics of tasting in mice. V. Glycine and cycloheximide. *Genet Res.*, 1988;52:207-12). Sequences were aligned with

20 Sequencer (Gene Codes, Ann Arbor, MI) and the single nucleotide polymorphisms, insertions and deletions identified.

#### EXAMPLE 9

**Preparation of tongue cDNA and expression studies.** Total RNA was extracted from anterior mouse tongue from the 129 and B6 strains (TRIZOL Reagent; GIBCOBRL). Total RNA (200 ng) was reverse transcribed using the Life Technologies SuperScript Kit. Following the reverse transcription, the samples were amplified using Advantage cDNA PCR Kit (Clontech, Palo Alto, CA). Primers were constructed to span exon 2 and 3, so that the genomic and cDNA product size would differ (Primer set 3A; 30 Left-5'TGCATTGGCCAGACTAGAAA3';

-72-

Right-5CGGCTGGCTATGACCTAT'). The expected product size for primer 3A is 418 bp for cDNA and 497 bp for genomic DNA. Single bands of these sizes were excised from the gel, purified and sequenced, confirming the intron-exon boundary and expression of mRNA of this gene in mouse tongue. Primers were 5 then designed to cover the whole cDNA, and, the sequences obtained and aligned, to confirm intron/exon boundaries.

#### EXAMPLE 10

Human gene expression. The human ortholog of the SAC1 gene was examined for mRNA expression in human tongue. Total RNA from human taste 10 papillae was obtained through biopsy, a procedure approved by the Committee on Studies Involving Human Beings at the University of Pennsylvania. The RNA was extracted as described above, reverse transcribed, and amplified, with human specific primers. Two bands were obtained of the expected size for genomic and cDNA. Sequencing of these bands confirmed the SAC1 gene is expressed in 15 human taste papillae.

#### EXAMPLE 11

Tissue Expression of SAC1. Oligonucleotide primers specific for different parts of the SAC1 gene were used to assay different tissues for SAC1 expression as shown in Table 2. Tissue specific cDNA pools were purchased from 20 OriGene Technologies Ltd. Primer pair 3A, amplifies parts of exons 2 and 3, with a small intron to differentiate between PCR product representing genomic DNA versus cDNA. Primer pair 6A amplifies parts of exons 4 and 5. This part of the protein encodes the 7TM domain, and may cross react with other GPCRs expressed in different tissues.

-73-

Table 2. Expression pattern of SAC1

Tissue	3A	6A
Brain	-	-
Heart	-	-
Kidney	+	+
Spleen	+	+
Thymus	+	+
Liver	-	+
Stomach	-	+
Sm Intestine	-	+
Muscle	-	+
Lung	-	+
Testis	+	+
Skin	-	-
Adrenal	+	-
Pancreas	+	+
Uterus	-	-
Prostrate	+	+
Embryo-8.5	-	-
9.5	-	-
12.5	-	-
19	+	-/+
Breast-virgin	-	+
Pregnant	-	+
Lactating	+	+
Involuting	-	-

-74-

## EXAMPLE 12

Primers for the SAC1 Locus (Seq. ID Nos.: 6-651) are:

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
28.MMHAP7B4. seq	CACTAGAGCTGCC ACCTTCC	CCCTCAGCACCA CTTTTGT	162	6-7
28.MMHAP7B4. seq	ACAAAAAGTGGTG CTGAGGG	CAGGAGACCCA AAGGATCAA	163	8-9
AA408705	GCTTCAGAAAATC GAGGCAC	GCATGGGCTATG ATAGGTGG	232	10-11
AA408705	TGTTGATCCCACA GCG	CAGGAAATGTCC ACTTCTGC		12-13
AA409223	TCTATCTTGCATC CAGCC	GTGCTGTGACTG TGCG		14-15
AA589460	CGCAGCATTATT TGGAG	CCGACCCTTAG GAGACAC		16-17
Agrin4	TGTGACTCCTCTT CCCCAC	TGAGCCACTCCA GATGTCAG	156	18-19
Agrin4	GTGTGTCAGCATC ACTGCCT	CCAACGTGCAGT CAAGAAAA	290	20-21
Agrin4	CGAGAGACAAAG TGGTGCTG	TTATGAAGGCC TCACCAAC	263	22-23
Agrin4	CCAGCTCCTAGAA TTGCCTG	GCAGTCTCCGA AACAAAGTC	298	24-25
Agrin4	ATAGAGGAATGG GTGCGATG	TACCAAGGAGGG GTCAGTCAG	299	26-27
Agrin4	TACAAGCGAGCTG ACCAATG	CCAATCAGCTCG AGTTAGCC	271	28-29
AgrinA	TGCCATTGTGGAT GTTCACT	GAGTCCGAGGTC GGTCAATA	575	30-31

-75-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
AgrinB	GCTGGCTTCTGTA GGTCAGG	TATGAGGGTCAA GGGTCAAGG	577	32-33
AgrinC	CGCTTGGTGAGA ACTAGCC	CATGTGGAGTTG TGGGAGTG	573	34-35
AgrinD	AATGGGCAGAAG ACAGATGG	TATCAGGGTCTG TGAAGCCC	507	36-37
AgrinE	ATACAGGACCCTT TACCCCG	CAGTGTCTTAG GTCCCCCA	587	38-39
Agrin	GCCTCTGTCTGCC ATCTCTC	ATAATGTTACCT GCAGGCGG	594	40-41
AI115523	CTGGAAACACCCA TGTCCCTC	CGGGCACATGG ACACTTTA	200	42-43
AI225779	GAGCATGAAGTGC AAGGTGA	CGTAGGTGGCAC AGTTGAGA	266	44-45
AI225779	GCTGTTAGTGAGG TCAGGGC	CGTAGGTGGCAC AGTTGAGA	104	46-47
AI225779	GAGCATGAAGTGC AAGGTGA	TCATTTCCCTAG CCTCGGTG	126	48-49
A022703	TCTAAGAAGATGA TGCAGACCC	TGTCCTTCAGGG ATAGTGCC		50-51
Cdc2l2	GGCTTCAGCCTCA AGTTCTG	AAAACAACCAA GTTGCCCTG	101	52-53
Cdc2l2	GGCACTGAAATGA CCTGGAT	AACAATTCAAGC AACCTCGG	265	54-55
Cdc2l2	CTGTTCCCTCCCA GACTCCA	TTCAGTCACGCA AACCTGAG	225	56-57
Cot	GCCCAGGACTTG TCACTGT	GGTAACCTGCAG CTCCACTC	284	58-59

-76-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
Cot	GGGACATGCTCTT GGTTCAT	GAACAAAGCCG GGTGATTAA	277	60-61
Cot	GCCCTCAGTTCTC CTAGCCT	GGCAGAGAAGA CTGGTGGAG	110	62-63
Cot	CCCAGACTTAGCG TCTCAGG	AGCAGAGACCTT TGGACTCG	277	64-65
Cot	GAAGGCTGAGTGA GTCCCAG	TTGCACGAGGAG AAGGTTTT	276	66-67
Cot	GATGCCAACGAGA CCTGAAT	AGAAGCCAAAAA CCCTCACCT	247	68-69
Cot	AAAAAGCCCTGCA AGAACTT	ATTCAAGGTCTCG TTGGCATC	107	70-71
D18346	TGTCCGCAGTGTG GAAACTA	ATGTCCAGGGTA GAGAGCCC	165	72-73
D18402	GGAGTTCTCCTAC CCTGGCT	GAGGCTCTGAGC AGTGTCAA	167	74-75
D4Bir1	GCGATGTTGTTG CG	CAGTGTCTTCC ACATT		76-77
D4Ertd296e	AGGCATATTGTAT AATAAATTGTAT GT	CCGGATGACTCT ACTTGAC	201	78-79
D4Hrb1	GCTGTTATGGGG TCGAGAA	AATTTCTGAAGC AGGGGGAT	194	80-81
D4Hrb1	TCCCCCTGCTTCA GAAATTAA	AGGGGGATGATT GTGAGTGA	192	82-83
D4mit313	CTTCTTTAATCAA TCTCTGTCTCTGTG	GGGCACATATGA ACCTCCTG	196	84-85
D4mit344	CCAAACTCTTAGC TTCTTCA	ACACAGAAGAC ACTGAAGAAC	187	86-87

-77-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
D4Mit51	CAGTTGTTAGAAG CAGGATCCC	AGGTGCATATAC CTGGGATACTC	123	88-89
D4Mit59	AGAGTTGGTCTC TTCCCCTG	TATCCAACACAT TTATGTCTGCG	108	90-91
D4Mit59	GCCAGTGTGCTGA AAGACTG	AGGGACCTGGA GACATCCTT	119	92-93
D4Nds16	CTGTAGGCTGCTT TTATCTTTG	TGCCCTTCAGC ACATGCCA		94-95
D4smh6b	TGCAGTGTGACAT GTGCATAGAT	GGAAAGCCAGG CTACGCAGAA	118	96-97
D4smh6b	CTGTAGGCTGCTT TTATCTTTG	TGCCCTTCAGC ACATGCCA	102	98-99
D4smh6b	TAGTGTGGTTCCT GACTAACCT	CGGTCTACATAG TGAGTGATTTC	181	100-101
D4smh6b	AAAAGCATCCTGC ATCCTTCTG	GGGTTATACAGA GAAACCCTGT	83	102-103
D4Xrf215	TTCCAAGCTCACA CATCAGC	GTGCTGCTCTGC ATTGAGTG	124	104-105
D4Xrf243	GACAGTGTGGGAG AATCCGT	CCCAAGGCATAG GTCACAAT	203	106-107
D4Xrf243	ATTGTGACCTATG CCTTGGG	CGAAGGACCGTC ATCTGAGT	105	108-109
D4Xrf472	GGCTTGATGTGA AAAAGGC	AGCTCCTCATCG CTCATGTT	245	110-111
D4Xrf472	TGGAACATCTCTG TCGGAAG	GGCTCTCATTGC CACCTTTA	193	112-113
D4X497	CCAGAGAACAGG AGACCTGC	GTGCTGGATACA CTGGCAGA	119	114-115

-78-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
D4Xrf497	GCGAGACGAGTG GGTAGTTC	ACACTGAAACCT CGCTTGCT	129	116-117
D4Xrf497	AGCAAGCGAGGTT TCAGTGT	ACGGGGCTTGAT CCTTTAT	204	118-119
Dshv4	AAGTCATGGGCC TCACCACCTGTC	TACTAGCTACCC TTCACATACC	100-300	120-121
Dshv5	ACCTAGCCACTGT CTCAGTCT	ACAGAAGCAGC ATTACACAG	100-300	122-123
Gnb1	TGGGACAGCTTCC TCAAGAT	AATGGGAATTGT GCTCTTGG	213	124-125
Gnb1	GGGCATCTGGCAA AGATTAA	AGATAACCTGTG TGTCCCGC	281	126-127
Gnb1	GATGTCCGAGAAG GGATGTG	TGTCA GCTTGA GTGCATCC	277	128-129
Gnb1	ACATGCAGGCTGT TTGACCT	TGTCA GCTTGA GTGCATCC	166	130-131
K00231	GTGCTCTGCAGAC AAACCAA	GAGCCATTGAA CCCTTAAA	154	132-133
K00231	TTTCAGGGTCAAA ATGGCTC	TCGACAGCAACT GTGCG		134-135
K00954	GGTGAGAGTGGG GAGATGAA	CCCGGGTGAGTT TAAGAACCC	237	136-137
k00954	GGTGAGAGTGGG GAGATGAA	AGGTAGGCCA ATTCCCTG	296	138-139
k00954	CCAGGGTTGCTGT ACTGAGA	CAGGTTAGGCC AATTCCCT	237	140-141
K01153	GGTCAGAGTCCTT CCTTCCC	TCCAACITCACA GGAAACCC	124	142-143

-79-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
K01153	TTTCCTGTGAAGT TGGAGGG	CACCCATATGGC AAACATCA	213	144-145
K01153	GGTCAGAGTCCTT CCTTCCC	TCCAACTTCACA GGAAACCC	125	146-147
K01153	TGATGTTGCCAT ATGGGTG	GCTTGCTGCTTC CGATATGT	181	148-149
K01599	GGAAAAGGGAGT CGCCATA	GAGCCGCCAAC TCTCACAC	166	150-151
K01599	AGGGGATAACCTG CATAGG	ACAAAATTGCTC ATTTGCCC	110	152-153
<u>M-05262</u>	CCATCCCCACTAG CCAGATA	GTCCCCCTTGTC ACAGCAAG	169	154-155
M107-H01	TGAGCACAGGATA GCTCCAC	AAAAGAACACC TGTTGGGG	217	156-157
M111-B04	TAAACCTCGGCTG TGTGAG	CCCTCAGTGACT TCCTGTGA	267	158-159
M134-C06	CAAAACCACATGG TTACCGA	GCCCTATTGCCA AATGACTT	264	160-161
M134-G01	GGCAGAAAGGAA TCAGAACG	CACATTAGCCAT TGTCTGG	161	162-163
M136-B01	TCCTTATGTCCA ACAGCCA	CATGGTCTGTGA TGTGACCA	164	164-165
M156-H05	ATACCCTGGTGA GAGCAGG	GCTGTCAAATGA GAAAGGCA	139	166-167
M184-B03	TATTTCATGCTGG GACCAAA	AGAGAAAAACA GTGGGGGTG	89	168-169
Mmp23	CGGGTCCTCTCTT CACCATA	CTACATTCCCT GAGCTGCC	297	170-171

-80-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
Mmp23	GTTGACCATGTCG GTAACCC	CCACCTCACCGA AACTGAAT	111	172-173
Mmp23	GGTGTGCGCTCA CAAACCT	GATGCACACACA AAAATCCG	197	174-175
Mmp23	ATCACCCACCAGA ACGAAAA	ACCCTCCAGGAG TAGGTGCT	255	176-177
PCEE	GATGAGACAGTGG GCAAGGT	TTGTCAATAGCA CCAAGCCA	154	178-179
PCEE	GCCTTAATAGCCC CCTTGTT	GCACTCAGCATT GCACAGAT	194	180-181
PCEE	GGACGGACAATTG TGGAAAA	CTATCACACCTC CGATGCCT	142	182-183
PCEE	CAAGCTGGTAGAA TCCCCAA	TCTTGGAGAAG CAGACCGT	209	184-185
Pkcz	TACAGCATATGCA TGCCAGG	ATTCCTCAGGGC ATTACACG	294	186-187
Pkcz	GCAATCTCTGTG TCCAGGC	ATTCCTCAGGGC ATTACACG	188	188-189
Pkcz	TACAGCATATGCA TGCCAGG	GGCCTGGACACA AGAGATTG	127	190-191
Pkcz	AAGTGGGTGGACA GTGAAGG	CAGCTTCCTCCA TCTTCTGG	201	192-193
Pkcz	AGAGCCTCCAGTA GATGGCA	TCGTGGACAAGC TCCTTCTT	297	194-195
Pkcz	CATCGAGTATGTC AATGGCG	TTGTCCAGTTT AGGTCCCG	156	196-197
Pkcz	CAGACTGGTTT CCGACAT	GTCAAAGTTGTC CAGGCCAT	132	198-199

-81-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
Pkcz	AGGACGGACCCCA AGATG	TGTCTCGCACTT CCTCACAG	130	200-201
Pkcz	CCAGAAAGATGGA GGAAGCTG	TCTACTGGAGGC TCTTGGGA	151	202-203
Pkcz	GAAAAACGACCA GATTACG	GATCTCAGCAGC ATAGAACCC	265	204-205
Pkcz	ACACATTAAGCTG ACGGACT	CAAACATAAGG ACACCCAGT	164	206-207
Pkcz	ACTGGGTGTCCTT ATGTTTG	CCTCTCTTGGG ATCCTTAT	193	208-209
Pkcz	GTCATAAAAGAGGA TCGACCA	GCTCTGTCTAGA AGTGCCTG	252	210-211
Pkcz	ACCAAGACCGAA GAGGGG	GGCATTACACGC TAACTTTCC	223	212-213
R74924	AGTGCCACCAACC TGGTAAG	AAGTGCCTGCAG GGATGC	165	214-215
R74924	TGCTTGGTGAGC AATGTTT	AGGGACACCCCTT ACCAGGTT	103	216-217
R74924	CTGATGCTTGGT GAGCAAT	GGGACACCCCTTA CCAGGTT		218-219
R75150	ACAGGACAAATGC TGGGTG	GTGGTAAAGAA CGCTTGGCT	217	220-221
R75150	GGTATCTCACTTG GTAGGAACCTC	AAGAACGCTTGG CTGGC		222-223
RER1 (1)	GCCGATCCTGGTG ATGTACT	ACAATGGCTCAA AACCGTTC		224-225
RER1 (2)	GCCTTGGGAATT ACCACCT	AGTACATCACCA GGATCGGC		226-227

-82-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
RER1	TAAAAGGCCATGC GATAAGC	AGAGCTCTGTGG GGTTCTCA		228-229
RER1	GAAGGGGACAGT GTTGGAGA	TCCATCAAGGAA GGATCCAC		230-231
Tp73	GGTGGGTAATGAT TGGACT	TGACGTGGAGG GAACTGCC	296-301	232-233
Tp73	TGAGATCTGGTGC CCTCTCT	GCCTGATCTAGG CTGGAAAA	222-229	234-235
Txgp1	AGGCAGAAAGCA GACAAGGA	CGACAGCACTTG TGACCACT	138	236-237
Txgp1	CTGCAGATGTAGA CCAGGCA	CTGTGGTGGATT GGACAGTG	269	238-239
Txgp1	TTGCCTAACACTC CCAAACC	TATTAGGAGCAC CACCAAGGC	244	240-241
Txgp1	ACCTGTCTTGTGG GTGGAAG	CTGTGGTGGATT GGACAGTG		242-243
U37351	GTGGCTTGGTGCT ATTGACA	GGGGCTATTAAG GCCATTT	160	244-245
V2R2	CAATTGAGGAATG GCTACCAA	TGGCTTCATGTC CATTGTGT	170	246-247
V2R2	CAGAACACACAA GGTAAATTGC	TCATGTTGCTG TCCAGTTG	163	248-249
TR1-like1(human)	GCCACCATGCTGG GCCCTGCTGCCT GGG	TCACTCATGTTT CCCCTGATTCC	2520	250-251
T1-ike2(human)	CTGATTTCCCTGTG TTCCCGT	CATGCTGGCCTA CTTCATCA	244	252-253

-83-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
T1-like3(human)	GCCTTGCAGGTCA GCTACGGTGCTAG CAT	TCACTCATGTTC CCCTGTATTCC	2441	254-255
T1-like4(human)	AGGAAGCAGAGA AAGGCCAG	TCAGAACTGCCT CTGAGCTG	274	256-257
T1-like5(human)	TCTTCACGTACTG GGGGAAC	ACTACAGCATCA GCAGCAGG	175	258-259
T1-like6(human)	AAGCTGAAGAACT TCCCCGT	TGGGCTACGACC TCTTGAT	211	260-261
h-Tr1like a	ATCTTCAGGCGCT CTGTCCT	GTACGACCTGAA GCTGTGGG		262-263
h-Tr1like b	ATCTTCAGGCGCT CTGTCC	GTACGACCTGAA GCTGTGGG		264-265
h-Tr1like c	ATCTTCAGGCGCT CTGTCC	GAGTACGACCTG AAGCTGTGG		266-267
h-Tr1like d	ATCTTCAGGCGCT CTGTCCT	TACGACCTGAAG CTGTGGG		268-269
h-Tr1like e	ATCTTCAGGCGCT CTGTCC	TACGACCTGAAG CTGTGGG		270-271
h-Tr1like f	GCTGTCCCGATGG TGAAC	ACCTTTGTGGC CAGGATG		272-273
h-Tr1like g	GCTGTCCCGATGG TGAAC	CACCTTTGTGG CCAGGAT		274-275
h-Tr1like h	GCTGTCCCGATGG TGAAC	CCTTTGTGGCC AGGATG		276-277
h-Tr1like I	CCTGAACCAGTGG GCTGT	ACCTTTGTGGC CAGGATG		278-279
h-Tr1like j	CCTGAACCAGTGG GCTGT	CACCTTTGTGG CCAGGAT		280-281

-84-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
h-Tr1like k	TCATGTTCCCT GATTCC	CATGCTGGCCTA CTTCATCA		282-283
h-Tr1like	ATGAGCAGGTAAC ACCTGGG	TCATCACCTGGG TCTCCTT		284-285
h-Tr1like m	ATGAGCAGGTAAC ACCTGGG	TTCATCACCTGG GTCTCCTT		286-287
mTr1like-1A	TGGGTTGTGTTCT CTGGTTG	CCTTTTACAGT CTGCCAGGT		288-289
mTr1like-1B	TGGGTTGTGTTCT CTGGTTG	GATCCCCTTTT ACAGTCTGC		290-291
mTr1like-2A	ACGGGGTTGGTAC TGTGTGT	CACCCATTGTTA GTGCTGGA		292-293
mTr1like-2B	ACGGGGTTGGTAC TGTGTGT	CACACACCCACC CATTGTTA		294-295
mTr1like-3A	TGCATTGCCAGA CTAGAAA	CGGCTGGCTAT GACCTAT		296-297
mTr1like-3B	TGCATTGCCAGA CTAGAAA	CGGCTGGCTAT GACCTATT		298-299
mTr1like-4A	GTTCTGCAGCATG ATGTCGT	GGCAGTTGTGAC TCTGTTGC		300-301
mTr1like-4B	GTTCTGCAGCATG ATGTCGT	CTGCAGGCAGTT GTGACTCT		302-303
mTr1like-5A	CCATCCTTTGCC TGTCTT	TCTGGAGGAACA TGTGATGG		304-305
mTr1like-5B	CACCATCCTTT GCCTGTC	GAACATGTGATG GGGCAAC		306-307
mTr1like-6A	CAAAGCAGCAGG AGGAGTG	AAATGTACTGGC CAGGCAAC		308-309

-85-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
mTr1like-6B	AGTGCTAGACCCA GCACCAAG	AAATGTACTGGC CAGGCAAC		310-311
mTr1like-7A	GCACTGACCAGTC TGTCAAC	GTCCCCAGAGAA AAGCACAG		312-313
mTr1like-7B	CAGTCTGTCACCA CCTCTGG	CAGTGGTCCCCA GAGAAAAG		314-315
mTr1like-8A	TACTATTGGGGC TTGTTGG	GCAGCACTATGT GCCTGGTA		316-317
mTr1like-8B	TACTATTGGGGC TTGTTGG	GCCTGGTATTG ATCGCTTT		318-319
mTr1like-9A	GCTCAGCTAGGGGA TGGAGAA	CAGCTCAGGGAC ACAATGAA		320-321
mTr1like-9B	TCCTACAGGCTAG GGCTCAG	CAGCTCAGGGAC ACAATGAA		322-323
mTr1like-10A	GGGACTGATGTGT GGCTTGT	AGGCGTCCCAGG AATAGAAG		324-325
mTr1like-10B	GGACTGATGTGT GCTTGT	AGGCGTCCCAGG AATAGAAG		326-327
mTr1like-11A	TGTTTCTGTTCTGG TGGCTG	ATCTGCAGGCAG GATCAGAC		328-329
mTr1like-11B	CTCAGTGGTGGGT GACAGTG	ATCTGCAGGCAG GATCAGAC		330-331
Mutation1 (mouse)	ACACACAGTACCA ACCCCGT	CCTGTGGTGATC AAGAAAGCA	182	332-333
Mutation2 (mouse)	TGCTTCTTGATCA CCACAGG	GCAACAGAGTC ACAACTGCC	131	334-335
Mutations1+2 (mouse)	ACACACAGTACCA ACCCCGT	GCAACAGAGTC ACAACTGCC	293	336-337

-86-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
34m15-T7	GGGTTTATGTGGC AAGCACT	ACTCCATTGCC TTTGTGG	118	338-339
34m15-SP6	CGCTACTTCGCTT TTATCCG	ATGATGACGTAC GACGACGA	150	340-341
37D20-T7	GAAAACAATCGG GGAGAAAGTC	TGAAATTATCAC ACGCCAGG	109	342-343
37D20-T7(3)*	AGTGAGAGGCCA GTCTCAA	GATCTGATGCC TCTTCTGC	247	344-345
37D20-SP6	GCTAGCCTGAAG CCAACAC	TGAACAGCATGC TTACCCAG	122	346-347
49O2-T7	TCCCTAGAGGCCT GTCTGTC	TCGTCTGGAGC CTCTTCTA	169	348-349
49O2-SP6	GATAGTCCCTAG CCAGCCC	GCCATAGCTCCT CACTGCTC	218	350-351
73B10-T7	CAGAGTGGCTCT GGTCTTC	TTGTGTTAGAT GCTCCTGC	237	352-353
73B10-SP6	TTATTCTGTGCTA GCCGCC	ATCAAGTCAACG TCCCCAAG	267	354-355
75M14	ACCTGGCCTGTGC TAATCTC	GCACCAACCTA AGAAAGCA	233	356-357
85G18	TCAGGCTAACCTC AAACTCACA	AAAGAAAAGAA AAGAAAAAGTC AGACA	113	358-359
118E21-T7	CCCAGAACTCCAT CCTCAAA	CCCAACCTGTGG TCAGCTAT	185	360-361
118E21-SP6	GGGGCAGGTGGGT AATAAGT	CAAAAGCCAA CTCCTTGAG	271	362-363
130A12-T7*	GCTCAGTGGGTAA GAGCACC	CTACCCCTGCCGC TAATCTCA	242	364-365

-87-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
130A12-SP6	CAGTTAGCACCCC ACCCCTAA	TCTGCACCTCTG TTCACCTG	114	366-367
138D7-T7	ACCTCTAGGGTTT ACGGGGA	CCTCAGGTAGTG CAAGCTCC	199	368-369
139J18-T7	TCAGTTACCAAGG GTTTCGG	ATAGGTTGTCAC AGGCCAGG	122	370-371
139J18-SP6	TCAGTTACCAAGG GTTTCGG	ATAGGTTGTCAC AGGCCAGG	122	372-373
147a15-T7*	GTGGTTGCTGGGA TTTGAAC	CAAGCAACCAA ACAACCAAA	101	374-375
147A15-SP6	TCCGGAGGACCAT AAATCTG	CACAGTCCCAGT CATTCCCT	249	376-377
151E4-T7	GTCCCCAAAGCTA GCACAGG	TCATGAGGCCACC ATGTGATT	240	378-379
151E4-SP6	GACCTTCGGAAGA GCAGTTG	AGTGTGTGTCGC CATATCCA	223	380-381
152O3-T7	CCTACTCTCTCTCC CCGCTT	GGAAAATGTTG GCCTTGAA	142	382-383
152O3-SP6	CTGGAGTGAAAGG CAGGAAG	AGGCAGGACCAT ATGAATAA	537	384-385
153B21SP6	TGAGAGTGGGAAT TCTGTTCA	GGATGTAATTGG TGGCAAGG	202	386-387
153B21T7	CTGTTGGAGGAGG TGGCCTA	TGCTTGTATGTT TTTCCTCGT	113	388-389
159J19SP6	TGAGAGTGCCCTC CTCTTTG	GAACCCCTGACC CCAGAC	200	390-391
159J19T7	TGAAGTGCAGATT TTTACATGG	GTTTTGGGGTGG AAAAGGAT	213	392-393

-88-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
189M12SP6*	CCGTCGACATTAA GGTGACA	GATACTGGGGTG GTGGGTAA	189	394-395
227G4-SP6	CCGTCGACATTAA GGTGACA	CGTCCCAGCTGT GTAACTGA	219	396-397
227G4-T7*	GGAAGCAAATGCT CCACTAAA	TATCCCTAGCCC CTTGTGTG	243	398-399
236C12-SP6	CCGTCGACATTAA GGTGACA	GGGTCCCTGTTGG TAGTGACC	209	400-401
238O5T7	TATAAGCAGCCCC TCATTGG	CAGGCCAGACA CTGCTTACA	244	402-403
238O5SP6	CCTTGGGATCTGG TGTGACT	TGGGTTTAGAGT ACGGCTGG	251	404-405
24718-T7	ACCCATTTCCCTAA TCCCCCTG	ATCTCTCCAGCC CCTCTCAG	177	406-407
280G12-T7*	GGGCTGGGAATTG AACCTAT	TGAATCCCTTAC AGCCTTGC	420	408-409
280G12-SP6	GCCCCATAAAATC CACTCCT	GCTCCGGAAGGC TAGAAGAT	233	410-411
284D21-T7	GGTTTGGGAGTGT TAGGCAA	ACTCAGTTGGCC TCTCCTCA	138	412-413
284D21-SP6	ACAGAAATCCCTC ATGCGA	TCAGTGTGGACC AGAAAGTCC	105	414-415
298E4	TCTGCAAGTCAGC TCTTGATAA	ACTCATAAGGGT CAAGCTGTCTG	100	416-417
298e4-T7(3)*	TCTCCCTTTTACCA ACTCCC	GCAAGGAGTCA AAAACAGCA	180	418-419
307ES	GCTAGTTGGGAA CAAACCA	ACTGCAAATGTC CAACTCCA	149	420-421

-89-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
338N4-T7	CAGTTACACAGCT GGGACGA	GCAAGAGCCTA GCAATCCAC	245	422-423
338N4-SP6	CAGTTAGCACCC CACCCCTA	TCTGCACCTCTG TTCACCTG	115	424-425
348P19-SP6	GGGTTCCACTTGA TGCTGAT	TGGTCTGTTCC TGGAGCTT	227	426-427
350D2-T7*	TGTAGGAAATGTT TCTGCACC	ACATGGAACAG GATTCTGGC	295	428-429
350D2-SP6	GCAGGGCAAACAG ACAGACAA	ATGGGGGATCCC TTACTGAC	217	430-431
360M12-T7	CGGTCAAGGAGTAG TGTGGGT	CAGCAGCTGATA TTGAGGCA	123	432-433
360M12-SP6	AATGATGAAGTGT CAGCCTCAG	CAACAGAACTCA AAGCCTGG	100	434-435
382A8-SP6	AGCAGGCACAGGT CTCTTGT	AAGAACAGGAC AGTGGTGGG	202	436-437
382A8-SP6(2)	CAGCGATTGGCTC TTCTCTT	GGGGCTTCCTTT CTGAGGTA	531	438-439
386N4-T7	AGCTCAGGTCCAG CTTGGTA	ATTTCCCTCC TGCTTCTC	107	440-441
386N4-SP6	CCAAGCCTCTGCT GGTTATC	TGAGGGTGGAG AATGGAAAG	109	442-443
387-T7	GCCCCATAAAATC CACTCCT	TTGCCTAACACT CCCAAACC	214	444-445
387-SP6	CAGTTACACAGCT GGGACGA	GCAAGAGCCTA GCAATCCAC	245	446-447
388I1	CAGCACCTTCCTC TGGTCTC	TGTCTCCAGAGG TTCTGCCT	137	448-449

-90-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
399I12-T7	TGGTGGTGTAAATA CTATTCCCTTG	TCTTTAATTTTT GGCTTTTGATA CA	102	450-451
399I12-SP6	CAGCTGTGTGCAT GTTGACC	CATCATGAAGAC TCAGGGCA	106	452-453
415A22SP6	GTCCACACCTGGC TTTTGTT	CAGCACTCAGTG AGGTTCCA	199	454-455
415G24SP6	ATGTAATGGAAGG GCTGCTG	CAGCACTCAGTG AGGTTCCA	113	456-457
417B22-SP6	AAACAGGCATGA AACTCAGGA	GGGTATCATTGT CACCTCCA	116	458-459
436P10-T7	CACAGGCCAAGTT GTTGTTG	CAGGGGACCTTC TGAATGAT	115	460-461
438C18-T7	AGCTCAGGTCCAG CTTGGTA	ACCACAAAATT TCCCCCTCC	115	462-463
438C18-SP6	CGGGACCTAAAAC TGGACAA	TGGGGACAGTTA CCAGGAAG	254	464-465
457N22-T7	CCGGAGGACCATA AATCTGA	CCTAAAAACAA GCCTGAGC	129	466-467
457N22-SP6	CCTTCAGAAATGT GTTTGGACA	TCCTGAGTTCAA ATCCCAGC	252	468-469
472018	CTTTCATTCTCCA CCCTCA	AGGTCTAGGGA GAGGTCCA	260	470-471
D4Mon1	AGGCCTACCCAAG GACATCT	GCAGTGAGCTGC AGAGTTG	201	472-473
D4Mon2	AGACACCCTAGGT CCTGCTG	TGATCTTCCAA ACGCATAAGA	151	474-475
D4Mon3	GCAAGCAACCTGA ACATGAA	GCTTACGATGGT CGTGAGGT	188	476-477

-91-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
D4Mon4	ACATGCCTGCCTA TCTTGCG	GGAACCTGTTT CCATGGTG	197	478-479
D4Mon5	ACCTTGTCCTGG TGTGAGC	TAGCTGGGACGT GGTATGGT	200	480-481
D4Mon6	CCATGGGAGACCA GAAGGTA	TGAGTGTCCCTCT GCCTGATG	206	482-483
D4Mon7	GCGCTGACATCCT CCTATGT	CCCACTATGGTC CCAGAGAA	187	484-485
D4Mon8	TTGCACGTCTTG TTTCGAG	AAAGGGGAATA GACCTGAGTAG AA	219	486-487
D4Mon9	CCAAGAGTCAGCC TTGGAGT	GGACAGGTAGCT CACCCAAC	200	488-489
Tr1like1cDNA mouse	TGCCAGCTTGGC TATCAT	TTCATTGTGTCC CTGAGCTG		490-491
Tr1like2cDNA mouse	AGCTTGGCTATC ATGGGTCTCAG	ACCACCGCCACT GTTCTCATCT		492-493
Tr1like_A1 (mouse)	TGTGGGGAAGA ACATAGAA	TGATGTGTGGCT TGTTCCTCTT	5935	494-495
Tr1like_A2 (mouse)	ATAGGTGGGAG GGAGCTAA	TGATGTGTGGCT TGTTCCTCTT	5903	496-497
TR1 like-2 (human)	TGTGCCTGTCACA GCAACTT	CATGCTAGCACC GTAGCTGA		498-499
TR1 like-3 (human)	GGAGACCTTCCCC TCCTTCT	GCTGTAGTTGAA GAGGGCGT		500-501
TR1 like-4 (human)	GTGCTTGGCTTCC TCCAG	CAGGTCGTACTC CATGTCCA		502-503
TR1 like-5 (human)	TGGAGTACGACCT GAAGCTG	ACTCATCCTGGC CACAAAAG		504-505

-92-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
TR1 like-6 (human)	GAACAGGAGGAC GCTGAGG	CTTTGTGGCCA GGATGAGT		506-507
TR1 like-7 (human)	TCACCTCACCTGG TTGTCAG	GTACGACCTGAA GCTGTGGG		508-509
TR1 like-8 (human)	GGCTGAGATCACA GGGTTGGGTCACT C	CCGTGCCTGTTG GAAGTTGCCTCT GCC		510-511
118e21-0	AATTCCCAGCAAC CACTCAC	CAGACACTCCAG AAGAGGGC	585	512-513
118e21-1	TGACTGCTCTTCC GAAGGTT	TTTGTGGAATAG CCAAAGCC	588	514-515
118-21-2	TCTCTCCTCTCTTC TCCCCC	AGCAGGGTGCAT CACCTTAT	551	516-517
118e21-3	TAGGAGTGCCCCA TAGGTTG	TCATTGTACCCA GCCAGTCA	518	518-519
118e21-4	AGGACTGAGCCTG GATGAGA	CTGGGCGTTTG TTTTGTTT	552	520-521
118e21-5	CTTCCTCCTGCAG CTACCAC	ACCCCTGCTACAA CGCAGACT	546	522-523
118e21-6	TCCAACCTTGACA CCCATT	AGCCAGGGCTAC ACAGAGAA	584	524-525
139J18T7(1)	CTGCTTTCTCTCA GCAACTG	ATTGCCGTTAG AAGCTAGG		526-527
139J18T7(2)	AACTGTACGTGGC TGCTGGT	ATTGCCGTTAG AAGCTAGG		528-529
Agrin(CA)n	GCCAGGTGACCC TATGAAA	GAGAGATGGCA GACAGAGGC	271	530-531
Agrin(TG)n	AGCTCTCTGTCCC TGGTGAA	TGCCAACCACCA GCCTCTCT	157	532-533

-93-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
repeat1	CTGAACCCTCCAC TCTCCTG	AGCCAGGGCTAC ACAGAGAA	205	534-535
repeat2	AGCCAGGGCTACA CAGAGAA	ACCCTGCTACAA CGCAGACT	153	536-537
repeat3	GCAAGTTTCAGGA GCTAGGG	CCCCAGAACCCAG AGACCATA	166	538-539
repeat4	CTAGGGGACTCTG CCAAGTG	CAAGACACCCA GTCCCAACT	195	540-541
repeat5	TACTTCCCCTTCC CGAACT	TCCTTGGTGCTT ACCCTCAC	232	542-543
repeat6	TGTTCCTGAGTTC ACAACGC	ATTCCCAGCAAC TACATGGC	269	544-545
repeat7	ACATGTCCACTGT GGCAAAA	TGTGATGAGTTT GAGGCCAG	246	546-547
repeat8	ATCAGACAGCCCA CAACCTC	TATGTGCCACCA CACCTGTC	206	548-549
repeat9	GCTCAAGGAAGG ACACACCT	TGCTCTAACAT TTTGAGCCAT	201	550-551
repeat10	GCTCAGCCCCTGA ATCAATA	GGGATCTGCCTG TCTTACCA	111	552-553
repeat11	GGAAGGTAGGGC CTGGTAAT	GCTCCAAGATCT GTGCGATT	277	554-555
repeat12	TTAGCGTTAGGGT GAGGGTG	GGAGACTACGG ACTTGTGGC	150	556-557
repeat13	CAGTTCTTCCCGA AAACCAC	TTTCTGGGAACT GAGATGGC	174	558-559
repeat14	GTTGGGGCTGCTC ATAGAAA	GCTGTGGCTCTC TTGGAGTT	422	560-561

-94-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
repeat15	CTCTGATTCCCA CATGCCT	AAGAGGGAGCA CTGAGGACA	152	562-563
repeat16	CAGCAGCAAATGA CCTTTCA	GAGGCAGGCAG ATTCTGAG	147	564-565
repeat17	GTTTCACATGTTG TGGTGGC	GGGACCTTGGG ATAGCATT	131	566-567
repeat18	TCAGACATCTCTG GCCTCCT	TTCACTAAGTTG CCCAGGCT	160	568-569
repeat19	TGCCTTTCTCAC ATTGTCTC	TTAGAAGCAGA GGCAGAGGC	250	570-571
repeat20	GACCTTGGAAGA GCAGTCG	TGGCAGCTCACA ATGTCTT	296	572-573
SHANRU1	GGTGTGGTGTAGG GGAAGAA	TTTCAACTGCAA ACACAAACAG	301	574-575
SHANRU2	AGGGCCAAGGAA GGAGAAT	GCAAATATATAG GGTACCGAGCTG	203	576-577
SHANRU3	CAGATTCTCCAGC TGTCAAGG	CTGTGTTCCGC ACCAAGT	229	578-579
SHANRU4	CTGCCCGTCCTTA TCTTCTG	ACGCACGCTCAC TCATACAC	289	580-581
SHANRU5	CAGCAGAGGTGAT GGGTTCT	TTGTCACACAGT GGTTAAATGC	203	582-583
SHANRU6	TAGAACCGTGGCT GAGGACT	CCGTAAGATAT GAAAGAACTTG GA	201	584-585
SHANRU7	TAATCCTGGCTTA GCGCTTG	TAGAAAGCACA GGGGACAGG	240	586-587
SHANRU8	CCTTCCTCGTCTG AGCTGTT	TTGGGACGTGAC CTGAGAAT	232	588-589

-95-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
SHANRU9	TATGTGTCTGGCC GTTGTTTC	GATGTGGGTGCA GGTGAAG	206	590-591
SHANRU10	CCCCTTCTGGAGT GTCTGAA	TCTAGGCAGGGC TACCTTTT	263	592-593
SHANRU11	GCTGAGCAGCCTC TAGCAA	ACCATGGCTTT CCCAGTAA	241	594-595
SHANRU12	CTGTGCCTTGTT GATCAGA	TGTGGCACTCTA CGGCATAA	261	596-597
SHANRU13	TGCATCACTATT AGCCTCAACC	AAGAATTGCAA AGACTGTGAGA	260	598-599
SHANRU14	AGCCAGCGCTACA CAGAGA	CTGGACCTTG AAGAGCAG	199	600-601
SHANRU15	GGTGGCTAAACC ATCCATA	GAGGGCAATGA GCAAAATGT	203	602-603
SHANRU16	GGTCCTGTCTTG GTTCAAGG	TAACACCCACAT CAGGCAAC	201	604-605
SHANRU17	TTTCATTTCTGGT GTTCTTT	AAACACAGGCG GAACGATAG	198	606-607
SHANRU18	CTATCGTCCGCC TGTGTTT	AAGGAAGAGGA TGGAGAAAGA	397	608-609
SHANRU19	CGGGTCTTAATGG AGCAGAG	TCCTCCCCAGTT ACCTAGCA	222	610-611
SHANRU20	CAGCAGGCAAGAT GACCTC	GTCCCTCACCAAG CCATGTTA	205	612-613
SHANRU21	AGCCTGGGCTAAG TTGTGTG	TATGGGCCAATG TTGTTCCCT	204	614-615
SHANRU22	ATGGTGGCTCAC ACCATCT	TTGTCCTCTGAT TGCAGCAT	193	616-617

-96-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
SHANRU23	CTTGGGTACATCAG GCTTTGT	AAGCTGCCCTGC TCTCTCTA	301	618-619
SHANRU24	ATGCTCAGCCTGC TTTGTTC	GCTGATAGCCCT GGGTTCTA	198	620-621
SHANRU25	TGTACGCACAAAT TGACTTGC	GAATCCACATTG CAAAGCCTA	222	622-623
SHANRU26	CACAGGCAAATGAA AGGGAAG	CCAGACTTCTCC AGCTCTCC	187	624-625
SHANRU27	TCCTCGAGAGGCT CTAGGTTT	TGCCTAGTCAAC CACAGGAG	237	626-627
SHANRU28	CCTGTGGTTGACT AGGCAGAA	GCCTGATAGCCT GGAATACA	406	628-629
SHANRU29	AAAGGGATGTGTG GCGTAAG	CAAAACCCAACC TTCTCAGC	195	630-631
SHANRU30	TGCACTGACCGTG ATAGAGG	CGGTGTAGCTCT GGCTGTCT	200	632-633
SHANRU31	CATCTCACCAACT CGCACTT	TTTCTGGGAACA AAGAGGCTA	418	634-635
SHANRU32	GAACCCAAGTGT GGGGTAA	TGGAAGCCCATC TGTCTCTT	222	636-637
SHANRU33	AAATGCAAGTGGG TGCTTCT	CCAGAAGAGGG CGTCAGAT	187	638-639
SHANRU34	GGTGTGCACCACC ATATTCA	GGGAATTATCAG CCAAAAAGC	201	640-641
SHANRU35	GCCCAACTGAAAG CTCAACT	GGAAGGGGGAT AACAAATTGAA	263	642-643
SHANRU36	TGCTAATTCAAG CACAGTGAGA	AGCTTGACACCT TGACAGCA	369	644-645

-97-

Marker	Forward	Reverse	Size, bp	SEQ. ID NO.
SHANRU37	AACCTGCAGAGAG GAGACCA	CTCCAAGGGGA GGACTCATT	201	646-647
SHANRU38	TTCAATTGAGTTT CTCTCCTCTGA	TGCAGGACCAA GAAGTAGGC	200	648-649
SHANRU39	CGAGATCTGATGC CCTCTTC	TGCTGAGAGCAG AAAAGGAA	200	650-651

Although the foregoing invention has been described in some detail by way of illustrating and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

5 All publications, patents, and web sites are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent, or web site was specifically and individually indicated to be incorporated by reference in its entirety.

-98-

CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a sequence variation of SEQ ID. NO 1, wherein said variation is associated with sensing carbohydrates, other sweeteners, or ethanol.  
5
2. An isolated polynucleotide comprising a sequence variation of SEQ ID. NO 2, wherein said variation is associated with sensing carbohydrates, other sweeteners, or ethanol.
3. An isolated polynucleotide comprising a sequence variation of SEQ ID. NO 4, wherein said variation is associated with altered sensation of carbohydrates, other sweeteners, or ethanol.  
10
4. The polynucleotide of Claim 1 wherein said variation is a missense mutation.
5. The polynucleotide of Claim 4 wherein said variation is a nonsense mutation.  
15
6. An isolated polypeptide comprising a variant form of SEQ ID. NO: 3, wherein said variant form is associated with altered preference for carbohydrates, other sweeteners, or ethanol.
7. An isolated polypeptide comprising a variant form of SEQ ID. NO 5, wherein said variant form is associated with altered preference for carbohydrates, other sweeteners, or ethanol.  
20
8. An isolated polynucleotide having at least 8 contiguous nucleotides of the polynucleotides of any one of the Claims 1-3 wherein said 8 contiguous nucleotides span said variation position.

-99-

9. An isolated polypeptide having at least four contiguous amino acids of the polypeptides of Claims 6 or 7 wherein said four contiguous amino acids span said variation position.
10. An isolated polynucleotide wherein said polynucleotide is substantially identical to the polynucleotide of Claim 8.
- 5 11. An isolated polypeptide wherein said polypeptide is substantially identical to the polypeptide of Claim 9.
12. An isolated polynucleotide having a sequence which is complementary to the polynucleotide of Claim 8 or 10.
- 10 13. A polynucleotide specific for the SAC1 locus wherein said polynucleotide hybridizes, under stringent conditions, to at least 8 contiguous nucleotides of the polynucleotide of Claim 1, 2, 3, or 4.
14. The polynucleotide according to Claim 13 wherein said polynucleotide is selected from the group consisting of SEQ ID. NOS 6-651 and homologous equivalents thereof.
- 15 15. A polynucleotide specific for the SAC1 locus wherein said polynucleotide that hybridizes, under stringent conditions, to at least 8 contiguous nucleotides of the polynucleotide of Claim 3.
16. The polynucleotide of Claim 15 wherein said polynucleotide is selected from the group consisting of SEQ ID. NOS 6-651 and homologous equivalents thereof.
- 20 17. A kit for the detection of the polynucleotide of any one of Claims 1-5, 8, and 10 comprising a polynucleotide that hybridizes, under stringent conditions, to at least 12 contiguous nucleotides of the polynucleotide of any one of the Claims 1-5, 8, and 10, and instructions relating to detection.
- 25

-100-

18. An isolated antibody which is immunoreactive to the polypeptide of  
Claim 9 or 11.
19. A method for analyzing a biomolecule in a biological sample, wherein said  
method comprising:
  - 5 a) altering SAC1 activity in a biological sample; and
  - b) measuring the activity.
20. A method for analyzing a polynucleotide in a biological sample  
comprising the steps of:
  - 10 a) contacting a polynucleotide in a biological sample with a probe  
wherein said probe hybridizes to the polynucleotides of Claim 8  
or 10 to form a hybridization complex; and
  - b) detecting the hybridization complex.
21. A method for analyzing the expression of SAC1 comprising the steps of  
a) contacting a biological sample with a probe wherein said probe  
15 comprises the polynucleotide of Claim 8 or 10; and
- b) detecting the expression of SAC1 mRNA transcript in said sample.
22. The method of Claim 19 wherein said step of measuring is an enzymatic  
assay.
23. The method of Claim 20 or 21 wherein said probe is immobilized on a  
20 solid support.
24. The method according to any one of the Claims 19-23 wherein said sample  
is derived from blood.
25. The method according to any one of the Claims 19-23 wherein said sample  
is derived from tongue.

-101-

26. The method according to any one of the Claims 19-23 wherein said sample is derived from pancreas.
27. The method according to any one of the Claims 19-23 wherein said sample is derived from a human.
- 5 28. A method for identifying susceptibility to obesity or diabetes which comprises comparing the nucleotide sequence of the suspected SAC1 allele with a wild type nucleotide sequence, wherein said difference between the suspected allele and the wild-type sequence identifies a sequence variation of the SAC1 nucleotide sequence.
- 10 29. An expression vector comprising the polynucleotide of Claim 3, 8, or 10.
30. A host cell comprising the expression vector of Claim 29.
31. A method of producing a polypeptide comprising culturing the cells of Claim 30 and recovering the polypeptide from the host cell.
32. An isolated polypeptide produced according to Claim 31.
- 15 33. A method for conducting a screening assay to identify a molecule which enhances or decreases the SAC1 activity comprising the steps of
  - a) contacting a biological sample with a molecule wherein said biological sample contains SAC1 activity; and
  - b) analyzing the SAC1 activity in said sample.
- 20 34. A pharmaceutical composition comprising
  - a) the polynucleotide of Claim 8 or 10, the polypeptide of Claim 9 or 11, the antibody of Claim 18 or the molecule of Claim 18; and
  - b) a suitable pharmaceutical carrier.

-102-

35. A method for treating or preventing obesity, diabetes, or alcoholism associated with expression of SAC1, wherein said method comprises administering to a subject an effective amount of the pharmaceutical composition of Claim 34.
- 5 36. A transgenic animal that carries an altered SAC1 allele.
37. The transgenic animal of Claim 36 is a knock out mouse.
38. The polypeptide of Claim 6 or 7, wherein said polypeptide is 7-transmembrane G protein coupled receptor (7TM GPCR).

## SEQUENCE LISTING

<110> Bachmanov, Alexander A  
Beauchamp, Gary K.  
Chatterjee, Aurobindo  
De Jong, Pieter J.  
Li, Shanru  
Li, Xia  
Ohmen, Jeffrey D  
Reed, Danielle R.  
Ross, David  
Tordoff, Michael G.

<120> GENE AND SEQUENCE VARIATION ASSOCIATED WITH SENSING  
CARBOHYDRATE COMPOUNDS AND OTHER SWEETNERS

<130> Gene & Sequence Variation.....

<140>

<141>

<150> 60/200,794

<151> 2000-04-28

<160> 652

<170> PatentIn Ver. 2.1

<210> 1

<211> 2577

<212> DNA

<213> Mouse

<400> 1  
atgccagctt tggctatcat gggcttcagc ctggctgctt tcctggagct tggatgggg 60  
gcctctttgt gtctgtcaca gcaattcaag gcacaagggg actacatact gggcgggcta 120  
tttccccctgg gctcaaccga ggaggccact ctcaaccaga gaacacaacc caacagcata 180  
ccgtgcaaca ggttctcacc ccttggtttg ttcctggcca tggctatgaa gatggctgtg 240  
gaggagatca acaatggatc tgccttgc tc cctggctgc ggctgggcta tgacctattt 300  
gacacatgct ccgagccagt ggtcaccatg aaatccagtc tcatgttctt ggccaagggtg 360  
ggcagtcaaaa gcattgtgc ctactgcaac tacacacagt accaaccctcg tgtgctggct 420  
gtcattggcc cccactcatac agagttgcc ctcattacag gcaagttctt cagcttcttc 480  
ctcatgccac aggtcagcta tagtgcacatg atggatggc taagtgaccg ggaaacgttt 540  
ccatccttct tccgcacatg gcccagtgcac cgggtgcac tgcaaggcagt tgtgactctg 600  
ttgcagaact tcagctggaa ctgggtggcc gccttaaggaa gtgatgatga ctatggccgg 660  
gaaggtctga gcatctttc tagtctggcc aatgcacgag gatctgcat cgcacatgag 720  
ggcctggcacaacatga cactagtggc caacagtgg gcaagggtgct ggatgtacta 780  
cgccaaagtga accaaagtta agtacaagtg gtggctgtt ttgcctctgc ccgtgctgctc 840

tactcccttt ttagttacag catccatcat ggcctctcac ccaaggatg ggtggccagt 900  
gagtcttggc tgacatctga cctggtcatg acacttccc atattgcccc tgtgggact 960  
gtgcttgggt ttttgcagcg gggtgcctca ctgcctgaat tttcccatta tgtggagact 1020  
caccttgcac tggccgctga cccagcattc tgcctcac tgaatgcggg gttggatctg 1080  
gaggaacatg tgatggggca acgctgtcca cggtgtgacg acatcatgtc gcagaaccta 1140  
tcatctggc tgttcagaa cctatcagct gggcaattgc accaccaaatttgc 1200  
tatgcagctg tgtacagtgt ggctcaagcc ctgcacaaca ccctacagtg caatgtctca 1260  
cattgccacg tattcagaaca tggtctaccc tggcagctcc tggagaacat gtacaatatg 1320  
agtttccatg ctcgagactt gacactacag ttgtatgtc aagggaaatgt agacatggaa 1380  
tatgacctga agatgtgggt gtggcagagc cctacacctg tattacatac tgtggcacc 1440  
ttcaacggca ccctcagct gcagcagct aaaatgtact ggccaggca ccaggtgcca 1500  
gtctcccagt gttcccgcca gtgc当地 ggc当地 ggc当地 ggc当地 1560  
tcctgtctgt atgactgcgt ggactgc当地 gc当地 gagct accggaagca tccagatgac 1620  
ttcacctgta ctccatgtaa ccaggaccag tggccccc当地 agaaaagcac agcctgtta 1680  
cctcgcaggc ccaagttct ggctggggg gagccagttg tgctgtcact cctcctgtg 1740  
ctttgcotgg tgctgggtct agcactggct gctctgggc tctctgtcca ccactggac 1800  
agccctcttg tccaggcctc aggtggctca cagttctgtt ttggccctgat ctgc当地 taggc 1860  
ctcttctgct tcagtgctt tctgttccca gggcggccaa gctctgccc ctgc当地 tgc当地 1920  
caacaaccaa tggctcacct ccctctcaca ggctgc当地 gc当地 acactctt cctgcaagca 1980  
gctgagacact ttgtggagtc tgagctgc当地 ctgagctggg ccaaactggct atgcagctac 2040  
cttcggggac tctggccctg gctagtggtt ctgtggccca ctttgggatg ggc当地 gacta 2100  
tgtgccttgt atttgatgc当地 ttcccacca gagtggtga cagactggct agtgc当地 gccc 2160  
acagagggtac tggagactg ccacgtgc当地 tcctgggtca gc当地 gggctt ggtgc当地 acatc 2220  
accaatgcaa tggtagctt ctctgtctt ctggccactt tcctggtaca gagccagctt 2280  
ggccgctaca accgtgc当地 tggctcacc ttgc当地 catcaccctgg 2340  
gtcttttg tgccctctctt ggccaaatgtg cagggtggctt accagccagc tggc当地 agatg 2400  
ggtgctatcc tagtctgtgc cttggccatc ctggtcaccc tccacctgccc caagtgc当地 tttat 2460  
tgtgc当地 tttt ggctgc当地 aaacc gctcaacacc caggagttct tcctgggaaag gaatgc当地 aag 2520  
aaagcagcag atgagaacag tggc当地 ggtt gaggcagctc agggacacaa tgaatga 2577

<210> 2  
<211> 11809  
<212> DNA  
<213> Mouse

<400> 2  
atctgagccct .tagacacagc actgggtgcca ggcaaacact cctgggccta catgcttggg 60  
gcctcttcat attccaaaag ctgtcttgg gtaagatgaa gttcctctgg cagtggcatg 120  
agtgtctgaag gctctttccc tgcccttac ctgcttctt gatagtctt ctgcatacca 180  
aacaggccct tgctcctgg gaaatggaaa ctatgaaatc aatagcttag gcttctctag 240  
gaaaggcttc cctggtcagt acaacctgtt tcacagcttc tatagaatag ttacatcagc 300  
cttctgaaga tggccttta gagcacatgc acccccaaga ttctaaatgt tcaataactaa 360  
ctgacccaaac cataaccttc tagccagccc tgctgtctt gttgtctggt acccaggtga 420  
ctgaggacat gactggtgga agggaaactag .gcccccttgt ctgtcagatg gccataccac 480  
gcatggctga tgcccagtgt ataagacccct acgttttcc actggtctta atgttaaacc 540  
ctaggacagt gtccctcagca tagctgggtgt gtgtgaatgc aaactttggg gcataatctt 600  
tccattaagc actgtgatataat atgttagtatt tccaaacaaat aaattataacc tacatgattg 660

ggtagatcat tctggatgg gtcacaggtg tgcaggatgc ctaattatgt gggggaaagaa 720  
catagaaata tatacgatgg gaggagacta accctagaa taaggctaaa gcatgtgtct 780  
ccagtcctga agactcaaag ggcaacgtga atcatgagac atgttcagga ctgaaggagt 840  
tgccatgtat ctgtccttga tgcataatcactatgagat ctgtgttacc 900  
tccatggc aggtgagaaa agaaacacct gaatggccta ccttaaaggaa ctaagtggaa 960  
aaataggatctt gaagataacc caggcactgt gtgacaaggc gggaaagaaaa ctagagatgc 1020  
tttcttcatg gcaacaacctt agagggtaca accttagtggt ttcttcgtt tactccactg 1080  
tatacaccccc atctgcttgg gctgtacatt gtctgaccat gcttataaca aaagtcacat 1140  
actactagcc aagactgaga acttagagcg actggccaga aagtaaagat acaacagttg 1200  
atatgtgtgc cacacacaga tccatgtgtatcatgatgc acgtgttttg 1260  
tggacatcctt cacaaggcag caggaaatg cttttttttt gctgtggac 1320  
accatatgac attgagatta cccgggtgcc cattccaaaca agagttataa gctccccctt 1380  
tggttgggtt ccagaaacctt gatttttttag caatagtcctt ctcacatcca gattaagagg 1440  
ggatggctt agcttagggtt actatgtatgc aactatgacc aaagcaactt gtggtaaaaa 1500  
gggtgttattt ggcttacact tccatatcac ttcatcaag tgaggacagg aactcaaata 1560  
gagtaggaat ttgggtgacaa gagctgtatgt agaggcaatg cagtgggtgcc acttagtggc 1620  
gcgcctcagtc tgctccctt cttaatagaa tgcaagacca ccagcccatg ggtggcacca 1680  
caatgggacc gggcccttcc ccatcggtca ctaagaaat gcccacatgc cagatctt 1740  
ggagacattt tctcaacggg ggcctactcc tttcagataa ctctatataa aattgacata 1800  
aaccagaaca gaggaggagg ctaagaagga aactgccaat tgcatacatg cacacacctg 1860  
gcccttagcag ctgcaggaaatg ctatgtttt atggcctttt ctcattttca tggaccagca 1920  
tgagcactct gcagagagatg atgcctgtatgc gcctgccaatg gcaggagtgc ttacactgaa 1980  
ggtaacagg atggcagggg ggcctcagag cttccaaatgt tcagaaccccc agcagaagag 2040  
ctgagaccct tgcccgagga ctcaggcggtt ttggaaaggc caggaaattt agccagagct 2100  
cttcttcaga tgggttacca tctgaagggtt agaccagata gccaactgtt gttgaggag 2160  
cacctctgca gcccctaccc ttggaaagata gaaagtgtct ctgtgacaag tatggccatt 2220  
gtgccccctt attccacatgtt caacagaaac ccttggaaatcc tgaacacttc tgcagctt 2280  
tttttacatgtt ctgcagggtt gctctaggaa tgaagggtgc cgagaggctt gggcgtaggc 2340  
aggtgacaag accacagttt ggggtcacag ctggcttactt ggatcactct tggacagagt 2400  
ttgttagata tggagtggag tatacacaag gcatcaggcg gggatattt aatgtatcac 2460  
cgagctccctt tggggcttgg cagccaagca cagcagttt tttgtttaaac aaatccacgg 2520  
ttccctcccccc ttgacgcagt acatctgtgg ctccaaaccc acacacccac ccattgttag 2580  
tgctggagac ttctacactac catgccagct ttggctatca tgggtctcag cctggctgct 2640  
ttccctggagc ttggatggg ggccttttgg tgcaggatggc agcaattcaa ggcacaagg 2700  
gactacatac tggggggctt attccctgtt ggctcaaccg aggaggccac tctcaaccag 2760  
agaacacaac ccaacacatccc cccgtgcaac aggtatggag gcttagtagct ggggtggag 2820  
tgaaccgaag cttggcagct ttggctccgtt ggtactacca atctggaaag aggtgggtat 2880  
cagttccat gtggctcag ttttttttgc ttggcttgc ttcctggccat ggctatgaaag 2940  
atggctgtgg aggagatcaa caatggatctt gcttgcctt ctgggtcg gctgggtat 3000  
gacctatttgc acacatgctc cgagccagtg gtcaccatga aatccagttt catgttctt 3060  
gccaagggtgg gcaatgttgc cattgtgttgc tactgtcaactt acacacatgtt gccaacccctt 3120  
gtgtggctg tcatcgcccccc ccaactcatca gagcttgcctt tcattacagg caagttctt 3180  
agcttcttcc tcatgccaca ggtgagccca cttcccttgc tttctcaacc gattgcaccc 3240  
attgagctctt catatcgaaa agtgcattt gatcaccacca ggtcagtttcatgatgc ctagccagca 3300  
tggatcggtt aagtgaccgg gaaacgtttt catttttttcc cccgtgacc 3360  
gggtgcagctt gcaaggcagttt gtcactctgtt tgcagaactt cagctggaaac tgggtggcc 3420  
ccttagggag tgcataatgc ttttttttttcatgatgc ctttttttttcatgatgc 3480  
atgcacgagg tatctgcatttgc acatgttgc gcaatgttgc acaacatgtt gccaacccctt 3540

aacagttggg caaggtgctg gatgtactac gccaaagtggaa ccaaaggtaaa gtacaagtgg 3600  
tggctgttgc tgcctctgcc cgtgctgtct actcccttt tagttacagc atccatcatg 3660  
gcctctcacc caaggtatgg gtggccagtg agtcttggct gacatctgac ctggcatg 3720  
caactcccaa tattgcccgt gtggccactg tgcttgggtt tttgcagcgg ggtgcccac 3780  
tgccctgaatt ttcccattat gtggagactc accttgcctt ggccgctgac ccagcattct 3840  
gtgcctcaact gaatgcggag ttggatctgg aggaacatgt gatggggcaa cgctgtccac 3900  
ggtgtgacga catcatgctg cagaacctat catctggct gttcagaac ctatcagctg 3960  
ggcaattgca ccaccaaata tttgcaacct atgcagctgt gtacagtgtg gtcagcccc 4020  
ttcacaacac cctacagtgc aatgtctcac attgccacgt atcagaacat gttctaccct 4080  
ggcaggtaaag ggtagggtt tttgctgggt tttgcctgtc cctgcaggaa cactgaacca 4140  
ggcagagcca aatcttggtg tgactggaga ggccttaccc tgactccact ccacagctcc 4200  
tggagaacat gtacaatatg agtttccatg ctcgagactt gacactacag tttgatgtgg 4260  
aaggaaatgt agacatggaa tatgacactg agatgtgggt gtggcagagc cctacacctg 4320  
tattacatac tggggcacc ttcaacggca cccttcagct gcagcagttt aaaaatgtact 4380  
ggccaggcaa ccaggttaagg acaagacagg caaaaaggat ggtggtaga agcttgcgg 4440  
tcttgggcca gtgctagcca aggggaggcc taacccaagg ctccatgtac aggtgccagt 4500  
ctccccagtgt tcccgccagt gcaaagatgg ccaggttcgc cgagtaaagg gctttcattc 4560  
ctgctgctat gactgcgtgg actgcaagggc gggcagctac cggaaagcatc caggtgaacc 4620  
gtcttcccta gacagtctgc acagccggc tagggggcag aagcattcaa gtctggcaag 4680  
cgccctcccg cggggctaat gtggagacag ttactgtggg ggctggctgg ggaggtcggt 4740  
ctccccatcag cagacccac attacttttcc ttccttccat cactacagat gacttcaccc 4800  
gtactccatg taaccaggac cagtggtccc cagagaaaaag cacagcctgc ttacctcgca 4860  
ggcccaagtt tctggcttgg ggggagccag ttgtgctgtc actccttcgt ctgctttgcc 4920  
tggtgctggg ttagactg gctgctctgg ggctctctgt ccaccactgg gacagccctc 4980  
ttgtccaggc ctcagggtggc tcacagttct gctttggct gatcgtccta ggccttct 5040  
gcctcagtgt cttctgttcc ccaaggccggc caagctctgc cagctgcctt gcacaacaac 5100  
caatggctca cttccctctc acaggctgcc tgagcacact cttctgcac gcaactgaga 5160  
ccttggta gtctgagctg ccactgagct gggcaaaactg gctatgcagc taccttcggg 5220  
gactctggc ctggctagt gtaactgtgg ccactttgt ggaggcagca ctatgtgcct 5280  
ggtatttgat cgctttccca ccagaggtgg tgacagactg gtcagtgctg cccacagagg 5340  
tactggagca ctgcccacgtg cgttccctggg tcagcctggg cttggtgac atcacaatg 5400  
caatgttagc tttcctctgc tttctggca ctttcctgtt acagagccag cctggccgct 5460  
acaaccgtgc ccgtggctc accttcgcca tgctagctt tttcatcacc tgggtctctt 5520  
ttgtgccccct cctggccaat gtgcagggtgg cttaccagcc agctgtgcag atgggtgcta 5580  
tcctagtctg tgccctggc atctggta cttccaccc gccaaggatgc tatgtgcttc 5640  
tttggctgcc aaagctcaac acccaggagt tttcctggg aaggaatgcc aagaaagcag 5700  
cagatgagaa cagtggcggt ggtggaggcag ctcaggacaa caatgaatga ccactgaccc 5760  
gtgaccccttc ctttagggaa cttagcccta ccagaaatct cctaagccaa caagccccga 5820  
atagtacctc agcctgagac gtgagacact taactataga cttggactcc actgaccccta 5880  
gcctcacaatg gaccccttcc ccaaacccttcc aaggccctgca gtgcacaaga tggaccctat 5940  
gagccccaccc atccttcaa agcaagatta tccttgatcc tattatgccc acctaaggcc 6000  
tgcccagggtt acccacaaaaa ggttcttgg gacttcatacc ccatactttg aattcagaaa 6060  
ttccccaggc agaccatggg agaccagaag gtactgttgc cctgaacatg cccagccctg 6120  
agccctcact cagcacccctg tccaggcgtc ccaggaatag aaggctggc atgtatgtgt 6180  
gtgtgtgtgt gtgtgtgtgt gtgtgtgtat gtacgtatgt atgtatgtat 6240  
caggacagaa caagaaagac atcaggcaga ggacactcag gaggtaggca acatccagcc 6300  
ttctccatcc cttagctgac cctagccctgtt aggagagaac caggtcgccg ccagcaccc 6360  
ggacagatca cacacagggt gcccgtcagc accacggccca gcccagccca cgcgggaccc 6420

ctggaatcag cttctagtag caaggacaga aaagttgccg caaggccccct tactggccag 6480  
 caccaggacat agagccacat gccttaagcg 6540  
 caggatcaga cccgggtcag ttctggactg gcccccacac ctgaatccccg gagcagctca 6600  
 gctggagaaa agagaaacaa gccacacatc agtcccataa aattaaacgc ttttttagt 6660  
 gttaaaata gcatttacac agaagcagca tttacacaga agcagotcta tgtcaactac 6720  
 ccagtcaactc agactttgac acagtgtcta gttagatgt gtggggccgc tggccggga 6780  
 tggcagtggc acatgatgat gggcagccac cagaacagaa acagaacagg gcccagctct 6840  
 gcagctcttgc tggcactgt caccaccac tgagactgag acagtggtca ggtgccaggt 6900  
 ctctctctgt tctctcttac tagctaccct tcacataacct tcagtacaaa ctgtgttgc 6960  
 atgtccaag tagcaggtgg ggaaaggggc atgcaaactg cccctttggg taactagctg 7020  
 ccacccttag agcaggcagg ctagcaataa ataaataagt tagacccac ctggcagcc 7080  
 agagagggtt gaaggctctg tctaaccctt caaaaatccc accttggcct gacaggtgag 7140  
 gcccattgaac ttagcgcacag tcagctgtg tccctgtgca cagttctgtg aggctttggg 7200  
 gcaaggggta ccaagagccc aagagagct ttcttgcctt aatggaggt cacttccaaa 7260  
 gaagggaccaggagggtt ccctgagact tggctgagg acttaaagt agagatgtct 7320  
 ctttacaaga ctctatagat acttgagctg taccaccatc agcagccca agagcagaca 7380  
 aaatgtcaag ccaatatccctt ggtgttatgg ctggccctcag gcccctctct gtggcctgt 7440  
 ccctctgccc tggcccaagag cccacagctg atctatctgt gctggccacc accacggcc 7500  
 ggcagagct cctggcacag caggagcaca gactcagcca caggcagcgc tgaagacatt 7560  
 ggttgcacat cacatgatgt ccacaaagaa ctcacagggg tttccatgg cttttggaa 7620  
 ggactggcgg ctacctgtaa gttctggagg gacagcagcc agctcccgga cgggtggccc 7680  
 tccagggtggc ccacccacta ctgcataaggc ctttgcatttggg ggggtgcagtgggggagcc 7740  
 tggggcaaca gctgaaggctt gacttcgagg gctactgcac cggctaagct ggctgacagg 7800  
 ccgctccac cagccgggtgc taccagaccc acttggtaact gtgtggctgtt attcaactgccc 7860  
 actacccca gctccagttt cccggcgctc ctctcgccctt ggggtccgat ggctgctcog 7920  
 tggggcccaactt ctgcattttgc tcccttagggg gagggaagg gacaacagag tggcactgag 7980  
 gctggccac ttccaggggc accagctgtt cccagacagt cagggcagga cctggtaagc 8040  
 ctggagatgg taggggaatg gcagccatgc agataccagg aacagctgag aggccgagaag 8100  
 cttagggcag tggcagacag caggacaaac agggggccacg ctggccaccc acacctaacc 8160  
 ccaatgctt aaccaagggt taatgttaca gctgagaaac taaaaaccag cgaaggccct 8220  
 gtgtgcccag cattccattt agccatcctg gtttgcaccac ccaaagaccc aaccagggtc 8280  
 caccacccca caggaccctg gtcattttat ttgcattttgc cctgtccctga aagttagtggg 8340  
 aacctgaaaa cacgtgttgc ctggggacat gctgagaggg acacaggggg acctggctt 8400  
 ccggcccgag agtccactct gctgttgcctt cagttcaagg cttgttgcacg acaaagcaag 8460  
 ggatagcaca agtcacacac cagttccagtg ctccaccaatg gtaatagga cgattttggg 8520  
 ccaagctgag cctgggtaca tggcagggcc tggccatggt caggattcac tggatagctt 8580  
 ccccttggc tttggccaccc tctggcccaa cctctccctga gtctttctctt ggaccttgc 8640  
 gcacaagtgt gccccactct gcttaagacc tccacatcag tccatctctt cctggggac 8700  
 acccaccctt caagatctt aatatccctg ggatatgtt taacactgat atgcttttac 8760  
 agtggctt gatacttta tctggcactc tgggtggatg caggctccat aactgataaa 8820  
 gcccatttctc ccccttagctt gggccctaga gaggccctt acctgttacg agtggcttact 8880  
 ttcatcttgc ccatatcattc tcctggcctc ttgccttgc caccttagcac accaggctgt 8940  
 ctccatttgc tctaaacggct tctaccacatc tcagcccttc cctgtccctac acactgactc 9000  
 ttgagatggc acccaccggg actcaaacac acaggcaggag cacagaggga agcgtcgccc 9060  
 ccaggcagag cgtgggagtg ggagggagtg ggaggagggg tggcacgcctt ctcacccatc 9120  
 ctctgtggc tcccagactt gccgttgcgg cagttcaagc cagggttgcctt gtaagcaggc 9180  
 gggaaaggcagg gcccgggtcc tgggtactgg taggggtacg tttgacccaa gggccagggt 9240  
 actgtgggtt gggccagttgg gggccacttgc tcctgttgc aggtccactt ggagccactt 9300

ttgggttca gggatgcgag gtctggcagg gagggaggga gggagggta agtgaaggca 9360  
aatgaatgag gccacagcaa ccctacccaa cgcacccct actcaactact gcacaggtcg 9420  
ccaaagacat agtagcaactg ctcagaaaag gtgatcttgc tCACGGTGTG cctcaggaaa 9480  
ccgtgctca gcatactgt ggcatacttt ctgcctccc ttgcctcctt gaagccctcc 9540  
acgtgtgtgt acagccagtc caccacatcc gcccctggcc acaggtccat caaagtcagg 9600  
gtagctgagc cctgggaagc tacgcccagaa tgaggaacag acggggccct tcccacacag 9660  
ccagggactc accaatgaca gcattggcaa tggtgatctt aagccacatc cggtcccgga 9720  
tctccagtc tgagtctggc aactgcata ggcggacaat ggcactcatg tcactcttca 9780  
cagtcagcgg tgcctcctca agctctgcag agcacacttc cctgagccca ggctcacagc 9840  
gtgaacctcc atggggttga gaggcaggggc cagggtcaaa ccttttatct cccatccttgc 9900  
ggagatgccc ctcatcgaaa cttgagctaa gacggggaga ttcttcccg tcccacagtg 9960  
caagtccacg taggcaaggc agccccccctc ccctccccgg agagaacaag ctgttagcta 10020  
tgtaggttag cagaaaagca aagcagaggc tgccatgtcc tcccaattcc cccctccgca 10080  
caggcctggc aggaccctca attcatgcag atgaccagta tggccaggcc tggagggata 10140  
tgtacatgta tctttgtta cacatttgc aagggttgg aagcaaaacaa aacccatcata 10200  
tgtaatggc ccctgtataa gctctgatga gcaccaaagc tcaaagctag aactgaccat 10260  
tgtccttcaa cctcagttc cttgggtggg ggggggtcct gtgagctgcc acttacgtgg 10320  
ggcgcggcaggc actgagctgg ttagtgagga agagctggc cgtgtgatgg cgctggagca 10380  
gggactcgta ccatacgcccc gcagggcacc cgtcagtgtc gctgtgtggg acagccagggc 10440  
agccgggtcg atgggtcgca ctgggtcagc tgcatagttt ccacagcaac ggattacagg 10500  
tggtaagtag gggggcagca cagaggcaga caagaaaagac ccccaagactg aacacagaaa 10560  
ccccaccccta ccccaccttt ccatagggta actcacccct tggatgggt aagtagctcc 10620  
gaggggttgg gtcccagcac ttggccactg tgagactgat gggctacag agttgagcag 10680  
accatgtgt aagtggggcc cgccacagcccc ctcccatcct gtgccactcc caccccaact 10740  
tggctcccac ctcaacctgt ctgggacacg atctcccgaa gcacccgtac agcgtcgtca 10800  
ttgctcatgt tctcaaagtt gacatgttc acctacgggg ttgtgggggt caggggttgg 10860  
tggtggtatg tggtgccctc ttgtccccac agtccccaca tggctccac ctgcagcaac 10920  
atgtcgcccc gctcaatgcg gccatcagca gccaaggcccc cggccatcat gatggatcca 10980  
atgttagatgc cgccatcacc cgggtcggtt ctctggccca cgatgctgat gcccaggaaag 11040  
tggtgccctct ctgcaggagg ggccgtgagc aggccccaa agctccccag gctgtacccca 11100  
cccccagcag gcacccacag cccacaaggc ctcacccatg ttgagactgat cgggtgatgat 11160  
gttcaggagc atgtggagt ctgtgatgt gctgaaggag gatgcctgcg gagggaccca 11220  
gtgaggggtct gtgtgggcac cattcagac agacacccca cccacccgtc gcctacccgg 11280  
tctgtctgcc tcaagcgctg ctggcagca cggcatttgc gttccgaac tagccgagag 11340  
gaggtgctct gctctgtggc gctgtcagc ctggggcagg agtcagaaaa gcacaaacat 11400  
gtataaccag ctggacgcgt caactacaaa tctccagcac gtactgacat gtgcacacgt 11460  
caccacccgg ctgtattgt ctcctcatc tgagtcaata aagctgctag attcaagctc 11520  
actgctcagt acagtggatg cactgtctgg aggtgtcccc aggtcccccc gccgatcccc 11580  
tctcgggtgc ccattggtcc gggcagctgt gggacagta ggggtgggtac gactgtggga 11640  
cttcagtcct aacagaatgc ggggtggcctg tgcatttcaa agtttatgca gtaactctgg 11700  
ggccacaggc gctaggagta ccaggctggg acctctaccc aaggatcaact gcttggaaaga 11760  
atatgtggaa tacccatccagg ctggagatgat accaaaggga taccaaggg 11800

<210> 3  
<211> 858  
<212> PRT  
<213> Mouse

&lt;400&gt; 3

Met Pro Ala Leu Ala Ile Met Gly Leu Ser Leu Ala Ala Phe Leu Glu  
1 5 10 15

Leu Gly Met Gly Ala Ser Leu Cys Leu Ser Gln Gln Phe Lys Ala Gln  
20 25 30

Gly Asp Tyr Ile Leu Gly Gly Leu Phe Pro Leu Gly Ser Thr Glu Glu  
35 40 45

Ala Thr Leu Asn Gln Arg Thr Gln Pro Asn Ser Ile Pro Cys Asn Arg  
50 55 60

Phe Ser Pro Leu Gly Leu Phe Leu Ala Met Ala Met Lys Met Ala Val  
65 70 75 80

Glu Glu Ile Asn Asn Gly Ser Ala Leu Leu Pro Gly Leu Arg Leu Gly  
85 90 95

Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val Thr Met Lys Ser  
100 105 110

Ser Leu Met Phe Leu Ala Lys Val Gly Ser Gln Ser Ile Ala Ala Tyr  
115 120 125

Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala Val Ile Gly Pro  
130 135 140

His Ser Ser Glu Leu Ala Leu Ile Thr Gly Lys Phe Phe Ser Phe Phe  
145 150 155 160

Leu Met Pro Gln Val Ser Tyr Ser Ala Ser Met Asp Arg Leu Ser Asp  
165 170 175

Arg Glu Thr Phe Pro Ser Phe Phe Arg Thr Val Pro Ser Asp Arg Val  
180 185 190

Gln Leu Gln Ala Val Val Thr Leu Leu Gln Asn Phe Ser Trp Asn Trp  
195 200 205

Val Ala Ala Leu Gly Ser Asp Asp Asp Tyr Gly Arg Glu Gly Leu Ser  
210 215 220

Ile Phe Ser Ser Leu Ala Asn Ala Arg Gly Ile Cys Ile Ala His Glu  
225 230 235 240

Gly Leu Val Pro Gln His Asp Thr Ser Gly Gln Gln Leu Gly Lys Val

245

250

255

Leu Asp Val Leu Arg Gln Val Asn Gln Ser Lys Val Gln Val Val Val  
 260 265 270

Leu Phe Ala Ser Ala Arg Ala Val Tyr Ser Leu Phe Ser Tyr Ser Ile  
 275 280 285

His His Gly Leu Ser Pro Lys Val Trp Val Ala Ser Glu Ser Trp Leu  
 290 295 300

Thr Ser Asp Leu Val Met Thr Leu Pro Asn Ile Ala Arg Val Gly Thr  
 305 310 315 320

Val Leu Gly Phe Leu Gln Arg Gly Ala Leu Leu Pro Glu Phe Ser His  
 325 330 335

Tyr Val Glu Thr His Leu Ala Leu Ala Asp Pro Ala Phe Cys Ala  
 340 345 350

Ser Leu Asn Ala Glu Leu Asp Leu Glu Glu His Val Met Gly Gln Arg  
 355 360 365

Cys Pro Arg Cys Asp Asp Ile Met Leu Gln Asn Leu Ser Ser Gly Leu  
 370 375 380

Leu Gln Asn Leu Ser Ala Gly Gln Leu His His Gln Ile Phe Ala Thr  
 385 390 395 400

Tyr Ala Ala Val Tyr Ser Val Ala Gln Ala Leu His Asn Thr Leu Gln  
 405 410 415

Cys Asn Val Ser His Cys His Val Ser Glu His Val Leu Pro Trp Gln  
 420 425 430

Leu Leu Glu Asn Met Tyr Asn Met Ser Phe His Ala Arg Asp Leu Thr  
 435 440 445

Leu Gln Phe Asp Ala Glu Gly Asn Val Asp Met Glu Tyr Asp Leu Lys  
 450 455 460

Met Trp Val Trp Gln Ser Pro Thr Pro Val Leu His Thr Val Gly Thr  
 465 470 475 480

Phe Asn Gly Thr Leu Gln Leu Gln Gln Ser Lys Met Tyr Trp Pro Gly  
 485 490 495

Asn Gln Val Pro Val Ser Gln Cys Ser Arg Gln Cys Lys Asp Gly Gln

500	505	510
Val Arg Arg Val Lys Gly Phe His Ser Cys Cys Tyr Asp Cys Val Asp		
515	520	525
Cys Lys Ala Gly Ser Tyr Arg Lys His Pro Asp Asp Phe Thr Cys Thr		
530	535	540
Pro Cys Asn Gln Asp Gln Trp Ser Pro Glu Lys Ser Thr Ala Cys Leu		
545	550	555
Pro Arg Arg Pro Lys Phe Leu Ala Trp Gly Glu Pro Val Val Leu Ser		
565	570	575
Leu Leu Leu Leu Leu Cys Leu Val Leu Gly Leu Ala Leu Ala Ala Leu		
580	585	590
Gly Leu Ser Val His His Trp Asp Ser Pro Leu Val Gln Ala Ser Gly		
595	600	605
Gly Ser Gln Phe Cys Phe Gly Leu Ile Cys Leu Gly Leu Phe Cys Leu		
610	615	620
Ser Val Leu Leu Phe Pro Gly Arg Pro Ser Ser Ala Ser Cys Leu Ala		
625	630	635
640		
Gln Gln Pro Met Ala His Leu Pro Leu Thr Gly Cys Leu Ser Thr Leu		
645	650	655
Phe Leu Gln Ala Ala Glu Thr Phe Val Glu Ser Glu Leu Pro Leu Ser		
660	665	670
Trp Ala Asn Trp Leu Cys Ser Tyr Leu Arg Gly Leu Trp Ala Trp Leu		
675	680	685
Val Val Leu Leu Ala Thr Phe Val Glu Ala Ala Leu Cys Ala Trp Tyr		
690	695	700
Leu Ile Ala Phe Pro Pro Glu Val Val Thr Asp Trp Ser Val Leu Pro		
705	710	715
720		
Thr Glu Val Leu Glu His Cys His Val Arg Ser Trp Val Ser Leu Gly		
725	730	735
Leu Val His Ile Thr Asn Ala Met Leu Ala Phe Leu Cys Phe Leu Gly		
740	745	750
Thr Phe Leu Val Gln Ser Gln Pro Gly Arg Tyr Asn Arg Ala Arg Gly		

755

760

765

Leu Thr Phe Ala Met Leu Ala Tyr Phe Ile Thr Trp Val Ser Phe Val  
 770 775 780

Pro Leu Leu Ala Asn Val Gln Val Ala Tyr Gln Pro Ala Val Gln Met  
 785 790 795 800

Gly Ala Ile Leu Val Cys Ala Leu Gly Ile Leu Val Thr Phe His Leu  
 805 810 815

Pro Lys Cys Tyr Val Leu Leu Trp Leu Pro Lys Leu Asn Thr Gln Glu  
 820 825 830

Phe Phe Leu Gly Arg Asn Ala Lys Lys Ala Ala Asp Glu Asn Ser Gly  
 835 840 845

Gly Gly Glu Ala Ala Gln Gly His Asn Glu  
 850 855

&lt;210&gt; 4

&lt;211&gt; 2559

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

atgctgggcc ctgctgtcct gggcctcagc ctctgggctc tcctgcaccc tgggacgggg 60  
 gccccattgt gcctgtcaca gcaacttagg atgaaggggg actacgtgt gggggggctg 120  
 ttccccctgg gcgaggccga ggaggctggc ctccgcagcc ggacacggcc cagcagccct 180  
 gtgtgcacca ggttctcctc aaacggcctg ctctggcac tggccatgaa aatggccgtg 240  
 gaggagatca acaacaagtc ggtatctgtg cccgggctgc gcctgggcta cgacctctt 300  
 gatacgtgtg cggagccgt ggtggccatg aagcccagcc tcatgttccct ggccaaggca 360  
 ggcagccgca acatgcgcgc ctactgcaac tacacgcagt accagccccg tttgtgtggct 420  
 gtcatcgccc cccactcgtc agagctcgcc atggtcaccc gcaagttctt cagttcttc 480  
 ctcatgcccc aggtcagcta cgggtctagc atggagctgc tgagcccccgg gtagaccccttc 540  
 ccctcccttc tccgcaccgt gcccagcgcac cgtgtgcagc tgacggccgc cgccggagctg 600  
 ctgcaggagt tcggctggaa ctgggtggcc gcccctggca gcgacgcacga gtacggccgg 660  
 cagggcctga gcatcttc gcccctggcc tcggcacgcg gcatctgcac cgcgcacgag 720  
 ggccctggc cgctgcggcccg tgccgatgac tcgcggctgg ggaagggtgca ggacgttctg 780  
 caccaggtga accagagcag cgtgcagggt gtgcgtctgt tcgcctccgt gcacgcgc 840  
 cacgcctct tcaactacag catcagcagc aggctctcgcc ccaagggtgtg ggtggccagc 900  
 gaggcctggc tgacctctga cctgggtcatg gggctgcccc gcatggccca gatggggcagc 960  
 gtgcgttggct tcctccagag ggggtccccag ctgcacgcgt tccccccatgta cgtgaagacgc 1020  
 cacctggccc tggccaccga cccggcccttc tgctctgccc tggggcagag ggagcagggt 1080  
 ctggaggagg acgtgggtggg ccagcgcgtgc cccgcgtgtg actgcacac gctgcagaac 1140  
 gtgagcgcag ggctaaatca ccaccagacg ttctctgtct acgcagctgt gtatagcgtg 1200  
 gcccaggccc tgcacaacac tcttcagtgc aacgcctcag gctgccccgc gcaggacccc 1260

gtgaagccct ggcagctcct ggagaacatg tacaaccctga cttccacgt gggcggt 1320  
 ccgctgcgt tcgacagcag cggaaacgtg gacatggagt acgacacgtaa gctgtgggtg 1380  
 tggcaggcgt cagtgcacag gctccacgac gtggcaggt tcaacggcag cctcaggaca 1440  
 gagegcctga agatccgctg gcacacgtct gacaaccaga agccgtgtc cccgtgtcg 1500  
 cggcagtgcc aggagggcca ggtgcgcgg gtcaagggtt tccactcctg ctgctacgac 1560  
 tgtgtggact gcgaggcggg cagttaccgg caaaacccag acgacatcgc ctgcacctt 1620  
 tgtggccagg atgagtgtc cccggagcga agcacacgtt gctccgcgg caggtctcgg 1680  
 ttccctggcat gggcgagcc ggctgtgtc ctgctgtcc tgctgtgag cctggcgctg 1740  
 ggccttgc tggctgttt gggctgttc gttcaccatc gggacagccc actggttcag 1800  
 gcctcggtt ggcctctggc ctgcttgc ctggctgtcc tggcctgtt ctgcctcagc 1860  
 gtccctctgt tccctggcca gcccagccct gcccgtatgcc tggccagca gccctgtcc 1920  
 cacctccgc tcacgggtgt cctgagcaca ctcttcctgc aggccggcga gatcttcgtg 1980  
 gagtcagaac tgcctctgag ctggcagac cggctgagtg gctgcctgcg gggccctgg 2040  
 gcctggctgg tggctgtgtt ggcctgtc gtggaggtcg cactgtgcac ctggtacctg 2100  
 gtggccttcc cgccggaggt ggtgacggac tggcacatgc tgccacaggaa ggcgtgggtg 2160  
 cactgcccga cactgctcctg ggtcagctt ggcctagcgc acgcccaccaa tgccacgcgt 2220  
 gccttctct tccctctggg cacttccctg gtgcggagcc agccggcccg atacaaccgt 2280  
 gcccgtggcc tcacctttgc catgtggcc tacttcatca cctgggttcc ctttgtgccc 2340  
 ctccctggcca atgtgcaggt ggtcctcagg cccggcgtgc agatggcgc ctcctgtct 2400  
 tgtgtcctgg gcatcctggc tgccctccac ctgcccaggt gttacctgtc catgcggcag 2460  
 ccagggctca acaccccccga gttcttcctg ggagggggcc ctggggatgc ccaaggccag 2520  
 aatgacggga acacaggaaa tcagggaaa catgagtga 2559

&lt;210&gt; 5

&lt;211&gt; 852

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

Met Leu Gly Pro Ala Val Leu Gly Leu Ser Leu Leu Trp Ala Leu Leu His

1

5

10

15

Pro Gly Thr Gly Ala Pro Leu Cys Leu Ser Gln Gln Leu Arg Met Lys

20

25

30

Gly Asp Tyr Val Leu Gly Leu Phe Pro Leu Gly Glu Ala Glu Glu

35

40

45

Ala Gly Leu Arg Ser Arg Thr Arg Pro Ser Ser Pro Val Cys Thr Arg

50

55

60

Phe Ser Ser Asn Gly Leu Leu Trp Ala Leu Ala Met Lys Met Ala Val

65

70

75

80

Glu Glu Ile Asn Asn Lys Ser Asp Leu Leu Pro Gly Leu Arg Leu Gly

85

90

95

Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val Ala Met Lys Pro  
100 105 110

Ser Leu Met Phe Leu Ala Lys Ala Gly Ser Arg Asp Ile Ala Ala Tyr  
115 120 125

Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala Val Ile Gly Pro  
130 135 140

His Ser Ser Glu Leu Ala Met Val Thr Gly Lys Phe Phe Ser Phe Phe  
145 150 155 160

Leu Met Pro Gln Val Ser Tyr Gly Ala Ser Met Glu Leu Leu Ser Ala  
165 170 175

Arg Glu Thr Phe Pro Ser Phe Phe Arg Thr Val Pro Ser Asp Arg Val  
180 185 190

Gln Leu Thr Ala Ala Ala Glu Leu Leu Gln Glu Phe Gly Trp Asn Trp  
195 200 205

Val Ala Ala Leu Gly Ser Asp Asp Glu Tyr Gly Arg Gln Gly Leu Ser  
210 215 220

Ile Phe Ser Ala Leu Ala Ser Ala Arg Gly Ile Cys Ile Ala His Glu  
225 230 235 240

Gly Leu Val Pro Leu Pro Arg Ala Asp Asp Ser Arg Leu Gly Lys Val  
245 250 255

Gln Asp Val Leu His Gln Val Asn Gln Ser Ser Val Gln Val Val Leu  
260 265 270

Leu Phe Ala Ser Val His Ala Ala His Ala Leu Phe Asn Tyr Ser Ile  
275 280 285

Ser Ser Arg Leu Ser Pro Lys Val Trp Val Ala Ser Glu Ala Trp Leu  
290 295 300

Thr Ser Asp Leu Val Met Gly Leu Pro Gly Met Ala Gln Met Gly Thr  
305 310 315 320

Val Leu Gly Phe Leu Gln Arg Gly Ala Gln Leu His Glu Phe Pro Gln  
325 330 335

Tyr Val Lys Thr His Leu Ala Leu Ala Thr Asp Pro Ala Phe Cys Ser  
340 345 350

Ala Leu Gly Glu Arg Glu Gln Gly Leu Glu Glu Asp Val Val Gly Gln  
355 360 365

Arg Cys Pro Gln Cys Asp Cys Ile Thr Leu Gln Asn Val Ser Ala Gly  
370 375 380

Leu Asn His His Gln Thr Phe Ser Val Tyr Ala Ala Val Tyr Ser Val  
385 390 395 400

Ala Gln Ala Leu His Asn Thr Leu Gln Cys Asn Ala Ser Gly Cys Pro  
405 410 415

Ala Gln Asp Pro Val Lys Pro Trp Gln Leu Leu Glu Asn Met Tyr Asn  
420 425 430

Leu Thr Phe His Val Gly Gly Leu Pro Leu Arg Phe Asp Ser Ser Gly  
435 440 445

Asn Val Asp Met Glu Tyr Asp Leu Lys Leu Trp Val Trp Gln Gly Ser  
450 455 460

Val Pro Arg Leu His Asp Val Gly Arg Phe Asn Gly Ser Leu Arg Thr  
465 470 475 480

Glu Arg Leu Lys Ile Arg Trp His Thr Ser Asp Asn Gln Lys Pro Val  
485 490 495

Ser Arg Cys Ser Arg Gln Cys Gln Glu Gly Gln Val Arg Arg Val Lys  
500 505 510

Gly Phe His Ser Cys Cys Tyr Asp Cys Val Asp Cys Glu Ala Gly Ser  
515 520 525

Tyr Arg Gln Asn Pro Asp Asp Ile Ala Cys Thr Phe Cys Gly Gln Asp  
530 535 540

Glu Trp Ser Pro Glu Arg Ser Thr Arg Cys Phe Arg Arg Arg Ser Arg  
545 550 555 560

Phe Leu Ala Trp Gly Glu Pro Ala Val Leu Leu Leu Leu Leu Leu  
565 570 575

Ser Leu Ala Leu Gly Leu Val Leu Ala Ala Leu Gly Leu Phe Val His  
580 585 590

His Arg Asp Ser Pro Leu Val Gln Ala Ser Gly Gly Pro Leu Ala Cys  
595 600 605

Phe Gly Leu Val Cys Leu Gly Ieu Val Cys Leu Ser Val Leu Leu Phe  
610 615 620

Pro Gly Gln Pro Ser Pro Ala Arg Cys Leu Ala Gln Gln Pro Leu Ser  
625 630 635 640

His Leu Pro Leu Thr Gly Cys Leu Ser Thr Leu Phe Leu Gln Ala Ala  
645 650 655

Glu Ile Phe Val Glu Ser Glu Leu Pro Leu Ser Trp Ala Asp Arg Leu  
660 665 670

Ser Gly Cys Leu Arg Gly Pro Trp Ala Trp Leu Val Val Leu Leu Ala  
675 680 685

Met Leu Val Glu Val Ala Leu Cys Thr Trp Tyr Leu Val Ala Phe Pro  
690 695 700

Pro Glu Val Val Thr Asp Trp His Met Leu Pro Thr Glu Ala Leu Val  
705 710 715 720

His Cys Arg Thr Arg Ser Trp Val Ser Phe Gly Leu Ala His Ala Thr  
725 730 735

Asn Ala Thr Leu Ala Phe Leu Cys Phe Leu Gly Thr Phe Leu Val Arg  
740 745 750

Ser Gln Pro Gly Arg Tyr Asn Arg Ala Arg Gly Leu Thr Phe Ala Met  
755 760 765

Leu Ala Tyr Phe Ile Thr Trp Val Ser Phe Val Pro Leu Leu Ala Asn  
770 775 780

Val Gln Val Val Leu Arg Pro Ala Val Gln Met Gly Ala Leu Leu Leu  
785 790 795 800

Cys Val Leu Gly Ile Leu Ala Ala Phe His Leu Pro Arg Cys Tyr Leu  
805 810 815

Leu Met Arg Gln Pro Gly Leu Asn Thr Pro Glu Phe Phe Leu Gly Gly  
820 825 830

Gly Pro Gly Asp Ala Gln Gly Gln Asn Asp Gly Asn Thr Gly Asn Gln  
835 840 845

Gly Lys His Glu  
850

<210> 6  
<211> 20  
<212> DNA  
<213> Mouse

<400> 6  
cactagagct gccacacctcc

20

<210> 7  
<211> 20  
<212> DNA  
<213> Mouse

<400> 7  
ccctcagcac cacttttgt

20

<210> 8  
<211> 20  
<212> DNA  
<213> Mouse

<400> 8  
acaaaaaagtg gtgctgaggg

20

<210> 9  
<211> 20  
<212> DNA  
<213> Mouse

<400> 9  
caggagaccc aaaggatcaa

20

<210> 10  
<211> 20  
<212> DNA  
<213> Mouse

<400> 10  
gcttcagaaa atcgaggcac

20

<210> 11  
<211> 20

<212> DNA

<213> Mouse

<400> 11

gcatgggcta tgatagggtgg

20

<210> 12

<211> 16

<212> DNA

<213> Mouse

<400> 12

tgttgatccc acagcgt

16

<210> 13

<211> 20

<212> DNA

<213> Mouse

<400> 13

caggaaatgt ccacttctgc

20

<210> 14

<211> 18

<212> DNA

<213> Mouse

<400> 14

tctatcttgc atccagcc

18

<210> 15

<211> 16

<212> DNA

<213> Mouse

<400> 15

gtgctgtgac tgtgcgt

16

<210> 16

<211> 18

<212> DNA

<213> Mouse

<400> 16  
cgcagcattt atttggag

18

<210> 17  
<211> 19  
<212> DNA  
<213> Mouse

<400> 17  
ccgacccttt aggagacac

19

<210> 18  
<211> 20  
<212> DNA  
<213> Mouse

<400> 18  
tgtgacttcc tcttcccac

20

<210> 19  
<211> 20  
<212> DNA  
<213> Mouse

<400> 19  
tgagccactc cagatgtcag

20

<210> 20  
<211> 20  
<212> DNA  
<213> Mouse

<400> 20  
ccaacgtgca gtcaagaaaa

20

<210> 21  
<211> 20  
<212> DNA  
<213> Mouse

<400> 21  
ccaacgtgca gtcaagaaaa

20

<210> 22  
<211> 20  
<212> DNA  
<213> Mouse

<400> 22  
cgagagacaa agtggtgctg

20

<210> 23  
<211> 20  
<212> DNA  
<213> Mouse

<400> 23  
ttatgaaggc cctcaccaac

20

<210> 24  
<211> 20  
<212> DNA  
<213> Mouse

<400> 24  
ccagctccta gaattgcctg

20

<210> 25  
<211> 20  
<212> DNA  
<213> Mouse

<400> 25  
gcagtctccc gaaacaagtc

20

<210> 26  
<211> 20  
<212> DNA  
<213> Mouse

<400> 26  
atagaggaat gggtgcgatg

20

<210> 27  
<211> 20

<212> DNA

<213> Mouse

<400> 27

taccaggagg ggtcagtcag

20

<210> 28

<211> 20

<212> DNA

<213> Mouse

<400> 28

tacaagcgag ctgaccaatg

20

<210> 29

<211> 20

<212> DNA

<213> Mouse

<400> 29

ccaaatcagct cgagtttagcc

20

<210> 30

<211> 20

<212> DNA

<213> Mouse

<400> 30

tgccattgtg gatgttcact

20

<210> 31

<211> 20

<212> DNA

<213> Mouse

<400> 31

gagtccgagg tcgggtcaata

20

<210> 32

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 32

gctggcttct gtagggtcagg

20

&lt;210&gt; 33

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 33

tatgagggtc aagggtcagg.

20

&lt;210&gt; 34

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 34

cgctttggtg agaactagcc

20

&lt;210&gt; 35

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 35

catgtggagt tgtggagtg

20

&lt;210&gt; 36

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 36

aatgggcaga agacagatgg

20

&lt;210&gt; 37

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 37

tatcagggtc tgtgaagccc

20

<210> 38  
<211> 20  
<212> DNA  
<213> Mouse

<400> 38  
atacaggacc ctttaccccg

20

<210> 39  
<211> 20  
<212> DNA  
<213> Mouse

<400> 39  
cagtgttctt aggtccccca

20

<210> 40  
<211> 20  
<212> DNA  
<213> Mouse

<400> 40  
gcctctgtctt gccatctctc

20

<210> 41  
<211> 20  
<212> DNA  
<213> Mouse

<400> 41  
ataatgttac ctgcaggcgg

20

<210> 42  
<211> 20  
<212> DNA  
<213> Mouse

<400> 42  
ctggaaacac ccatgtcctc

20

<210> 43  
<211> 20

<212> DNA

<213> Mouse

<400> 43

cggcacatg gacacttta

20

<210> 44

<211> 20

<212> DNA

<213> Mouse

<400> 44

gagcatgaag tgcaagggtga

20

<210> 45

<211> 20

<212> DNA

<213> Mouse

<400> 45

cgttaggtggc acagttgaga

20

<210> 46

<211> 20

<212> DNA

<213> Mouse

<400> 46

gctgttagtg aggtcagggc

20

<210> 47

<211> 20

<212> DNA

<213> Mouse

<400> 47

cgttaggtggc acagttgaga

20

<210> 48

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 48

gagcatgaag tgcaaggtga

20

&lt;210&gt; 49

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 49

tcatttcctt agcctcggtg

20

&lt;210&gt; 50

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 50

tctaagaaga tgatgcagac cc

22

&lt;210&gt; 51

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 51

tgcccttcag ggatagtgcc

20

&lt;210&gt; 52

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 52

ggcttcagcc tcaagttctg

20

&lt;210&gt; 53

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 53

aaaacaacca agttgccctg

20

<210> 54  
<211> 20  
<212> DNA  
<213> Mouse

<400> 54  
ggcactgaaa tgacctggat

20

<210> 55  
<211> 20  
<212> DNA  
<213> Mouse

<400> 55  
aacaattcaa gcaacacctgg

20

<210> 56  
<211> 20  
<212> DNA  
<213> Mouse

<400> 56  
ctgttccttc ccagactcca

20

<210> 57  
<211> 20  
<212> DNA  
<213> Mouse

<400> 57  
ttcagtcacg caaacacctgag

20

<210> 58  
<211> 20  
<212> DNA  
<213> Mouse

<400> 58  
gcccaggact ttgtcactgt

20

<210> 59  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 59

ggtaacctgc agctccactc

20

&lt;210&gt; 60

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 60

gggacatgct cttggttcat

20

&lt;210&gt; 61

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 61

gaacaaagcc gggtgattta

20

&lt;210&gt; 62

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 62

gccctcagtt ctcctagcct

20

&lt;210&gt; 63

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 63

ggcagagaag actgggtggag

20

&lt;210&gt; 64

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 64  
cccagactta gcgtctcagg

20

<210> 65  
<211> 20  
<212> DNA  
<213> Mouse

<400> 65  
agcagagacc tttggactcg

20

<210> 66  
<211> 20  
<212> DNA  
<213> Mouse

<400> 66  
gaaggctgag tgagtcccg

20

<210> 67  
<211> 20  
<212> DNA  
<213> Mouse

<400> 67  
ttgcacgagg agaaggttt

20

<210> 68  
<211> 20  
<212> DNA  
<213> Mouse

<400> 68  
gatgccaacg agacctgaat

20

<210> 69  
<211> 20  
<212> DNA  
<213> Mouse

<400> 69  
agaagccaaa accctcacct

20

<210> 70  
<211> 20  
<212> DNA  
<213> Mouse

<400> 70  
aaaaaaggccct gcaagaacct

20

<210> 71  
<211> 20  
<212> DNA  
<213> Mouse

<400> 71  
attcaggctc cgttggcata

20

<210> 72  
<211> 20  
<212> DNA  
<213> Mouse

<400> 72  
tgtccgcagt gtggaaacta

20

<210> 73  
<211> 20  
<212> DNA  
<213> Mouse

<400> 73  
atgtccaggg tagagagccc

20

<210> 74  
<211> 20  
<212> DNA  
<213> Mouse

<400> 74  
ggagttctcc taccctggct

20

<210> 75  
<211> 20

<212> DNA

<213> Mouse

<400> 75

gaggctctga gcagtgtaaa

20

<210> 76

<211> 14

<212> DNA

<213> Mouse

<400> 76

gcgatgttgt tgcg

14

<210> 77

<211> 18

<212> DNA

<213> Mouse

<400> 77

cagtgtcttt ccacattt

18

<210> 78

<211> 27

<212> DNA

<213> Mouse

<400> 78

aggcatattt tataataat ttgttgt

27

<210> 79

<211> 19

<212> DNA

<213> Mouse

<400> 79

ccggatgact ctacttgac

19

<210> 80

<211> 20

<212> DNA

<213> Mouse

<400> 80  
gctgtttatg gggtcgagaa

20

<210> 81  
<211> 20  
<212> DNA  
<213> Mouse

<400> 81  
aatttctgaa gcagggggat

20

<210> 82  
<211> 20  
<212> DNA  
<213> Mouse

<400> 82  
tccccctgct tcagaaatta

20

<210> 83  
<211> 20  
<212> DNA  
<213> Mouse

<400> 83  
agggggatga ttgtgagtga

20

<210> 84  
<211> 27  
<212> DNA  
<213> Mouse

<400> 84  
cttcttaat caatctctgt ctctgtg

27

<210> 85  
<211> 20  
<212> DNA  
<213> Mouse

<400> 85  
gggcacatat gaacctcctg

20

<210> 86  
<211> 20  
<212> DNA  
<213> Mouse

<400> 86  
ccaaaactctt agcttcttca

20

<210> 87  
<211> 21  
<212> DNA  
<213> Mouse

<400> 87  
acacagaaga cactgaagaa c

21

<210> 88  
<211> 22  
<212> DNA  
<213> Mouse

<400> 88  
cagttgttag aagcaggatc cc

22

<210> 89  
<211> 23  
<212> DNA  
<213> Mouse

<400> 89  
aggtgtcatat acctgggata ctc

23

<210> 90  
<211> 21  
<212> DNA  
<213> Mouse

<400> 90  
agagtttgggt ctcttccccct g

21

<210> 91  
<211> 23

<212> DNA

<213> Mouse

<400> 91

tatccaaacac atttatgtct gcg

23

<210> 92

<211> 20

<212> DNA

<213> Mouse

<400> 92

gccagtggtgc tgaaaagactg

20

<210> 93

<211> 20

<212> DNA

<213> Mouse

<400> 93

agggacacctgg agacatccctt

20

<210> 94

<211> 23

<212> DNA

<213> Mouse

<400> 94

ctgtaggctg ctttatctt ttg

23

<210> 95

<211> 20

<212> DNA

<213> Mouse

<400> 95

tgcgccttca gcacatgcga

20

<210> 96

<211> 23

<212> DNA

<213> Mouse

<400> 96  
tgca  
gtgtgtga catgtgcata gat

23

<210> 97  
<211> 21  
<212> DNA  
<213> Mouse

<400> 97  
gaaa  
aggccag gctacgcaga a

21

<210> 98  
<211> 23  
<212> DNA  
<213> Mouse

<400> 98  
ctgtaggctg ctttatctt ttg

23

<210> 99  
<211> 20  
<212> DNA  
<213> Mouse

<400> 99  
tgccc  
cttca gcacatgcca

20

<210> 100  
<211> 22  
<212> DNA  
<213> Mouse

<400> 100  
tagt  
gtggtt cctgactaac ct

22

<210> 101  
<211> 22  
<212> DNA  
<213> Mouse

<400> 101  
cggt  
ctacat agtgatgtat tc

22

<210> 102  
<211> 22  
<212> DNA  
<213> Mouse

<400> 102  
aaaagcatcc tgcatcccttc tg

22

<210> 103  
<211> 22  
<212> DNA  
<213> Mouse

<400> 103  
gggttataca gagaaacccct gt

22

<210> 104  
<211> 20  
<212> DNA  
<213> Mouse

<400> 104  
ttccaagctc acacatcagc

20

<210> 105  
<211> 20  
<212> DNA  
<213> Mouse

<400> 105  
tgctgtct gcattgagtg

20

<210> 106  
<211> 20  
<212> DNA  
<213> Mouse

<400> 106  
gacagtgtgg gagaatccgt

20

<210> 107  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 107

cccaaggcat aggtcacaat

20

&lt;210&gt; 108

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 108

attgtgacct atgccttggg

20

&lt;210&gt; 109

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 109

cgaaggaccg tcatctgagt

20

&lt;210&gt; 110

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 110

ggctttgatg tgaaaaaggc

20

&lt;210&gt; 111

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 111

agctcctcat cgctcatgtt

20

&lt;210&gt; 112

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 112  
tggaacatct ctgtcggaaag

20

<210> 113  
<211> 20  
<212> DNA  
<213> Mouse

<400> 113  
ggctctcatt gccaccctta

20

<210> 114  
<211> 20  
<212> DNA  
<213> Mouse

<400> 114  
ccagagaaca ggagacacctgc

20

<210> 115  
<211> 20  
<212> DNA  
<213> Mouse

<400> 115  
gtgctggata cactggcaga

20

<210> 116  
<211> 20  
<212> DNA  
<213> Mouse

<400> 116  
gcgagacgag tggtagttc

20

<210> 117  
<211> 20  
<212> DNA  
<213> Mouse

<400> 117  
acactgaaac ctcgcttgct

20

<210> 118  
<211> 20  
<212> DNA  
<213> Mouse

<400> 118  
agcaagcgag gtttcagtgt

20

<210> 119  
<211> 20  
<212> DNA  
<213> Mouse

<400> 119  
acggggcttg atccttttat

20

<210> 120  
<211> 25  
<212> DNA  
<213> Mouse

<400> 120  
aagttcatgg gcctcaccac ctgtc

25

<210> 121  
<211> 22  
<212> DNA  
<213> Mouse

<400> 121  
tactagctac cttcacata cc

22

<210> 122  
<211> 21  
<212> DNA  
<213> Mouse

<400> 122  
acctagccac tgtctcagtc t

21

<210> 123  
<211> 21

<212> DNA

<213> Mouse

<400> 123

acagaaggcag catttacaca g

21

<210> 124

<211> 20

<212> DNA

<213> Mouse

<400> 124

tggcacagct tcctcaagat

20

<210> 125

<211> 20

<212> DNA

<213> Mouse

<400> 125

aatgggaatt gtgctcttgg

20

<210> 126

<211> 20

<212> DNA

<213> Mouse

<400> 126

gggcatctgg caaagattta

20

<210> 127

<211> 20

<212> DNA

<213> Mouse

<400> 127

agataaacctg tgtgtccccgc

20

<210> 128

<211> 20

<212> DNA

<213> Mouse

<400> 128  
gatgtccgag aagggatgtg

20

<210> 129  
<211> 20  
<212> DNA  
<213> Mouse

<400> 129  
tgtcagctt gagtgcattc

20

<210> 130  
<211> 20  
<212> DNA  
<213> Mouse

<400> 130  
acatgcaggc tgtttgacct

20

<210> 131  
<211> 20  
<212> DNA  
<213> Mouse

<400> 131  
tgtcagctt gagtgcattc

20

<210> 132  
<211> 20  
<212> DNA  
<213> Mouse

<400> 132  
gtgctctgca gacaaaccaa

20

<210> 133  
<211> 20  
<212> DNA  
<213> Mouse

<400> 133  
gagccatttt gacccttaaa

20

<210> 134  
<211> 20  
<212> DNA  
<213> Mouse

<400> 134  
tttcagggtc aaaaatggctc

20

<210> 135  
<211> 17  
<212> DNA  
<213> Mouse

<400> 135  
tcgacagcaa ctgtgcg

17

<210> 136  
<211> 20  
<212> DNA  
<213> Mouse

<400> 136  
ggtgagatgt gggagatgaa

20

<210> 137  
<211> 20  
<212> DNA  
<213> Mouse

<400> 137  
cccggtgag tttaagaacc

20

<210> 138  
<211> 20  
<212> DNA  
<213> Mouse

<400> 138  
ggtgagatgt gggagatgaa

20

<210> 139  
<211> 20

<212> DNA  
<213> Mouse

<400> 139  
aggtaggcc caatttctcg

20

<210> 140  
<211> 20  
<212> DNA  
<213> Mouse

<400> 140  
ccagggttgc tgtactgaga

20

<210> 141  
<211> 20  
<212> DNA  
<213> Mouse

<400> 141  
caggttaggc ccaatttct

20

<210> 142  
<211> 20  
<212> DNA  
<213> Mouse

<400> 142  
ggtcagagtc cttccttccc

20

<210> 143  
<211> 20  
<212> DNA  
<213> Mouse

<400> 143  
tccaacttca cagggaaaccc

20

<210> 144  
<211> 20  
<212> DNA  
<213> Mouse

&lt;400&gt; 144

tttcctgtga agttggaggg

20

&lt;210&gt; 145

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 145

cacccatatg gcaaacatca

20

&lt;210&gt; 146

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 146

ggtcagagtc ctteccccc

20

&lt;210&gt; 147

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 147

tccaaacttca cagggaaaccc

20

&lt;210&gt; 148

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 148

tgatgttgc catatgggtg

20

&lt;210&gt; 149

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 149

gcttgctgct tccgatatgt

20

<210> 150  
<211> 19  
<212> DNA  
<213> Mouse

<400> 150  
gaaaaaggga gtcgccata

19

<210> 151  
<211> 20  
<212> DNA  
<213> Mouse

<400> 151  
gagccgccta actctcacac

20

<210> 152  
<211> 19  
<212> DNA  
<213> Mouse

<400> 152  
aggggataac ctgcataagg

19

<210> 153  
<211> 20  
<212> DNA  
<213> Mouse

<400> 153  
acaaaaattgc tcatttgccc

20

<210> 154  
<211> 20  
<212> DNA  
<213> Mouse

<400> 154  
ccatccccac tagccagata

20

<210> 155  
<211> 20

<212> DNA  
<213> Mouse

<400> 155  
gtcccccttg tcacagcaag

20

<210> 156  
<211> 20  
<212> DNA  
<213> Mouse

<400> 156  
tgagcacagg atagctccac

20

<210> 157  
<211> 20  
<212> DNA  
<213> Mouse

<400> 157  
aaaagaacac ctgtttgggg

20

<210> 158  
<211> 19  
<212> DNA  
<213> Mouse

<400> 158  
taaacacctgg ctgtgtgag

19

<210> 159  
<211> 20  
<212> DNA  
<213> Mouse

<400> 159  
ccctcagtga cttccctgtga

20

<210> 160  
<211> 20  
<212> DNA  
<213> Mouse

<400> 160  
caaaaccaca tggttaccca

20

<210> 161  
<211> 20  
<212> DNA  
<213> Mouse

<400> 161  
gccctattgc caaatgactt

20

<210> 162  
<211> 20  
<212> DNA  
<213> Mouse

<400> 162  
ggcagaaagg aatcagaagg

20

<210> 163  
<211> 20  
<212> DNA  
<213> Mouse

<400> 163  
cacattagcc attgtcctgg

20

<210> 164  
<211> 20  
<212> DNA  
<213> Mouse

<400> 164  
tccttatgt ccaacagccca

20

<210> 165  
<211> 20  
<212> DNA  
<213> Mouse

<400> 165  
catggtctgt gatgtgacca

20

<210> 166  
<211> 20  
<212> DNA  
<213> Mouse

<400> 166  
atacccttgg tgagagcagg

20

<210> 167  
<211> 20  
<212> DNA  
<213> Mouse

<400> 167  
gctgtcaaat gagaaaggca

20

<210> 168  
<211> 20  
<212> DNA  
<213> Mouse

<400> 168  
tatttcatgc tgggaccaaa

20

<210> 169  
<211> 20  
<212> DNA  
<213> Mouse

<400> 169  
agagaaaaac agtgggggtg

20

<210> 170  
<211> 20  
<212> DNA  
<213> Mouse

<400> 170  
cggttcctct cttcaccata

20

<210> 171  
<211> 20

<212> DNA

<213> Mouse

<400> 171

ctacatttcc ctgagctgcc

20

<210> 172

<211> 20

<212> DNA

<213> Mouse

<400> 172

gttgaccatg tcggtaaccc

20

<210> 173

<211> 20

<212> DNA

<213> Mouse

<400> 173

ccacacctacg gaaactgaat

20

<210> 174

<211> 20

<212> DNA

<213> Mouse

<400> 174

ggtgtttggc tcacaaacct

20

<210> 175

<211> 20

<212> DNA

<213> Mouse

<400> 175

gatgcacaca caaaaatccg

20

<210> 176

<211> 20

<212> DNA

<213> Mouse

<400> 176  
atcacccacc agaacgaaaa

20

<210> 177  
<211> 20  
<212> DNA  
<213> Mouse

<400> 177  
accctccagg agtaggtgct

20

<210> 178  
<211> 20  
<212> DNA  
<213> Mouse

<400> 178  
gatgagacag tgggcaaggt

20

<210> 179  
<211> 20  
<212> DNA  
<213> Mouse

<400> 179  
ttgtcaatag caccaagcca

20

<210> 180  
<211> 20  
<212> DNA  
<213> Mouse

<400> 180  
gccttaatag ccccccttgtt

20

<210> 181  
<211> 20  
<212> DNA  
<213> Mouse

<400> 181  
gcactcagca ttgcacagat

20

<210> 182  
<211> 20  
<212> DNA  
<213> Mouse

<400> 182  
ggacggacaa ttctggaaaa

20

<210> 183  
<211> 20  
<212> DNA  
<213> Mouse

<400> 183  
ctatcacacc tccgatgcct

20

<210> 184  
<211> 20  
<212> DNA  
<213> Mouse

<400> 184  
caagctggta gaatcccaa

20

<210> 185  
<211> 20  
<212> DNA  
<213> Mouse

<400> 185  
tctttggaga agcagaccgt

20

<210> 186  
<211> 20  
<212> DNA  
<213> Mouse

<400> 186  
tacagcatat gcatgccagg

20

<210> 187  
<211> 20

<212> DNA

<213> Mouse

<400> 187

atccctcagg gcattacacg

20

<210> 188

<211> 20

<212> DNA

<213> Mouse

<400> 188

gcaatctctt gtgtccaggc

20

<210> 189

<211> 20

<212> DNA

<213> Mouse

<400> 189

atccctcagg gcattacacg

20

<210> 190

<211> 20

<212> DNA

<213> Mouse

<400> 190

tacagcatat gcatgccagg

20

<210> 191

<211> 20

<212> DNA

<213> Mouse

<400> 191

ggcctggaca caagagattg

20

<210> 192

<211> 20

<212> DNA

<213> Mouse

<400> 192  
aagtgggtgg acagtgaagg

20

<210> 193  
<211> 20  
<212> DNA  
<213> Mouse

<400> 193  
cagcttcctc catcttctgg

20

<210> 194  
<211> 20  
<212> DNA  
<213> Mouse

<400> 194  
agagcctcca gtagatggca

20

<210> 195  
<211> 20  
<212> DNA  
<213> Mouse

<400> 195  
tcgtggacaa gtccttctt

20

<210> 196  
<211> 20  
<212> DNA  
<213> Mouse

<400> 196  
catcgagtagt gtcaatggcg

20

<210> 197  
<211> 20  
<212> DNA  
<213> Mouse

<400> 197  
ttgtccagtt ttagtcccg

20

<210> 198  
<211> 20  
<212> DNA  
<213> Mouse

<400> 198  
cagactgggt tttccgacat

20

<210> 199  
<211> 20  
<212> DNA  
<213> Mouse

<400> 199  
gtcaaagttg tccaggccat

20

<210> 200  
<211> 18  
<212> DNA  
<213> Mouse

<400> 200  
aggacggacc ccaagatg

18

<210> 201  
<211> 20  
<212> DNA  
<213> Mouse

<400> 201  
tgtctcgac ttcctcacag

20

<210> 202  
<211> 20  
<212> DNA  
<213> Mouse

<400> 202  
ccagaagatg gaggaagctg

20

<210> 203  
<211> 20

<212> DNA  
<213> Mouse

<400> 203  
tctactggag gctcttggga

20

<210> 204  
<211> 20  
<212> DNA  
<213> Mouse

<400> 204  
gaaaaacgac cagatttacg

20

<210> 205  
<211> 20  
<212> DNA  
<213> Mouse

<400> 205  
gatctcagca gcatagaacc

20

<210> 206  
<211> 20  
<212> DNA  
<213> Mouse

<400> 206  
acacattaag ctgacggact

20

<210> 207  
<211> 20  
<212> DNA  
<213> Mouse

<400> 207  
caaacataag gacacccagt

20

<210> 208  
<211> 20  
<212> DNA  
<213> Mouse

<400> 208  
actgggtgtc cttatgtttg

20

<210> 209  
<211> 20  
<212> DNA  
<213> Mouse

<400> 209  
cctctctttg ggatccttat

20

<210> 210  
<211> 20  
<212> DNA  
<213> Mouse

<400> 210  
gtcataaaaga ggatcgacca

20

<210> 211  
<211> 20  
<212> DNA  
<213> Mouse

<400> 211  
gctctgtcta gaagtgcctg

20

<210> 212  
<211> 18  
<212> DNA  
<213> Mouse

<400> 212  
accaagacccg aagagggg

18

<210> 213  
<211> 22  
<212> DNA  
<213> Mouse

<400> 213  
ggcattacac gctaactttt cc

22

<210> 214  
<211> 20  
<212> DNA  
<213> Mouse

<400> 214  
agtgccacca acctggtaag

20

<210> 215  
<211> 18  
<212> DNA  
<213> Mouse

<400> 215  
aagtgcctgc agggatgc

18

<210> 216  
<211> 20  
<212> DNA  
<213> Mouse

<400> 216  
tgctttggtg agcaatgttt

20

<210> 217  
<211> 20  
<212> DNA  
<213> Mouse

<400> 217  
agggacaccc ttaccaggtt

20

<210> 218  
<211> 20  
<212> DNA  
<213> Mouse

<400> 218  
ctgatgctt ggtgagcaat

20

<210> 219  
<211> 19

<212> DNA  
<213> Mouse

<400> 219  
gggacaccct taccaggtt

19

<210> 220  
<211> 20  
<212> DNA  
<213> Mouse

<400> 220  
acaggacaaa tgctggggtt

20

<210> 221  
<211> 20  
<212> DNA  
<213> Mouse

<400> 221  
gtggtaaaga acgcttggct

20

<210> 222  
<211> 24  
<212> DNA  
<213> Mouse

<400> 222  
ggtatctcac ttggtaggaa cctc

24

<210> 223  
<211> 17  
<212> DNA  
<213> Mouse

<400> 223  
aagaacgctt ggctggc

17

<210> 224  
<211> 20  
<212> DNA  
<213> Mouse

&lt;400&gt; 224

gcccgcattcg gtgatgtact

20

&lt;210&gt; 225

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 225

acaatggctc aaaaccgttc

20

&lt;210&gt; 226

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 226

gccttggaa tttaccacct

20

&lt;210&gt; 227

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 227

agtacatcac caggatcgcc

20

&lt;210&gt; 228

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 228

taaaaggcca tgcgataaagc

20

&lt;210&gt; 229

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 229

agagctctgt ggggttctca

20

<210> 230

<211> 20

<212> DNA

<213> Mouse

<400> 230

gaaggggaca gtgttggaga

20

<210> 231

<211> 20

<212> DNA

<213> Mouse

<400> 231

tccatcaagg aaggatccac

20

<210> 232

<211> 19

<212> DNA

<213> Mouse

<400> 232

ggtgggtaat gattggact

19

<210> 233

<211> 19

<212> DNA

<213> Mouse

<400> 233

tgacgtggag ggaactgcc

19

<210> 234

<211> 20

<212> DNA

<213> Mouse

<400> 234

tgagatctgg tgccctctct

20

<210> 235

<211> 20

<212> DNA  
<213> Mouse

<400> 235  
gcctgatcta ggctggaaaa

20

<210> 236  
<211> 20  
<212> DNA  
<213> Mouse

<400> 236  
aggcagaaag cagacaagga

20

<210> 237  
<211> 20  
<212> DNA  
<213> Mouse

<400> 237  
cgacagcact tgtgaccact

20

<210> 238  
<211> 20  
<212> DNA  
<213> Mouse

<400> 238  
ctgcagatgt agaccaggca

20

<210> 239  
<211> 20  
<212> DNA  
<213> Mouse

<400> 239  
ctgtggtgga ttggacagtg

20

<210> 240  
<211> 20  
<212> DNA  
<213> Mouse

<400> 240  
ttgcctaaca ctcccaaacc

20

<210> 241  
<211> 20  
<212> DNA  
<213> Mouse

<400> 241  
tatttaggc accaccaggc

20

<210> 242  
<211> 20  
<212> DNA  
<213> Mouse

<400> 242  
acctgtcttg tgggtggaag

20

<210> 243  
<211> 20  
<212> DNA  
<213> Mouse

<400> 243  
ctgtggtggc ttggacagtgc

20

<210> 244  
<211> 20  
<212> DNA  
<213> Mouse

<400> 244  
gtggcttggc gctattgaca

20

<210> 245  
<211> 20  
<212> DNA  
<213> Mouse

<400> 245  
ggggcttata agggcatttt

20

<210> 246

<211> 21

<212> DNA

<213> Mouse

<400> 246

caattgagga atggctacca a

21

<210> 247

<211> 20

<212> DNA

<213> Mouse

<400> 247

tggcttcatg tccattgtgt

20

<210> 248

<211> 22

<212> DNA

<213> Mouse

<400> 248

cagaaccaca aaggtaaatt gc

22

<210> 249

<211> 21

<212> DNA

<213> Mouse

<400> 249

tcatgtttgc tgtccagttt g

21

<210> 250

<211> 29

<212> DNA

<213> Homo sapiens

<400> 250

gccaccatgc tgggcctgc tgtcctggg

29

<210> 251

<211> 24

<212> DNA

<213> Homo sapiens

<400> 251

tcactcatgt ttccccctgat ttcc

24

<210> 252

<211> 20

<212> DNA

<213> Homo sapiens

<400> 252

ctgatttcct gtgttcccgt

20

<210> 253

<211> 20

<212> DNA

<213> Homo sapiens

<400> 253

catgctggcc tacttcatca

20

<210> 254

<211> 29

<212> DNA

<213> Homo sapiens

<400> 254

gccttgagg tcagctacgg tgcttagcat

29

<210> 255

<211> 24

<212> DNA

<213> Homo sapiens

<400> 255

tcactcatgt ttccccctgat ttcc

24

<210> 256

<211> 20

<212> DNA

<213> Homo sapiens

<400> 256  
aggaaggcaga gaaaggccag

20

<210> 257  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 257  
tcagaactgc ctctgagctg

20

<210> 258  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 258  
tcttcacgtt ctgggggaac

20

<210> 259  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 259  
actacagcat cagcagcagg

20

<210> 260  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 260  
aagctgaaga acttcccggt

20

<210> 261  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 261  
tgggctacga cctctttgtat

20

<210> 262  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 262  
atcttcaggc gctctgtcct

20

<210> 263  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 263  
gtacgacctg aagctgtggg

20

<210> 264  
<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 264  
atcttcaggc gctctgtcc

19

<210> 265  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 265  
gtacgacctg aagctgtggg

20

<210> 266  
<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 266  
atcttcaggc gctctgtcc

19

<210> 267  
<211> 21

<212> DNA

<213> Homo sapiens

<400> 267

gagtacgacc tgaagctgtg g

21

<210> 268

<211> 19

<212> DNA

<213> Homo sapiens

<400> 268

atcttcaggc gctctgtcc

19

<210> 269

<211> 19

<212> DNA

<213> Homo sapiens

<400> 269

tacgacactga agctgtggg

19

<210> 270

<211> 19

<212> DNA

<213> Homo sapiens

<400> 270

atcttcaggc gctctgtcc

19

<210> 271

<211> 19

<212> DNA

<213> Homo sapiens

<400> 271

tacgacactga agctgtggg

19

<210> 272

<211> 18

<212> DNA

<213> Homo sapiens

&lt;400&gt; 272

gctgtccccga tggtaaac

18

&lt;210&gt; 273

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 273

accttttgcg gccaggatg

19

&lt;210&gt; 274

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 274

gctgtccccga tggtaaac

18

&lt;210&gt; 275

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 275

cacctttgtt ggccaggat

19

&lt;210&gt; 276

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 276

gctgtccccga tggtaaac

18

&lt;210&gt; 277

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 277

ccttttgcgg ccaggatg

18

<210> 278  
<211> 18  
<212> DNA  
<213> Homo sapiens

<400> 278  
cctgaaccag tgggctgt

18

<210> 279  
<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 279  
accttttgta gccaggatg

19

<210> 280  
<211> 18  
<212> DNA  
<213> Homo sapiens

<400> 280  
cctgaaccag tgggctgt

18

<210> 281  
<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 281  
cacctttgtt ggccaggat

19

<210> 282  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 282  
tcatgttcc cctgatttcc

20

<210> 283  
<211> 20

<212> DNA

<213> Homo sapiens

<400> 283

catgctggcc tacttcatca

20

<210> 284

<211> 20

<212> DNA

<213> Homo sapiens

<400> 284

atgagcaggt aacacacctggg

20

<210> 285

<211> 20

<212> DNA

<213> Homo sapiens

<400> 285

tcatcacctg ggtctccctt

20

<210> 286

<211> 20

<212> DNA

<213> Homo sapiens

<400> 286

atgagcaggt aacacacctggg

20

<210> 287

<211> 20

<212> DNA

<213> Homo sapiens

<400> 287

ttcatcacct gggctccctt

20

<210> 288

<211> 20

<212> DNA

<213> Mouse

<400> 288  
tgggttgtt tctctggttt

20

<210> 289  
<211> 21  
<212> DNA  
<213> Mouse

<400> 289  
ccttttaca gtctgccagg t

21

<210> 290  
<211> 20  
<212> DNA  
<213> Mouse

<400> 290  
tgggttgtt tctctggttt

20

<210> 291  
<211> 21  
<212> DNA  
<213> Mouse

<400> 291  
gatccccttt ttacagtctg c

21

<210> 292  
<211> 20  
<212> DNA  
<213> Mouse

<400> 292  
acggggttgg tactgtgtgt

20

<210> 293  
<211> 20  
<212> DNA  
<213> Mouse

<400> 293  
cacccattgt tagtgctgga

20

<210> 294  
<211> 20  
<212> DNA  
<213> Mouse

<400> 294  
acggggttgg tactgtgtgt

20

<210> 295  
<211> 20  
<212> DNA  
<213> Mouse

<400> 295  
cacacaccca cccattgtta

20

<210> 296  
<211> 20  
<212> DNA  
<213> Mouse

<400> 296  
tgcattggcc agactagaaa

20

<210> 297  
<211> 19  
<212> DNA  
<213> Mouse

<400> 297  
cggtctggct atgacctat

19

<210> 298  
<211> 20  
<212> DNA  
<213> Mouse

<400> 298  
tgcattggcc agactagaaa

20

<210> 299  
<211> 20

<212> DNA

<213> Mouse

<400> 299

cggctgggct atgacctatt

20

<210> 300

<211> 20

<212> DNA

<213> Mouse

<400> 300

gttctgcagc atgatgtcgt

20

<210> 301

<211> 20

<212> DNA

<213> Mouse

<400> 301

ggcagttgtg actctgttgc

20

<210> 302

<211> 20

<212> DNA

<213> Mouse

<400> 302

gttctgcagc atgatgtcgt

20

<210> 303

<211> 20

<212> DNA

<213> Mouse

<400> 303

ctgcaggcag ttgtgactct

20

<210> 304

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 304

ccatcctttt tgccctgtctt

20

&lt;210&gt; 305

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 305

tctggaggaa catgtgatgg

20

&lt;210&gt; 306

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 306

caccatcctt tttgcctgtc

20

&lt;210&gt; 307

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 307

gaacatgtga tggggcaac

19

&lt;210&gt; 308

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 308

caaaggcagca ggaggagtg

19

&lt;210&gt; 309

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 309

aaatgtactg gccaggcaac

20

<210> 310  
<211> 20  
<212> DNA  
<213> Mouse

<400> 310  
agtgctagac ccāgcaccag

20

<210> 311  
<211> 20  
<212> DNA  
<213> Mouse

<400> 311  
aaatgtactg gccaggcaac

20

<210> 312  
<211> 20  
<212> DNA  
<213> Mouse

<400> 312  
gcactgacca gtctgtcacc

20

<210> 313  
<211> 20  
<212> DNA  
<213> Mouse

<400> 313  
gtccccagag aaaagcacag

20

<210> 314  
<211> 20  
<212> DNA  
<213> Mouse

<400> 314  
cagtctgtca ccacacctgg

20

<210> 315  
<211> 20

<212> DNA

<213> Mouse

<400> 315

cagtggtccc cagagaaaag

20

<210> 316

<211> 20

<212> DNA

<213> Mouse

<400> 316

tactattcgg ggcttgg

20

<210> 317

<211> 20

<212> DNA

<213> Mouse

<400> 317

gcagcactat gtgcctggta

20

<210> 318

<211> 20

<212> DNA

<213> Mouse

<400> 318

tactattcgg ggcttgg

20

<210> 319

<211> 20

<212> DNA

<213> Mouse

<400> 319

gcctggatt tgatcgctt

20

<210> 320

<211> 20

<212> DNA

<213> Mouse

<400> 320  
gctcagctag ggatggagaa

20

<210> 321  
<211> 20  
<212> DNA  
<213> Mouse

<400> 321  
cagctcaggg acacaatgaa

20

<210> 322  
<211> 20  
<212> DNA  
<213> Mouse

<400> 322  
tcctacaggg tagggctcag

20

<210> 323  
<211> 20  
<212> DNA  
<213> Mouse

<400> 323  
cagctcaggg acacaatgaa

20

<210> 324  
<211> 20  
<212> DNA  
<213> Mouse

<400> 324  
gggactgatg tgtggcttgt

20

<210> 325  
<211> 20  
<212> DNA  
<213> Mouse

<400> 325  
aggcgtccca ggaatagaag

20

<210> 326  
<211> 21  
<212> DNA  
<213> Mouse

<400> 326  
ggactgatgt gtggcttgtt t

21

<210> 327  
<211> 20  
<212> DNA  
<213> Mouse

<400> 327  
aggcgtccca ggaatagaag

20

<210> 328  
<211> 20  
<212> DNA  
<213> Mouse

<400> 328  
tgtttctgtt ctgggtggctg

20

<210> 329  
<211> 20  
<212> DNA  
<213> Mouse

<400> 329  
atctgcaggc aggatcagac

20

<210> 330  
<211> 20  
<212> DNA  
<213> Mouse

<400> 330  
ctcagtggtg ggtgacagtg

20

<210> 331  
<211> 20

<212> DNA

<213> Mouse

<400> 331

atctgcaggc aggatcagac

20

<210> 332

<211> 20

<212> DNA

<213> Mouse

<400> 332

acacacagta ccaaccccggt

20

<210> 333

<211> 20

<212> DNA

<213> Mouse

<400> 333

cctgtggta tcaagaagca

20

<210> 334

<211> 20

<212> DNA

<213> Mouse

<400> 334

tgcttcgttga tcaccacagg

20

<210> 335

<211> 20

<212> DNA

<213> Mouse

<400> 335

gcaacagagt cacaactgcc

20

<210> 336

<211> 20

<212> DNA

<213> Mouse

<400> 336  
acacacagta ccaacccgt

20

<210> 337  
<211> 20  
<212> DNA  
<213> Mouse

<400> 337  
gcaacagagt cacaactgcc

20

<210> 338  
<211> 20  
<212> DNA  
<213> Mouse

<400> 338  
gggtttatgt ggcaagcact

20

<210> 339  
<211> 20  
<212> DNA  
<213> Mouse

<400> 339  
actccatttg ccttttgtgg

20

<210> 340  
<211> 20  
<212> DNA  
<213> Mouse

<400> 340  
cgctacttcg cttttatccg

20

<210> 341  
<211> 20  
<212> DNA  
<213> Mouse

<400> 341  
atgatgacgt acgacgacga

20

<210> 342  
<211> 21  
<212> DNA  
<213> Mouse

<400> 342  
gaaaacaatc ggggagaagt. c

21

<210> 343  
<211> 20  
<212> DNA  
<213> Mouse

<400> 343  
tcaaattatc acacgccagg.

20

<210> 344  
<211> 20  
<212> DNA  
<213> Mouse

<400> 344  
agtgagaggc ccagtctcaa

20

<210> 345  
<211> 20  
<212> DNA  
<213> Mouse

<400> 345  
gatctgatgc cctcttctgc

20

<210> 346  
<211> 20  
<212> DNA  
<213> Mouse

<400> 346  
gctagccttg aagccaaacac

20

<210> 347  
<211> 20

<212> DNA

<213> Mouse

<400> 347

tgaacagcat gcttacccag

20

<210> 348

<211> 20

<212> DNA

<213> Mouse

<400> 348

tcccttagagg cctgtctgtc

20

<210> 349

<211> 20

<212> DNA

<213> Mouse

<400> 349

tctgtctcgga gcctcttcta

20

<210> 350

<211> 20

<212> DNA

<213> Mouse

<400> 350

gatagtccct tagccagccc

20

<210> 351

<211> 20

<212> DNA

<213> Mouse

<400> 351

gccccatagctc ctcactgctc

20

<210> 352

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 352

cagagtgggc tctggtcttc

20

&lt;210&gt; 353

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 353

ttgtgttcag atgctcctgc

20

&lt;210&gt; 354

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 354

ttatttctgt gctagccgccc

20

&lt;210&gt; 355

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 355

atcaagtcaa cgtccccaag

20

&lt;210&gt; 356

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 356

acctggcctg tgctaatctc

20

&lt;210&gt; 357

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 357

gcaccaaccc taagaaagca

20

<210> 358  
<211> 22  
<212> DNA  
<213> Mouse

<400> 358  
tcaggctaac ctcaaactca ca

22

<210> 359  
<211> 27  
<212> DNA  
<213> Mouse

<400> 359  
aaagaaaaaga aaagaaaaag tcagaca

27

<210> 360  
<211> 20  
<212> DNA  
<213> Mouse

<400> 360  
cccagaactc catcctcaaa

20

<210> 361  
<211> 20  
<212> DNA  
<213> Mouse

<400> 361  
cccaacacctg ggtcagctat

20

<210> 362  
<211> 20  
<212> DNA  
<213> Mouse

<400> 362  
ggggcaggtg ggtaataagt

20

<210> 363  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 363

caaaagccca actccttgag

20

&lt;210&gt; 364

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 364

gctcagtggg taagagcacc

20

&lt;210&gt; 365

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 365

ctaccctgccc gctaattctca

20

&lt;210&gt; 366

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 366

cagtttagcac cccaccctaa

20

&lt;210&gt; 367

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 367

tctgcacccctc tgttcacctg

20

&lt;210&gt; 368

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 368  
acctctaggg tttacgggga

20

<210> 369  
<211> 20  
<212> DNA  
<213> Mouse

<400> 369  
cctcaggtag tgcaagctcc

20

<210> 370  
<211> 20  
<212> DNA  
<213> Mouse

<400> 370  
tcagttacca agggttcgg

20

<210> 371  
<211> 20  
<212> DNA  
<213> Mouse

<400> 371  
atagggtgtc acaggccagg

20

<210> 372  
<211> 20  
<212> DNA  
<213> Mouse

<400> 372  
tcagttacca agggttcgg

20

<210> 373  
<211> 20  
<212> DNA  
<213> Mouse

<400> 373  
atagggtgtc acaggccagg

20

<210> 374  
<211> 20  
<212> DNA  
<213> Mouse

<400> 374  
gtggttgctg ggatttgaac

20

<210> 375  
<211> 20  
<212> DNA  
<213> Mouse

<400> 375  
caagcaacca aacaacccaa

20

<210> 376  
<211> 20  
<212> DNA  
<213> Mouse

<400> 376  
tccggaggac cataaatctg

20

<210> 377  
<211> 20  
<212> DNA  
<213> Mouse

<400> 377  
cacagtccca gtcattccct

20

<210> 378  
<211> 20  
<212> DNA  
<213> Mouse

<400> 378  
gtccccaaag ctagcacagg

20

<210> 379  
<211> 20

<212> DNA

<213> Mouse

<400> 379

tcatgagcca ccatgtgatt

20

<210> 380

<211> 20

<212> DNA

<213> Mouse

<400> 380

gaccttcgga agagcagttg

20

<210> 381

<211> 20

<212> DNA

<213> Mouse

<400> 381

agtgtgtgtc gccatatcca

20

<210> 382

<211> 20

<212> DNA

<213> Mouse

<400> 382

cctactctct ctcccccgtt

20

<210> 383

<211> 20

<212> DNA

<213> Mouse

<400> 383

ggaaaaatgtt tggccttgaa

20

<210> 384

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 384

ctggagtgaa aggccaggaaag

20

&lt;210&gt; 385

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 385

aggccggcacc atatgaataa

20

&lt;210&gt; 386

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 386

tgagagtgaaa aattctgttc a

21

&lt;210&gt; 387

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 387

ggatgtaaaa ggtggcaagg

20

&lt;210&gt; 388

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 388

ctgttggagg aggtggccta

20

&lt;210&gt; 389

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 389

tgcttgatg ttttcctcg t

21

<210> 390  
<211> 20  
<212> DNA  
<213> Mouse

<400> 390  
tgagagtgcc ctcctcttg

20

<210> 391  
<211> 18  
<212> DNA  
<213> Mouse

<400> 391  
gaaccctga ccccagac

18

<210> 392  
<211> 22  
<212> DNA  
<213> Mouse

<400> 392  
tgaagtgcag attttacat gg

22

<210> 393  
<211> 20  
<212> DNA  
<213> Mouse

<400> 393  
gttttgggggt ggaaaaggat

20

<210> 394  
<211> 20  
<212> DNA  
<213> Mouse

<400> 394  
ccgtcgacat ttaggtgaca

20

<210> 395  
<211> 20

<212> DNA

<213> Mouse

<400> 395

gatactgggg tgggtggtaa

20

<210> 396

<211> 20

<212> DNA

<213> Mouse

<400> 396

ccgtcgacat ttaggtgaca

20

<210> 397

<211> 20

<212> DNA

<213> Mouse

<400> 397

cgtcccgagct gtgttaactga

20

<210> 398

<211> 21

<212> DNA

<213> Mouse

<400> 398

ggaaagcaaat gctccactaa a

21

<210> 399

<211> 20

<212> DNA

<213> Mouse

<400> 399

tatcccttagc cccttgcgtg

20

<210> 400

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 400

ccgtcgacat ttagtgaca

20

&lt;210&gt; 401

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 401

gggtcctgtt ggttagtgacc

20

&lt;210&gt; 402

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 402

tataaggcgc ccctcattgg

20

&lt;210&gt; 403

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 403

caggccagac actgcttaca

20

&lt;210&gt; 404

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 404

ccttggatc tggtgtgact

20

&lt;210&gt; 405

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 405

tgggtttaga gtacggctgg

20

<210> 406  
<211> 20  
<212> DNA  
<213> Mouse

<400> 406  
accatttcc taatccctg

20

<210> 407  
<211> 20  
<212> DNA  
<213> Mouse

<400> 407  
atctctccag cccctctcag

20

<210> 408  
<211> 20  
<212> DNA  
<213> Mouse

<400> 408  
gggctggaa ttgaacctat

20

<210> 409  
<211> 20  
<212> DNA  
<213> Mouse

<400> 409  
tgaatccctt acagccttgc

20

<210> 410  
<211> 20  
<212> DNA  
<213> Mouse

<400> 410  
gccccataaa atccactcct

20

<210> 411  
<211> 20

<212> DNA

<213> Mouse

<400> 411

gctccggaaag gctagaagat

20

<210> 412

<211> 20

<212> DNA

<213> Mouse

<400> 412

ggtttggag tgtaggcaa

20

<210> 413

<211> 20

<212> DNA

<213> Mouse

<400> 413

actcagttgg cctctccctca

20

<210> 414

<211> 19

<212> DNA

<213> Mouse

<400> 414

acagaaaatcc ctcatgcga

19

<210> 415

<211> 21

<212> DNA

<213> Mouse

<400> 415

tcagtgtgga ccagaaaatgc c

21

<210> 416

<211> 22

<212> DNA

<213> Mouse

<400> 416  
tctgcaagtc agctcttgat aa

22

<210> 417  
<211> 23  
<212> DNA  
<213> Mouse

<400> 417  
actcataagg gtcaagctgt ctg

23

<210> 418  
<211> 20  
<212> DNA  
<213> Mouse

<400> 418  
tctccccctt taccactccc

20

<210> 419  
<211> 20  
<212> DNA  
<213> Mouse

<400> 419  
gcaaggagtc aaaaacagca

20

<210> 420  
<211> 20  
<212> DNA  
<213> Mouse

<400> 420  
gctagttggg gaacaaaacca

20

<210> 421  
<211> 20  
<212> DNA  
<213> Mouse

<400> 421  
actgcaaatg tccaaactcca

20

<210> 422  
<211> 20  
<212> DNA  
<213> Mouse

<400> 422  
cagttacaca gctgggacga

20

<210> 423  
<211> 20  
<212> DNA  
<213> Mouse

<400> 423  
gcaagagcct agcaatccac

20

<210> 424  
<211> 20  
<212> DNA  
<213> Mouse

<400> 424  
cagtttagca ccccacccct

20

<210> 425  
<211> 20  
<212> DNA  
<213> Mouse

<400> 425  
tctgcacctc tgttcacctg

20

<210> 426  
<211> 20  
<212> DNA  
<213> Mouse

<400> 426  
gggttccact tgatgctgat

20

<210> 427  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 427

tggctgttt cctggagctt

20

&lt;210&gt; 428

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 428

tgttaggaaat gtttctgcac c

21

&lt;210&gt; 429

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 429

acatggaaaca ggattctggc

20

&lt;210&gt; 430

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 430

gcaggcaaacc agacagacaaa

20

&lt;210&gt; 431

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 431

atgggggatc ccttactgac

20

&lt;210&gt; 432

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 432  
cggtcaggag tagtgtgggt

20

<210> 433  
<211> 20  
<212> DNA  
<213> Mouse

<400> 433  
cagcagctga tattgaggca

20

<210> 434  
<211> 22  
<212> DNA  
<213> Mouse

<400> 434  
aatgatgaag tgtcagcctc ag

22

<210> 435  
<211> 20  
<212> DNA  
<213> Mouse

<400> 435  
caacagaact caaagcctgg

20

<210> 436  
<211> 20  
<212> DNA  
<213> Mouse

<400> 436  
agcaggcaca ggtcttttgt

20

<210> 437  
<211> 20  
<212> DNA  
<213> Mouse

<400> 437  
aagaacagga cagtgggtggg

20

<210> 438  
<211> 20  
<212> DNA  
<213> Mouse

<400> 438  
cagcgattgg ctcttccttt

20

<210> 439  
<211> 20  
<212> DNA  
<213> Mouse

<400> 439  
ggggcttcct ttctgaggtt

20

<210> 440  
<211> 20  
<212> DNA  
<213> Mouse

<400> 440  
agtcaggttc cagcttggta

20

<210> 441  
<211> 20  
<212> DNA  
<213> Mouse

<400> 441  
attttccctt cctgcttctc

20

<210> 442  
<211> 20  
<212> DNA  
<213> Mouse

<400> 442  
ccaaaggctct gctggttatc

20

<210> 443  
<211> 20

<212> DNA

<213> Mouse

<400> 443

tgagggtgga gaatggaaag

20

<210> 444

<211> 20

<212> DNA

<213> Mouse

<400> 444

gccccataaaa atccactcct

20

<210> 445

<211> 20

<212> DNA

<213> Mouse

<400> 445

ttgccttaaca ctcggaaacc

20

<210> 446

<211> 20

<212> DNA

<213> Mouse

<400> 446

cagttacaca gctgggacga

20

<210> 447

<211> 20

<212> DNA

<213> Mouse

<400> 447

gcaagagcct agcaatccac

20

<210> 448

<211> 20

<212> DNA

<213> Mouse

<400> 448  
cagcaccctc ctctggtctc

20

<210> 449  
<211> 20  
<212> DNA  
<213> Mouse

<400> 449  
tgtctccaga ggttctgcct

20

<210> 450  
<211> 24  
<212> DNA  
<213> Mouse

<400> 450  
tggtggtgta atactattcc tttg

24

<210> 451  
<211> 26  
<212> DNA  
<213> Mouse

<400> 451  
tccttaattt ttggctttt gataca

26

<210> 452  
<211> 20  
<212> DNA  
<213> Mouse

<400> 452  
cagctgtgtg catgttgacc

20

<210> 453  
<211> 20  
<212> DNA  
<213> Mouse

<400> 453  
catcatgaag actcaggga

20

<210> 454  
<211> 20  
<212> DNA  
<213> Mouse

<400> 454  
gtccacacacct ggctttgtt

20

<210> 455  
<211> 20  
<212> DNA  
<213> Mouse

<400> 455  
cagcactcag tgaggttcca

20

<210> 456  
<211> 20  
<212> DNA  
<213> Mouse

<400> 456  
atgtaatgga aaggctgctg

20

<210> 457  
<211> 20  
<212> DNA  
<213> Mouse

<400> 457  
cagcactcag tgaggttcca

20

<210> 458  
<211> 21  
<212> DNA  
<213> Mouse

<400> 458  
aaacaggcat gaaaactcagg a

21

<210> 459  
<211> 20

<212> DNA

<213> Mouse

<400> 459

gggtatcatt gtcacctcca

20

<210> 460

<211> 20

<212> DNA

<213> Mouse

<400> 460

cacaggccaa gttgttgtt

20

<210> 461

<211> 20

<212> DNA

<213> Mouse

<400> 461

caggggacct tctgaatgat

20

<210> 462

<211> 20

<212> DNA

<213> Mouse

<400> 462

agctcaggtc cagttggta

20

<210> 463

<211> 20

<212> DNA

<213> Mouse

<400> 463

accacaaaat tttccctcc

20

<210> 464

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 464

cgggacctaa aactggacaa

20

&lt;210&gt; 465

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 465

tggggacagt taccaggaag

20

&lt;210&gt; 466

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 466

ccggaggacc ataaaatctga

20

&lt;210&gt; 467

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 467

cctcaaaaac aagcctgagc

20

&lt;210&gt; 468

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 468

ccttcagaaa tgtgtttgga ca

22

&lt;210&gt; 469

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 469

tcctgagttc aaatcccagc

20

<210> 470  
<211> 20  
<212> DNA  
<213> Mouse

<400> 470  
cttccatc tccaccctca

20

<210> 471  
<211> 20  
<212> DNA  
<213> Mouse

<400> 471  
aggtcctagg gagaggtcca

20

<210> 472  
<211> 20  
<212> DNA  
<213> Mouse

<400> 472  
aggcctaccc aaggacatct

20

<210> 473  
<211> 20  
<212> DNA  
<213> Mouse

<400> 473  
gcagtgagct gcagagtttg

20

<210> 474  
<211> 20  
<212> DNA  
<213> Mouse

<400> 474  
agacacccta ggtcctgctg

20

<210> 475  
<211> 22

<212> DNA

<213> Mouse

<400> 475

tgatcttcc aaacgcataa ga

22

<210> 476

<211> 20

<212> DNA

<213> Mouse

<400> 476

gcaagcaacc tgaacatgaa

20

<210> 477

<211> 20

<212> DNA

<213> Mouse

<400> 477

gcttacgatg gtcgtgaggt

20

<210> 478

<211> 20

<212> DNA

<213> Mouse

<400> 478

acatgcctgc ctatcttgc

20

<210> 479

<211> 20

<212> DNA

<213> Mouse

<400> 479

ggaacctgtt ttccatggtg

20

<210> 480

<211> 20

<212> DNA

<213> Mouse

<400> 480  
accttggtcc tgggtgtgagc

20

<210> 481  
<211> 20  
<212> DNA  
<213> Mouse

<400> 481  
tagctgggac gtggtatggt

20

<210> 482  
<211> 20  
<212> DNA  
<213> Mouse

<400> 482  
ccatgggaga ccagaaggta

20

<210> 483  
<211> 20  
<212> DNA  
<213> Mouse

<400> 483  
tgagtgtcct ctgcctgatg

20

<210> 484  
<211> 20  
<212> DNA  
<213> Mouse

<400> 484  
gcgcgtacat cctcctatgt

20

<210> 485  
<211> 20  
<212> DNA  
<213> Mouse

<400> 485  
cccaactatgg tccccagagaa

20

<210> 486  
<211> 20  
<212> DNA  
<213> Mouse

<400> 486  
ttgcacgtct ttgtttcgag

20

<210> 487  
<211> 24  
<212> DNA  
<213> Mouse

<400> 487  
aaaggggaat agacctgagt agaa

24

<210> 488  
<211> 20  
<212> DNA  
<213> Mouse

<400> 488  
ccaagagtca gccttggagt

20

<210> 489  
<211> 20  
<212> DNA  
<213> Mouse

<400> 489  
ggacaggttag ctcacccaac

20

<210> 490  
<211> 19  
<212> DNA  
<213> Mouse

<400> 490  
tgccagctt ggctatcat

19

<210> 491  
<211> 20

<212> DNA

<213> Mouse

<400> 491

ttcattgtgt ccctgagctg

20

<210> 492

<211> 24

<212> DNA

<213> Mouse

<400> 492

agctttggct atcatgggtc tcag

24

<210> 493

<211> 22

<212> DNA

<213> Mouse

<400> 493

accaccgcca ctgttctcat ct

22

<210> 494

<211> 20

<212> DNA

<213> Mouse

<400> 494

tgtggggaa gaacatagaa

20

<210> 495

<211> 22

<212> DNA

<213> Mouse

<400> 495

tgatgtgtgg cttgtttctc tt

22

<210> 496

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 496

ataggtgggg aggagctaa

20

&lt;210&gt; 497

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 497

tcatgtgtgg cttgtttctc tt

22

&lt;210&gt; 498

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 498

tgtgcctgtc acagcaactt

20

&lt;210&gt; 499

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 499

catgctagca ccgttagctga

20

&lt;210&gt; 500

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 500

ggagaccccttc ccctccctct

20

&lt;210&gt; 501

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 501

gctgttagttg aagagggcgt

20

<210> 502  
<211> 18  
<212> DNA  
<213> Homo sapiens

<400> 502  
gtgcttggct tcctccag

18

<210> 503  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 503  
caggtcgtagtccatgtcca

20

<210> 504  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 504  
tggagtacga cctgaagctg

20

<210> 505  
<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 505  
actcatcctg gccacaaaaag

20

<210> 506  
<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 506  
gaacaggagg acgctgagg

19

<210> 507  
<211> 20

<212> DNA

<213> Homo sapiens

<400> 507

cgtttgtggc caggatgagt

20

<210> 508

<211> 20

<212> DNA

<213> Homo sapiens

<400> 508

tcacacctacc tggttgtcag

20

<210> 509

<211> 20

<212> DNA

<213> Homo sapiens

<400> 509

gtacgacactg aagctgtggg

20

<210> 510

<211> 27

<212> DNA

<213> Homo sapiens

<400> 510

ggctgagatc acagggttgg gtcactc

27

<210> 511

<211> 27

<212> DNA

<213> Homo sapiens

<400> 511

ccgtgcctgt tggaagttgc ctctgcc

27

<210> 512

<211> 20

<212> DNA

<213> Mouse

&lt;400&gt; 512

aatccccagc aaccactcac

20

&lt;210&gt; 513

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 513

cagacactcc agaagagggc

20

&lt;210&gt; 514

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 514

tgactgctct tccgaaggtt

20

&lt;210&gt; 515

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 515

tttgtggaat agccaaagcc

20

&lt;210&gt; 516

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 516

tctctcctct cttctccccc

20

&lt;210&gt; 517

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 517

agcagggtgc atcaccttat

20

<210> 518  
<211> 20  
<212> DNA  
<213> Mouse

<400> 518  
taggagtgcc ccataggttg

20

<210> 519  
<211> 20  
<212> DNA  
<213> Mouse

<400> 519  
tcattgtacc cagccagtc

20

<210> 520  
<211> 20  
<212> DNA  
<213> Mouse

<400> 520  
aggactgagc ctggatgaga

20

<210> 521  
<211> 20  
<212> DNA  
<213> Mouse

<400> 521  
ctgggcgttt tgttttgttt

20

<210> 522  
<211> 20  
<212> DNA  
<213> Mouse

<400> 522  
cttccttcctg cagctaccac

20

<210> 523  
<211> 20

<212> DNA

<213> Mouse

<400> 523

accctgctac aacgcagact

20

<210> 524

<211> 20

<212> DNA

<213> Mouse

<400> 524

tccaacccttg acacccattt

20

<210> 525

<211> 20

<212> DNA

<213> Mouse

<400> 525

agccagggtc acacagagaa

20

<210> 526

<211> 20

<212> DNA

<213> Mouse

<400> 526

ctgctttcc tcagcaactg

20

<210> 527

<211> 20

<212> DNA

<213> Mouse

<400> 527

attcggcggtt agaagctagg

20

<210> 528

<211> 20

<212> DNA

<213> Mouse

<400> 528  
aactgtacgt ggctgctgg

20

<210> 529  
<211> 20  
<212> DNA  
<213> Mouse

<400> 529  
attcggcgtt agaagctagg

20

<210> 530  
<211> 20  
<212> DNA  
<213> Mouse

<400> 530  
gccaggtgac ccttatgaaa

20

<210> 531  
<211> 20  
<212> DNA  
<213> Mouse

<400> 531  
gagagatggc agacagaggg

20

<210> 532  
<211> 20  
<212> DNA  
<213> Mouse

<400> 532  
agctctctgt ccctggtgaa

20

<210> 533  
<211> 20  
<212> DNA  
<213> Mouse

<400> 533  
tgccaaccac tagcctctct

20

<210> 534

<211> 20

<212> DNA

<213> Mouse

<400> 534

ctgaaccctc cactctcctg

20

<210> 535

<211> 20

<212> DNA

<213> Mouse

<400> 535

agccagggct acacagagaa

20

<210> 536

<211> 20

<212> DNA

<213> Mouse

<400> 536

agccagggct acacagagaa

20

<210> 537

<211> 20

<212> DNA

<213> Mouse

<400> 537

accctgctac aacgcagact

20

<210> 538

<211> 20

<212> DNA

<213> Mouse

<400> 538

gcaagtttca ggagctaggg

20

<210> 539

<211> 20

<212> DNA

<213> Mouse

<400> 539

cccccagaacc agagaccata

20

<210> 540

<211> 20

<212> DNA

<213> Mouse

<400> 540

cccccagaacc agagaccata

20

<210> 541

<211> 20

<212> DNA

<213> Mouse

<400> 541

ctaggggact ctgccaaatg

20

<210> 542

<211> 20

<212> DNA

<213> Mouse

<400> 542

caagacaccc agtcccaact

20

<210> 543

<211> 20

<212> DNA

<213> Mouse

<400> 543

tacttccctt ttcccgaaact

20

<210> 544

<211> 20

<212> DNA

<213> Mouse

<400> 544  
tccttggtgc ttaccctcac

20

<210> 545  
<211> 20  
<212> DNA  
<213> Mouse

<400> 545  
tgttcctgag ttcacaacgc

20

<210> 546  
<211> 20  
<212> DNA  
<213> Mouse

<400> 546  
atccccagca actacatggc

20

<210> 547  
<211> 20  
<212> DNA  
<213> Mouse

<400> 547  
acatgtccac tgtggcaaaa

20

<210> 548  
<211> 20  
<212> DNA  
<213> Mouse

<400> 548  
tgtcatgagt ttgaggccag

20

<210> 549  
<211> 20  
<212> DNA  
<213> Mouse

<400> 549  
atcagacagc ccacaacctc

20

<210> 550  
<211> 20  
<212> DNA  
<213> Mouse

<400> 550  
tatgtgccac cacacctgtc

20

<210> 551  
<211> 20  
<212> DNA  
<213> Mouse

<400> 551  
gctcaaggaa ggacacacac

20

<210> 552  
<211> 22  
<212> DNA  
<213> Mouse

<400> 552  
tgctcttaac atttttagcc at

22

<210> 553  
<211> 20  
<212> DNA  
<213> Mouse

<400> 553  
gctcagcccc tgaatcaata

20

<210> 554  
<211> 20  
<212> DNA  
<213> Mouse

<400> 554  
gggatctgcc tgtcttacca

20

<210> 555  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 555

ggaaggtagg gcctggtaat

20

&lt;210&gt; 556

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 556

gctccaagat ctgtgcgatt

20

&lt;210&gt; 557

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 557

ttagcgtag ggtgagggtg

20

&lt;210&gt; 558

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 558

ggagactacg gacttgtggc

20

&lt;210&gt; 559

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 559

cagttcttccc cgaaaaccac

20

&lt;210&gt; 560

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 560  
tttctggaa ctgagatggc

20

<210> 561  
<211> 20  
<212> DNA  
<213> Mouse

<400> 561  
gttggggctg ctcatagaaa

20

<210> 562  
<211> 20  
<212> DNA  
<213> Mouse

<400> 562  
gctgtggctc tcttggagtt

20

<210> 563  
<211> 20  
<212> DNA  
<213> Mouse

<400> 563  
ctctgatttc ccacatgcct

20

<210> 564  
<211> 20  
<212> DNA  
<213> Mouse

<400> 564  
aagagggagc actgaggaca

20

<210> 565  
<211> 20  
<212> DNA  
<213> Mouse

<400> 565  
cagcagcaaa tgaccttca

20

<210> 566  
<211> 20  
<212> DNA  
<213> Mouse

<400> 566  
gaggcaggca gatttctgag

20

<210> 567  
<211> 20  
<212> DNA  
<213> Mouse

<400> 567  
gttccacatg ttgtggtggc

20

<210> 568  
<211> 20  
<212> DNA  
<213> Mouse

<400> 568  
gggaccttg ggataggatt

20

<210> 569  
<211> 20  
<212> DNA  
<213> Mouse

<400> 569  
tcagacatct ctggccctct

20

<210> 570  
<211> 20  
<212> DNA  
<213> Mouse

<400> 570  
ttcactaagt tgcccaggct

20

<210> 571  
<211> 22

<212> DNA

<213> Mouse

<400> 571

tgccttttc tcacattgtc tc

22

<210> 572

<211> 20

<212> DNA

<213> Mouse

<400> 572

ttagaaggcag aggcagaggc

20

<210> 573

<211> 20

<212> DNA

<213> Mouse

<400> 573

gacccttgga agagcagtcg

20

<210> 574

<211> 20

<212> DNA

<213> Mouse

<400> 574

tggcagctca caatgtcttt

20

<210> 575

<211> 20

<212> DNA

<213> Mouse

<400> 575

ggtgtgggtgt aggggaagaa

20

<210> 576

<211> 22

<212> DNA

<213> Mouse

&lt;400&gt; 576

tttcaactgc aaacacaaaac ag

22

&lt;210&gt; 577

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 577

aggggccaagg aaggagaat

19

&lt;210&gt; 578

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 578

gcaaataatat agggtaaccga gctg

24

&lt;210&gt; 579

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 579

cagattctcc agctgtcagg

20

&lt;210&gt; 580

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 580

ctgtgtttcc gcaccaagt

19

&lt;210&gt; 581

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 581

ctgccccgtcc ttatcttctg

20

<210> 582  
<211> 20  
<212> DNA  
<213> Mouse

<400> 582  
acgcacgctc actcatacac

20

<210> 583  
<211> 20  
<212> DNA  
<213> Mouse

<400> 583  
cagcagaggt gatgggttct

20

<210> 584  
<211> 22  
<212> DNA  
<213> Mouse

<400> 584  
ttgtcacaca gtggtaaat gc

22

<210> 585  
<211> 20  
<212> DNA  
<213> Mouse

<400> 585  
tagaaccgtg gctgaggact

20

<210> 586  
<211> 24  
<212> DNA  
<213> Mouse

<400> 586  
ccgtaagata tgaaagaact tgga

24

<210> 587  
<211> 20

<212> DNA

<213> Mouse

<400> 587

taatcctggc ttagcgcttg

20

<210> 588

<211> 20

<212> DNA

<213> Mouse

<400> 588

tagaaagcac aggggacagg

20

<210> 589

<211> 20

<212> DNA

<213> Mouse

<400> 589

ccttcctcgt ctgagctgtt

20

<210> 590

<211> 20

<212> DNA

<213> Mouse

<400> 590

ttgggacgtg acctgagaat

20

<210> 591

<211> 20

<212> DNA

<213> Mouse

<400> 591

tatgtgtctg gccgttgttc

20

<210> 592

<211> 19

<212> DNA

<213> Mouse

&lt;400&gt; 592

gatgtgggtg caggtgaag

19

&lt;210&gt; 593

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 593

ccctttctgg agtgtctgaa

20

&lt;210&gt; 594

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 594

tctaggcagg gctacccccc t

21

&lt;210&gt; 595

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 595

gctgagcagc ctctagcaa

19

&lt;210&gt; 596

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 596

accatggctt ttccccagtaa

20

&lt;210&gt; 597

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 597

ctgtgccttt ggtgatcaga

20

<210> 598  
<211> 20  
<212> DNA  
<213> Mouse

<400> 598  
tgtggcactc tacggcataa

20

<210> 599  
<211> 23  
<212> DNA  
<213> Mouse

<400> 599  
tgcatcacta ttaaggctca acc

23

<210> 600  
<211> 23  
<212> DNA  
<213> Mouse

<400> 600  
aagaatttgc aaagactgtg aga

23

<210> 601  
<211> 20  
<212> DNA  
<213> Mouse

<400> 601  
ctggaccttt ggaagagcag

20

<210> 602  
<211> 20  
<212> DNA  
<213> Mouse

<400> 602  
ggtggctcaa accatccata

20

<210> 603  
<211> 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 603

gagggcaatg agcaaaaatgt

20

&lt;210&gt; 604

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 604

ggtcctgtct ctggttcagg

20

&lt;210&gt; 605

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 605

taacacccac atcaggcaac

20

&lt;210&gt; 606

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 606

tttcatttcc tggtgttcct tt

22

&lt;210&gt; 607

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 607

aaacacaggc ggaacgatag

20

&lt;210&gt; 608

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Mouse

<400> 608  
ctatcggtcc gcctgtgttt

20

<210> 609  
<211> 21  
<212> DNA  
<213> Mouse

<400> 609  
aaggaagagg atggagaaaag a

21

<210> 610  
<211> 20  
<212> DNA  
<213> Mouse

<400> 610  
cggttcttaa tggagcagag

20

<210> 611  
<211> 20  
<212> DNA  
<213> Mouse

<400> 611  
tcctccccag ttacctagca

20

<210> 612  
<211> 19  
<212> DNA  
<213> Mouse

<400> 612  
cagcaggcaa gatgacctc

19

<210> 613  
<211> 20  
<212> DNA  
<213> Mouse

<400> 613  
gtccctcacc agccatgtta

20

<210> 614  
<211> 20  
<212> DNA  
<213> Mouse

<400> 614  
agcctgggct aagttgtgtg

20

<210> 615  
<211> 20  
<212> DNA  
<213> Mouse

<400> 615  
tatgggccaa tgggtttcct

20

<210> 616  
<211> 20  
<212> DNA  
<213> Mouse

<400> 616  
atgggtggctc acaaccatct

20

<210> 617  
<211> 20  
<212> DNA  
<213> Mouse

<400> 617  
ttgtcctctg attgcagcat

20

<210> 618  
<211> 20  
<212> DNA  
<213> Mouse

<400> 618  
cttgggtcat caggctttgt

20

<210> 619  
<211> 20

<212> DNA

<213> Mouse

<400> 619

aagctgccct gctctctcta,

20

<210> 620

<211> 20

<212> DNA

<213> Mouse

<400> 620

atgctcagcc tgctttgttt

20

<210> 621

<211> 20

<212> DNA

<213> Mouse

<400> 621

gctgatagcc ctggggttcta

20

<210> 622

<211> 21

<212> DNA

<213> Mouse

<400> 622

tgtacgcaca aattgacttg c

21

<210> 623

<211> 21

<212> DNA

<213> Mouse

<400> 623

gaatccacat tgcaaaggct a

21

<210> 624

<211> 20

<212> DNA

<213> Mouse

<400> 624  
cacaggcaaa tgaagggaag

20

<210> 625  
<211> 20  
<212> DNA  
<213> Mouse

<400> 625  
ccagacttct ccagctctcc

20

<210> 626  
<211> 21  
<212> DNA  
<213> Mouse

<400> 626  
tcctcgagag gctcttaggtt t

21

<210> 627  
<211> 20  
<212> DNA  
<213> Mouse

<400> 627  
tgccctagtc accacaggag

20

<210> 628  
<211> 21  
<212> DNA  
<213> Mouse

<400> 628  
cctgtggttt actaggcaga a

21

<210> 629  
<211> 20  
<212> DNA  
<213> Mouse

<400> 629  
gcctgatagc ctggaataca

20

<210> 630  
<211> 20  
<212> DNA  
<213> Mouse

<400> 630  
aaaggatgt gtggcgtaag

20

<210> 631  
<211> 20  
<212> DNA  
<213> Mouse

<400> 631  
caaaacccaa ccttctcagc

20

<210> 632  
<211> 20  
<212> DNA  
<213> Mouse

<400> 632  
tgcaactgacc gtgatagagg

20

<210> 633  
<211> 20  
<212> DNA  
<213> Mouse

<400> 633  
cggtgttagct ctggctgtct

20

<210> 634  
<211> 20  
<212> DNA  
<213> Mouse

<400> 634  
catctcacca actcgcaatt

20

<210> 635  
<211> 21

<212> DNA

<213> Mouse

<400> 635

tttctggaa caaagaggct a

21

<210> 636

<211> 20

<212> DNA

<213> Mouse

<400> 636

gaacccaagt gttgggtaa

20

<210> 637

<211> 20

<212> DNA

<213> Mouse

<400> 637

tggaagccca tctgtctctt

20

<210> 638

<211> 20

<212> DNA

<213> Mouse

<400> 638

aaatgcaagt gggtgcttct

20

<210> 639

<211> 19

<212> DNA

<213> Mouse

<400> 639

ccagaagagg gcgtcagat

19

<210> 640

<211> 20

<212> DNA

<213> Mouse

<400> 640  
ggtgtgcacc accatattca

20

<210> 641  
<211> 21  
<212> DNA  
<213> Mouse

<400> 641  
gggaattatc agccaaaaag c

21

<210> 642  
<211> 20  
<212> DNA  
<213> Mouse

<400> 642  
gcccaactga aagctcaact

20

<210> 643  
<211> 21  
<212> DNA  
<213> Mouse

<400> 643  
ggaaggggga taacaattga a

21

<210> 644  
<211> 23  
<212> DNA  
<213> Mouse

<400> 644  
tgctaatttc aagcacagtg aga

23

<210> 645  
<211> 20  
<212> DNA  
<213> Mouse

<400> 645  
agcttgacac cttgacagca

20

<210> 646

<211> 20

<212> DNA

<213> Mouse

<400> 646

aacctgcaga gaggagacca

20

<210> 647

<211> 20

<212> DNA

<213> Mouse

<400> 647

ctccaagggg aggactcatt

20

<210> 648

<211> 24

<212> DNA

<213> Mouse

<400> 648

ttcaatttag tttctctctt ctga

24

<210> 649

<211> 20

<212> DNA

<213> Mouse

<400> 649

tgcaggacca agaagttaggc

20

<210> 650

<211> 20

<212> DNA

<213> Mouse

<400> 650

cgagatctga tgcccttcttc

20

<210> 651

<211> 20

<212> DNA

<213> Mouse

<400> 651

tgctgagagc agaaaaggaa

20

<210> 652

<211> 166

<212> DNA

<213> Mouse

<400> 652

gcagtgagct gcagagtttgcagaatgagg gcactctaaa ctcataatg gaggaggccc 60  
ttccctcaca ctccagatgg ctgataggtg gcattacatg gtccancgcg cgcacgcgt 120  
cagatgcaat ctccacattc ataaccagat gtccttgggtt aggct 166